

OR 1630

Vitamin A deficiency is associated with body mass index and body adiposity in women with recommended intake of vitamin A

La deficiencia de vitamina A se asocia con el índice de masa corporal y la adiposidad corporal en mujeres con ingesta recomendada de vitamina A

Claudia Bento¹, Andréa C. Matos², Adryana Cordeiro³ and Andréa Ramalho⁴

¹Human Nutrition. University Federal of Rio de Janeiro (UFRJ). Rio de Janeiro, Brazil.

²Medical Clinic. University Federal of Rio de Janeiro (UFRJ). Department of Nutrition and Dietetics. University Federal Fluminense. Micronutrients Research Center (NPqM) of Institute of Nutrition Josué de Castro (INJC) Rio de Janeiro, Brazil. ³Faculty of Medicine. University Federal of Rio de Janeiro (UFRJ). Micronutrients Research Center (NPqM)/Federal University of Rio de Janeiro. Rio de Janeiro, Brazil. ⁴Public Health. Fundação Oswaldo Cruz (FIOCRUZ). Rio de Janeiro, Brazil. Social Applied Nutrition Department of UFRJ. Micronutrients Research Center (NPqM) of Institute of Nutrition Josué de Castro (INJC). Rio de Janeiro, Brazil

Received: 10/10/2017

Accepted: 02/06/2018

Correspondence: Adryana Cordeiro. Department of Social Applied Nutrition - Micronutrients Research Center (NPqM). Federal University of Rio de Janeiro. Av. Brigadeiro Trompovsky, s/n. Sub-solo, Bloco J. 21941-590 Ilha do Fundão, Rio de Janeiro. Brazil

e-mail: adrynutri@yahoo.com.br

DOI: 10.20960/nh.1630

ABSTRACT

Introduction: evidence indicates that vitamin A is involved in regulating fat mass. A low consumption of vitamin A has been reported in individuals with obesity, as have lower

concentrations of this vitamin, than in eutrophic individuals when their dietary intake of vitamin A is not significantly different.

Objective: to investigate vitamin A nutritional status and its association with body mass index (BMI) and body fat in women who have the recommended dietary intake of vitamin A.

Methods: cross-sectional study with 200 women, paired by age and by the dietary intake of vitamin A recommended. Participants were divided into four groups, according to BMI. Anthropometric data were evaluated (weight, BMI and waist circumference [WC]), as well as the diagnosis of night blindness (NB). Lipid and glycemic profiles were measured. The cut-off points for deficiency of serum concentrations of retinol and β -carotene were $< 1.05 \mu\text{mol/l}$ and $40 \mu\text{g/dl}$, respectively. The recommended dietary intake of vitamin A was $700 \mu\text{g/day}$.

Results: there was a significant drop in retinol concentrations according to BMI ($p < 0.001$) and WC ($p < 0.001$). We found β -carotene to behave similarly ($p = 0.005$; $p < 0.001$). We found NB in 7.5% of overweight (OW) cases and 20.0% of obesity class II (OII), and no functional alteration was found in the eutrophic group (EU). Inadequate levels of retinol and β -carotene increased the odds ratio for the occurrence of OW, obesity class I (OI) and OII, as well as inadequate WC.

Conclusion: even with recommended intake of vitamin A, we found a biochemical and functional inadequacy of vitamin A nutritional status, associated with overweight, obesity and body adiposity.

Key words: Overweight. Obesity. Retinol. β -carotene. Night blindness. Body adiposity.

RESUMEN

Introducción: la evidencia indica que la vitamina A está involucrada en la regulación de la masa grasa. Un bajo consumo de vitamina A ha sido reportado en individuos con obesidad, ya que tienen concentraciones más bajas de esta vitamina que los individuos eutróficos a pesar de que su ingesta dietética de vitamina A no es significativamente diferente.

Objetivo: investigar el estado nutricional de la vitamina A y su asociación con el índice de masa corporal (IMC) y la grasa corporal en las mujeres que tienen la ingesta dietética recomendada de vitamina A.

Métodos: estudio transversal con 200 mujeres, emparejado por edad y por la ingesta dietética de vitamina A recomendada. Se dividieron en cuatro grupos, de acuerdo con el IMC. Los datos antropométricos fueron evaluados (peso, índice de masa corporal [IMC] y circunferencia de la cintura [CC]), así como el diagnóstico de ceguera nocturna (CN). Se midieron los perfiles lipídicos y glicémicos. Los puntos de corte para la deficiencia de las concentraciones séricas de retinol y β -caroteno fueron $< 1,05 \mu\text{mol/l}$ y $40 \mu\text{g/dl}$, respectivamente. La ingesta dietética recomendada de vitamina A fue de $700 \mu\text{g/día}$.

Resultados: hubo una disminución significativa de las concentraciones de retinol según el IMC ($p < 0,001$) y CC ($p < 0,001$). Se observó un comportamiento similar del β -caroteno ($p = 0,005$, $p < 0,001$). Encontramos NB en el 7,5% de los casos con sobrepeso (OW) y el 20,0% de los casos con obesidad clase II (OII), y no encontramos alteración funcional en la UE. Niveles inadecuados de retinol y β -caroteno aumentaron la *odds ratio* para la ocurrencia de OW, obesidad clase I (OI) y OII, al igual que los niveles insuficientes de CC.

Conclusión: incluso con la ingesta recomendada de vitamina A, encontramos una deficiencia bioquímica y funcional del estado nutricional de vitamina A asociada al sobrepeso, la obesidad y la adiposidad corporal.

Palabras clave: Sobrepeso. Obesidad. Retinol. β -caroteno. Ceguera nocturna. Adiposidad corporal.

INTRODUCTION

Obesity is a disease of rising prevalence now considered as one of the greatest public health problems in modern society. It is estimated that 65% of the world's population lives in countries where overweight and obesity kill more people than malnutrition (1). Over the last 20 years, the prevalence of female obesity has risen from 8% to 16.9% (2). Of total primary healthcare users, 89.7% are women, and 30.6% of those women

are overweight, while 19.5% are obese, testament to the enormity of the socioeconomic impact that obesity has in Brazil (3,4).

Obesity creates a predisposition to chronic, non-communicable diseases, which has a major impact on the health of the working-age population. There are well-established associations between a high BMI and increased morbidity, including from cardiovascular disease, type 2 diabetes mellitus, high blood pressure, stroke, and dyslipidemia (5). The key point is that obesity associated with insulin resistance is responsible for mechanisms that trigger the lipid abnormalities observed in individuals with obesity. These alterations may be responsible for increasing the risk of cardiovascular disease and also contribute to increasing oxidative stress in obesity (6).

There is evidence that vitamin A is involved in the regulation of body fat, which suggests that when vitamin A is deficient (VAD) there occurs an increase in the recruitment of preadipocytes into adipocytes, inhibition of apoptosis, and an increase of adaptive thermogenesis (7,8). Additionally, there are studies supporting that the increase of BMI can be associated with lowest concentrations of serum vitamin A, which suggests that VAD may play a part in the growth of obesity (9,10).

Using anthropometric indicators as predictors of antioxidant insufficiency, we found BMI and waist circumference to have an inverse association with retinol and β -carotene in humans (11). In some studies with adult participants, serum β -carotene was found to have a negative association with BMI, body fat percentage, waist circumference and waist-hip ratio (12,13).

There is also research assuming increased BMI to relate directly to oxidative stress, given that the greater the degree of obesity, the more intense the inflammatory condition to which these individuals are exposed, due to the intra-abdominal adipose tissue increasing production of proinflammatory cytokines, which leads to an increase in oxidative stress and consequently, to greater demand for nutrients with antioxidant action, such as vitamin A (14,15). However, there is a dearth of research comparing serum retinol and carotenoid concentrations and standards for vitamin A intake and their relation to BMI and body adiposity. Some studies have demonstrated lower serum retinol and carotenoid concentrations in individuals with obesity than in eutrophic individuals when their dietary intake of sources of these nutrients, as reported in food frequency questionnaires, is not significantly different (16,17).

The aim of this study was to analyze vitamin A nutritional status through biochemical and functional indicators and its association with anthropometric parameters in women from different BMI ranges with recommended intake of vitamin A.

MATERIAL AND METHODS

This is an analytical cross-sectional study with adult women, scheduled for nutritional care in a Nutrition Ambulatory in the referred Municipal Health Center of the city of Rio de Janeiro, in the period from January 2012 to October 2014. All women who attended inclusion criteria were invited to participate in the study. The research was previously approved by the Research Ethics Committee of the Hospital Universitário Clementino Fraga Filho of the Federal University of Rio de Janeiro (protocol of research no. 011/05).

After the initial approach and clarifications about the goals and procedures of the study, all the women who agreed to participate in the study gave a formal authorization by signing the informed consent, thus joining the first stage of the research that comprised an anthropometric assessment and a food intake survey.

The inclusion criteria were: female gender (aged 20-59 years), from all ethnic groups and social classes, scheduled for nutritional care of Health Unit. The following exclusion criteria were applied in the first phase of the research: inadequacy of vitamin A intake (evaluated in the first phase of research), pregnant women and/or nursing mothers, liver diseases (except for non-alcoholic liver disease), malabsorptive syndromes, acute infection, alcoholism, kidney insufficiency, HIV virus carriers, cancer, and use of vitamin A supplements over the last six months.

After the first phase, the sample was divided into four groups according to BMI ranges (18): eutrophic group (EU), overweight group (OW), obesity class I group (OI) and obesity class II (OII) group.

Sample size

A sample size calculation was performed on the basis of the study. It estimated the prevalence of Brazilian adults inadequacy intake of micronutrients (POF 2008-2009) that demonstrated elevated inadequacy of vitamin A intake in women about 78,5% in

urban and rural areas studied (19). Consequently, the prevalence of adequacy was 21.5%.

To obtain a sample size with 95% of confidence interval, considering an adequacy prevalence of 21.5% with a sampling error of 6%, 155 women would be needed with a recommended dietary intake of vitamin A, according to the Institute of Medicine at the end of research. Based on this calculation, the number of attendance evaluated in a period of 18 months, considering the average weekly attendance in the six months prior to the accomplishment of the present research, was 1,080 women (Fig. 1).

Anthropometric measures

In this stage of the research, weight and height of the women participating in the study were measured and BMI was calculated according to the cut-off points recommended by the World Health Organization (WHO) (18). Additionally, waist circumference (WC) was measured, using inadequate the cut-off > 80 cm for women, as proposed by the International Diabetes Federation (IDF) (20).

Food frequency questionnaire for vitamin A

Women involved in the study were asked separately to recall all the dishes, snacks, or other foods eaten in the 24 hours prior to the survey, regardless of whether the food was eaten inside or outside the house. During data collection, each woman was prompted to make sure that no meal or snack was forgotten. Next, a detailed list of all the ingredients of the dishes, snacks or other foods mentioned in the reported consumed food were collected from each woman via interview. The study participants were then asked to recall how many days they had consumed each of the 15 locally available vitamin A rich foods in the past seven days (21). During training of data collectors, it was stressed that small quantities of food eaten less than one tablespoon should be excluded. This was important as foods eaten below the aforementioned quantities would not contribute significantly to nutrient adequacy but would inflate the score. Minimum consumption of one tablespoon of food was better correlated with probability of adequacy. Food frequency questions were double coded and entered into a custom built spreadsheet, which calculated an overall mean daily intake of vitamin A using the vitamin A content of foods as published in the table of the

Instituto de Nutrición de Centro América y Panamá (INCAP) (22). Portion size was assessed using a Photographic Atlas of Food Portion Sizes (21).

Vitamin A intake was compared to the values of the daily intake recommended by the Institute of Medicine in 2001. The cut-off point adopted for the recommended dietary intake of vitamin A was 700 µg/day.

A month after the first procedure, all the women participating in the first stage of the study were again scheduled to receive information about the nutritional diagnosis, including intake of vitamin A, in addition to receiving the relevant dietary guidelines. Women who had reached the recommended dietary intake of this vitamin were invited to follow up the previously established research protocol and, to that aim, they were instructed to appear for the blood collection procedure after a 12-hour fast.

In the second phase of the research, only women who showed the recommended dietary intake of vitamin A were included in accordance with the results of the food intake survey conducted in the previous phase.

Biochemical measures of vitamin A and diagnosis of night blindness (NB)

The nutritional status of vitamin A was assessed according to biochemical and functional parameters. For biochemical assessment of vitamin A, serum concentrations of retinol and β-carotene were quantified by high-performance liquid chromatography with ultraviolet detector (HPLC-UV), and the following cut-off points were used: < 1.05 µmol/l and 40 µg/dl to indicate inadequacy of retinol and β-carotene, respectively (23,24).

In the functional assessment of the nutritional status of vitamin A, the presence of NB was investigated through an interview standardized by the WHO (25).

The interview was composed of the following questions:

1. Do you have difficulty to see during the day?
2. Do you have difficulty to see in low light or at night?
3. Do you have NB?

Cases of NB were considered when the answer to question 1 was “No” and, at least, one answer to questions 2 or 3 was “Yes”. If the interviewed had any ophthalmic problem corrected by glasses or contact lens, he/she was asked about the ability to see with them.

Diagnosis of DM2

The diagnosis of DM2 took place in accordance with the American Diabetes Association (ADA) (26) guidelines previously endorsed by the Brazilian Society of Diabetes, which classifies the levels of blood glucose into ≥ 126 mg/dl (fasting) or > 200 mg/dl (one-off).

Others biochemical measures

For biochemical evaluation, a blood sample was obtained by venipuncture after a 12-hour fast period. Laboratory tests were conducted to evaluate: lipid profile (total cholesterol, triglycerides, high-density lipoprotein cholesterol [HDL-c], low density lipoprotein cholesterol [LDL-c]) and the following concentrations were considered as normal values: total cholesterol < 200 mg/dl; triglycerides < 150 mg/dl; HDL-c > 50 mg/dl; LDL-c < 150 mg/dl. Determinations of triglycerides, total cholesterol and HDL-c were performed by the enzymatic colorimetric method. Reagents for these biochemical evaluations were purchased from Labtest Diagnóstica S.A., Minas Gerais, Brazil. LDL-c fraction was determined in accordance with the Friedewald's formula.

Statistical analysis

In the statistical analysis, the measures of central tendency and dispersion were calculated and expressed as means and standard deviations for clinical, dietary and biochemical variables. The analysis of variance (ANOVA) and the Bonferroni test for multiple comparisons were used. Pearson correlation coefficient was applied for serum concentrations of retinol and β -carotene with variables related to markers of oxidative stress. The odds ratio was estimated to evaluate the strength of the association between serum concentrations of retinol and β -carotene with biochemical, clinical and anthropometric variables, either the association of inadequacy of serum concentrations of both micronutrients according to BMI ranges. The significance level adopted was 5% ($p < 0.05$). Statistical analysis was performed using the Statistical Package for the Social Sciences (SPSS) for Windows version 17.

RESULTS

The sample group comprised 200 adult women, who were divided into four groups according to their BMI. Their characteristics are described in table I.

A statistically significant difference between the classes was observed when considering the variable BMI (EU/OW $p = 0.04$; EU/OI $p = 0.02$; EU/OII $p = 0.03$; OW/OI $p = 0.02$; OW/OII $p < 0.00$; OI/OII $p = 0.01$).

Even with the recommended daily intake of vitamin A, the percentage of vitamin A (retinol and β -carotene) inadequacy was, respectively, 7.5 and 8.8% in the EU, 12.5 and 17.5% in the OW, 17.5 and 25% in the OI and 22.5 and 27.5% in the OII.

When comparing the groups according to the occurrence of NB, a significant difference between them was found ($p = 0.002$). The breakdown for NB was as follows: 7.5% in overweight women and 12.5% and 20% in women with obesity class I and class 2, respectively. None noted in the eutrophic group of women. The number of women with NB increased in accordance with BMI: three women in the overweight group, seven in the OI group, and ten in the OII group.

The mean of serum retinol and β -carotene concentrations of the groups is presented in table II.

Overweight women were found to be 14.0, 5.1 and 3.5 times more likely to have inadequate retinol, β -carotene concentrations and NB, respectively. The odds ratio increased to 35.3, 17.0 and 7.4, respectively, in OI, and to 65.4, 26.7 and 9.2, respectively, in OII. Women in the overweight, OI and OII groups were 5.1, 7.2 and 8.0 times more likely to have an inadequate WC, respectively (Table III).

A negative correlation was found between the serum retinol and β -carotene concentrations with inadequate WC in the OW groups ($r = -0.32/-0.46$; $p < 0.001$), OI ($r = -0.56/-0.70$; $p < 0.001$) and OII ($r = -0.68/-0.85$; $p < 0.001$), respectively.

A strong correlation was found between serum retinol and β -carotene concentrations in all the groups studied. Proportionately to the increase in BMI, the negative correlation with serum retinol and β -carotene levels increased (Table IV).

DISCUSSION

Our research showed an inverse relationship between an increase in BMI and alterations in vitamin A nutritional status, as identified in the women assessed to have lower levels of serum retinol and β -carotene, as well as through the diagnosis of NB.

Researchers suggest an inverse association between intake and serum concentrations of antioxidant nutrients, such as vitamin A, in the presence of obesity (27,28). Our findings showed that the biochemical and functional vitamin A profiles of overweight and women with obesity are in a much worse state than those of eutrophic women, even if their vitamin A intake falls within the recommended dietary guidelines. The findings draw attention to an important facet of VAD on which to reflect, by demonstrating that even while ingesting the recommended daily vitamin A allowance, the increased demand for the nutrient was found to relate to an increase in BMI and body adiposity, and in the midst of such weight increases, nutritional vitamin A requirements can be much higher than current recommended allowances.

We found serum retinol and β -carotene concentrations to be significantly higher in eutrophic women compared to women with overweight, obesity class I and obesity class II. In addition, the odds ratio were more elevated for serum retinol and β -carotene deficiency, night blindness and inadequate waist circumference in women with obesity class II. There is a research that compared eutrophic women with those with obesity and the last one had lower concentrations of β -carotene in the blood, a higher concentration of endothelial adhesion molecules and greater concentrations of blood sugar and insulin, confirming that a rise in body fat leads to increased resistance to insulin, endothelial dysfunction and oxidative stress, and these relate to the processes involved in obesity (29,30). Retinol binding protein 4 (RBP4), which has been referred to as an adipokine that is positively associated with BMI, when analyzed in overweight women confirmed prior research that serum RBP4 is higher relative to "normal" levels in lean adults (31). Relatively high serum RBP4 in overweight subjects despite lower retinol levels suggested that some of the serum RBP4 in overweight individuals may be functioning unbound to retinol. Elevated RBP4 with excess body fat has been associated with insulin resistance mediated through suppressed GLUT4 transporter expression (32).

Otherwise, in a study developed by Nuss et al. (2017) (33), β -carotene was not significantly associated with any indicator of body composition (BMI, fat mass and lean mass) in a group of women. Although body composition influences serum nutrient profiles, parameters related to high body-fat mass, such as oxidative stress and inflammation may influence such changes in nutrient levels (33).

There is a study reporting low vitamin A intake as a factor causing VAD in overweight individuals (28). Indeed, inadequate vitamin A consumption was noted in 70% of the 738 women assessed during the first phase of our research, and these findings are in line with the most recent nationwide dietary survey, which revealed a 78.0% magnitude of inadequate vitamin A intake across all the regions studied, both in urban and rural Brazil. Such findings are a cause for concern, given the important role this nutrient plays in cellular metabolism.

We found that the percentages of β -carotene insufficiency were higher than those of retinol, and we also noted a moderate positive correlation between serum retinol and β -carotene concentrations in EU, and a negative strong one in the other groups (OW, OI and OII). Our findings corroborate those of Mecocci et al. (2008) (34), as they show a drop in serum β -carotene concentrations and, at the same time, a drop in circulating retinol values, suggesting more β -carotene is being mobilized for conversion to retinol. These findings may be related to the greater mobilization of β -carotene for conversion into retinol, since β -carotene is known to be a potent precursor to retinol. This finding is noteworthy given how a drop in serum β -carotene concentrations leads to a rise in oxidative stress (34), suggesting that this antioxidant could be being rerouted to other important functions, like combatting oxidative stress, in order to maintain an adequate retinol nutritional status. It is thus important to maintain serum retinol concentrations in order to keep β -carotene functioning as an antioxidant.

There are also studies showing that a buildup of visceral fat and increase in BMI are linked to an increase in lipid peroxidation and decrease in serum retinol and β -carotene concentrations, and thus accompanies a negative antioxidant system (35,36). Furthermore, existing research into vitamin A consumption shows, in such circumstances, vitamin intake to be below the recommended daily allowances (37-39). The main limitation of this study is related to the assessment of dietary intake, the same limitations presented by all studies based on self-reported dietary recall, in particular, underestimation of consumption. Nevertheless, we see no reason to believe that underreporting could have been different between the groups we assessed.

This study demonstrated that even with the recommended intake of vitamin A, a biochemical and functional inadequacy of vitamin A nutritional status associated with overweight, obesity and body adiposity was found. The findings support the

hypothesis that vitamin A deficiency in these individuals could arise from the higher metabolism of this nutrient, to which they are more exposed than eutrophic individuals. Demonstrating that these conditions may represent an important cause of vitamin A depletion, as well as aggravating factors for vitamin A deficiency.

ACKNOWLEDGMENTS

We thank the National Council of Technological for their research support to this study.

AUTHORSHIP

Bento C, Matos AC, Cordeiro A participated in the conception and design of the study and data analysis, and performed the statistical analysis. Matos AC, Cordeiro A, Ramalho A drafted the manuscript and performed the interpretation of data. Ramalho A coordinated the manuscript and gave final approval.

REFERENCES

1. World Health Organization (WHO). 10 Facts about obesity. Consulted May 2014. Accessed Feb 24th 2017. Available from: <http://www.who.int/features/factfiles/obesity/facts/en/>
2. Instituto Brasileiro de Geografia e Estatística. Pesquisa de Orçamentos Familiares. POF 2008-2009. Accessed March 22nd 2017. Available from: http://www.ibge.gov.br/home/presidencia/noticias/noticia_visualiza.php?id_noticia=1699&id_pagina=1
3. Ministério da Saúde. Indicadores e Dados Básicos do Brasil – DATASUS 2009. Accessed Dec 22nd 2016. Available from: <http://www.datasus.gov.br/db>
4. Ministério da Saúde. Sistema de Vigilância Alimentar e Nutricional – SISVAN 2012. Accessed March 23rd 2013. Available from: <http://www.saude.gov.br/nutricao>
5. Campbell PT. Obesity: a certain and avoidable cause of câncer. *Lancet* 2014;384(9945):727-8.
6. Godala M, Materek-Kuśmierkiewicz I, Moczulski D, et al. The risk of plasma vitamin A, C, E and D deficiency in patients with metabolic syndrome: a case-control study. *Adv Clin Exp Med* 2017;26(4):581-6.

7. Schreiber R, Taschler U, Preiss-Landl K, et al. Retinyl ester hydrolases and their roles in vitamin A homeostasis. *Biochim Biophys Acta* 2012;23:113-23.
8. Pang XY, Wang S, Jurczak MJ, et al. Retinol saturase modulates lipid metabolism and the production of reactive oxygen species. *Arch Biochem Biophys* 2017;633:93-102.
9. Rumana Yasmeen R, Shanmugam M, Jeyakumar A, et al. The contribution of vitamin A to autocrine regulation of fat depots. *Biochim Biophys Acta* 2012;1821(1):190-7.
10. Vincenta H, Bourguignonb C, Weltmanc A, et al. Effects of antioxidant supplementation on insulin sensitivity, endothelial adhesion molecules, and oxidative stress in normal-weight and overweight young adults. *Metabolism* 2009;58(2):254-62.
11. Stenzel AP, Carvalho R, Jesus P, et al. Serum antioxidant associations with metabolic characteristics in metabolically healthy and unhealthy adolescents with severe obesity: an observational study. *Nutrients* 2018;10:150.
12. Canas J, Damaso L, Altomare J, et al. Insulin resistance and adiposity in relation to serum b-carotene levels. *J Pediatr* 2012;161(1):58-64.
13. Martínez-Tomás R, Pérez-Llamas F. Daily intake of fruit and vegetable soups processed in different ways increases human serum b-carotene and lycopene concentrations and reduces levels of several oxidative stress markers in healthy subjects. *Food Chem* 2012;127-33.
14. Syauqy A, Hsu CY, Rau HH, et al. Association of dietary patterns with components of metabolic syndrome and inflammation among middle-aged and older adults with metabolic syndrome in Taiwan. *Nutrients* 2018;10(2):143.
15. Bressan J, Hermsdorff L, Zulet A, et al. Impacto hormonal e inflamatório de diferentes composições dietéticas: ênfase em padrões alimentares e fatores dietéticos específicos. *Arq Bras Endocrinol Metab* 2009;572-81.
16. Nyssonen K. Lycopene, lutein and B-carotene as determinants of LDL conjugated dienes in serum. *Atherosclerosis* 2010;200:565-72.
17. Wei X, Peng R, Cao J, et al. Serum vitamin A status is associated with obesity and the metabolic syndrome among school-age children in Chongqing, China. *Asia Pac J Clin Nutr* 2016;25(3):563-70.

18. World Health Organization (WHO). Physical status: the use and interpretation of anthropometry: report of a WHO expert committee. Geneva: WHO; 1995.
19. Araujo MC, Verly Junior E, Junger WL, et al. Independent associations of income and education with nutrient intakes in Brazilian adults: 2008-2009 National Dietary Survey. *Public Health Nutr* 2014;17(12):2740-52.
20. International Diabetes Federation. The Consensus Worldwide Definition of the Metabolic Syndrome, 2006. Accessed June 20th 2017. Available from: http://www.idf.org/webdata/docs/MetS_def_update2006.pdf
21. Pinheiro ABV, Lacerda EM, Benzecry EH, et al. Tabela para avaliação de consumo alimentar em medidas caseiras. Vol. 5; 2001. p. 352.
22. Instituto de Nutrición Centro América y Panamá. Tabla de composición de alimentos para uso en América Latina, 2006. Accessed May 19th 2012. Available from: <http://www.incap.org.gt>
23. World Health Organization (WHO). Indicators for assessing vitamin A deficiency and their application in monitoring and evaluating intervention programmes. Micronutrient Series, WHO/NUT. 10. Geneva: WHO; 1996.
24. IVACG (International Vitamin A Consultative Group). Improving the vitamin A status of populations. USAID-ILSI. 2003.
25. World Health Organization (WHO). Indicators for assessing vitamin A deficiency and their application in monitoring and evaluating intervention programs. WHO/NUT/96.10. Geneva: WHO; 1996.
26. American Diabetes Association (ADA). Classification and diagnosis of diabetes: standards of medical care in diabetes – 2018. *Diabetes Care* 2018;41(Suppl 1):S13- S27.
27. Kabat GC, Heo M, Ochs-Balcom HM, et al. Longitudinal association of measures of adiposity with serum antioxidant concentrations in postmenopausal women. *Eur J Clin Nutr* 2016;70(1):47-53.
28. Lefebvre P, Letois F, Sultan A, et al. Nutrient deficiencies in patients with obesity considering bariatric surgery: a cross-sectional study. *Surg Obes Relat Dis* 2014;10(3):540-6.
29. França BK. Peroxidação lipídica e obesidade: métodos para aferição do estresse oxidativo em obesos. *GE J Port Gastreterol* 2013;20(5)199-206.

30. Barbosa K, Bressan J, Zulet M, et al. Influence of dietary intake on plasma biomarkers of oxidative stress in humans. *An Sist Sanit Navar* 2008;31(3):259-80.
31. Comucci EB, Vasques AC, Geloneze B, et al. Serum levels of retinol binding protein 4 in women with different levels of adiposity and glucose tolerance. *Arq Bras Endocrinol Metabol* 2014;58(7):709-14.
32. Smith U, Kahn BB. Adipose tissue regulates insulin sensitivity: role of adipogenesis, de novo lipogenesis and novel lipids. *J Intern Med* 2016;280(5):465-75.
33. Nuss ET, Valentine AR, Zhang Z, et al. Serum carotenoid interactions in premenopausal women reveal B-carotene is negatively impacted by body fat. *Exp Biol Med* 2017;242(12):1262-70.
34. Paganini-Hill A, Kawas CH, Corrada MM. Antioxidant vitamin intake and mortality: The Leisure World Cohort Study. *Am J Epidemiol* 2015;181(2):120-6.
35. Burri B, Neidlinger T, Clifford A. Serum carotenoid depletion follows first-order kinetics in healthy adult women fed naturally low carotenoid diets. *J Nutr* 2001;131(8):2096-100.
36. Shaker O, Shehaby AA, Zakaria A, et al. Plasma visfatin and retinol binding protein-4 levels in patients with type 2 diabetes mellitus and their relationship to adiposity and fatty liver. *Clin Biochem* 2011;1457-63.
37. Matos AC, Souza GG, Ramalho A, et al. Effect of vitamin A supplementation on clinical evolution in patients undergoing coronary artery bypass grafting, according to serum levels of zinc. *Nutr Hosp* 2012;27(6):1981-6.
38. Berry D, Noy N. Signaling by vitamin A and retinol-binding protein in regulation of insulin responses and lipid homeostasis. *Biochim Biophys Acta* 2012;1821(1):168-76.
39. Wang L, Gaziano J, Norkus E, et al. Associations of plasma carotenoids with risk factors and biomarkers related to cardiovascular disease in middle-aged and older women. *Am J Clin Nutr* 2008;88(3):747-54.

Table I. General characteristics of the sample stratified by BMI groups

Variables	EU	OW	OI	OII	p-value
BMI (kg/m ²)	22.8 ± 1.1	27.3 ± 1.1	33.0 ± 0.8	37.4 ± 2.6	0.003*
Age (years)	53.0 ± 2.2	52 ± 2.1	53.0 ± 2.3	54.0 ± 2.5	0.831
WC (cm)	79.3 ± 6.1	112.3 ± 10.2	120.3 ± 9.5	120.3 ± 9.5	0.005*
DAIVA (µg)	795.1 ± 49,9	781.5 ± 34.5	756.6 ± 33.7	732.3 ± 26.0	0.065

Mean and standard deviation. BMI: body mass index; DAIVA: daily intake of vitamin A; EU: eutrophic; OI: obesity class I; OII: obesity class II; OW: overweight; WC: waist circumference. *p ≤ 0.005.

**Nutrición
Hospitalaria**

Table II. Mean of serum retinol ($\mu\text{mol/l}$) and β -carotene ($\mu\text{g/dl}$) concentrations in the groups studied

	Retinol ($\mu\text{mol/l}$)	β -carotene ($\mu\text{g/dl}$)
Eutrophic	1.3 \pm 0.2	61.2 \pm 12.1
Overweight	1.0 \pm 0.3	43.5 \pm 5.9
Obesity class I	0.9 \pm 0.3	37.1 \pm 5.3
Obesity class II	0.7 \pm 0.3	33.1 \pm 4.9
<i>p</i> *	< 0.001**	< 0.005*

Mean and standard deviation. ANOVA test: * $p \leq 0.005$; ** $p \leq 0.001$.

**Nutrición
Hospitalaria**

Table III. Multiple logistic regression analysis of serum retinol and β -carotene deficiency, night blindness and inadequate waist circumference (> 80 cm) in the groups studied

	OW/EU	OR	OI/EU	OR	OII/EU	OR
Variables	OR [†]	95%*	OR [†]	95%*	OR [†]	95%*
Retinol (< 1.05 μ mol/l)	14.0	7.0-27.3	35.3	17.6-68.8	65.4	32.7-127.5
β -carotene (< 40 μ g/dl)	5.1	2.5-9.9	17.0	8.5-24.6	26.7	13.3-52.0
NB	3.5	1.7-6.8	7.4	3.7-14.4	9.2	4.6-17.9
Inadequate WC	5.1	2.5-9.9	7.2	3.6-14.0	8.0	4.0-15.6

O1/EU: obesity class I/eutrophic; O2/EU: obesity class II/eutrophic; OW/EU: overweight/eutrophic. OR[†]: odds ratio; OR 95%*: confidence interval of 95%; NB: night blindness; WC: waist circumference.

**Nutrición
Hospitalaria**

Table IV. Correlation between serum retinol and β -carotene concentrations in each group studied

Groups	Retinol		β -carotene	
	<i>r</i>	<i>p</i>	<i>r</i>	<i>p</i>
Eutrophic	0.52	< 0.001*	0.66	< 0.001*
Overweight	-0.75	< 0.001*	-0.81	< 0.001*
Obesity class I	-0.74	< 0.001*	-0.86	< 0.001*
Obesity class II	-0.84	< 0.001*	-0.90	< 0.001*

Pearson's correlation. * $p \leq 0.001$.

**Nutrición
Hospitalaria**




Fig. 1. Flow chart describing participant recruitment of the study.

