

OR 1666

**Vitamin A supplementation according to zinc status on oxidative stress levels in cardiac surgery patients**

*Suplemento de vitamina A según el estado de zinc en los niveles de estrés oxidativo en pacientes con cirugía cardíaca*

Andréa Matos<sup>1,2</sup>, Gisele Souza<sup>2</sup>, Vanessa Moreira<sup>2</sup>, Mariana Luna<sup>2</sup> and Andréa Ramalho<sup>2</sup>

<sup>1</sup>Nutrition and Dietetic Department. Fluminense Federal University. Rio de Janeiro, Brazil. <sup>2</sup>Social Applied Nutrition Department. Micronutrients Research Center (NPqM). Federal University of Rio de Janeiro. Rio de Janeiro, Brazil

**Received:** 06/11/2017

**Accepted:** 31/01/2018

**Correspondence:** Andréa Matos. Nutrition and Dietetic Department. Fluminense Federal University. Mário Santos Braga Street, 30, Niterói, Rio de Janeiro. Brazil

e-mail: amatosnut@gmail.com

DOI: 10.20960/nh.1666

**ABSTRACT**

**Introduction:** it is widely reported that oxidative stress increases in patients undergoing cardiovascular surgery and causes negative outcomes, such as reduced immunological responses and postoperative complications. In this context, vitamin A and zinc serves an important function in the immunological system because it plays a role in fighting oxidative stress.

**Objectives:** to evaluate the effect of vitamin A supplementation on oxidative stress levels in cardiac surgery patients based on zinc nutritional status.

**Methods:** a control group (G1) and a vitamin A intervention group (G2- 5,000 IU of vitamin A daily) were subdivided based on their zinc nutritional status. The patients' serum concentrations of retinol,  $\beta$ -carotene, zinc and malondialdehyde were assessed before surgery (T0), on the 3<sup>rd</sup> (T1) and 21<sup>st</sup> postoperative day (T2).

**Results:** in the individuals with adequate zinc concentrations, the retinol concentrations were

significantly higher in G2 than in G1 at T1 and T2. In G2, the  $\beta$ -carotene concentrations were significantly higher in individuals with adequate zinc concentrations compared to those who had inadequate concentrations at T1 ( $p = 0.024$ ) and T2 ( $p = 0.043$ ). After surgery, malondialdehyde levels were significantly higher in individuals who had inadequate zinc concentrations, compared to those who had adequate zinc concentrations. Analyzing the clinical evolution, operative mortality, hospitalization time and length of stay in Intensive Care Unit (ICU) were lower in G2.

**Conclusion:** vitamin A has contributed to the reduction of oxidative stress, particularly in patients with adequate zinc concentrations and, this way, may help decrease the risk of postoperative complications.

**Key words:** Vitamin A. Zinc. Oxidative stress. Malondialdehyde. Cardiac surgery.

## RESUMEN

**Introducción:** se ha informado ampliamente que el estrés oxidativo aumenta en pacientes sometidos a cirugía cardiovascular y provoca resultados negativos, como respuestas inmunológicas reducidas y complicaciones postoperatorias. En este contexto, la vitamina A y el zinc desempeñan una función importante en el sistema inmunológico porque este desempeña un papel en la lucha contra el estrés oxidativo.

**Objetivos:** evaluar el efecto de la suplementación con vitamina A en los niveles de estrés oxidativo en pacientes sometidos a cirugía cardíaca basado en el estado nutricional del zinc.

**Métodos:** los pacientes fueron subdivididos en un grupo control (G1) y un grupo de intervención con vitamina A (G2, que recibió 5.000 UI de vitamina A diariamente) en función de su estado nutricional de zinc. Se evaluaron las concentraciones séricas de retinol,  $\beta$ -caroteno, zinc y malondialdehído de los pacientes antes de la cirugía (T0), en el tercer día postoperatorio (T1) y el vigesimoprimer día postoperatorio (T2).

**Resultados:** en los individuos con concentraciones adecuadas de zinc, las concentraciones de retinol fueron significativamente mayores en el G2 que en el G1 en T1 y T2. En el G2, las concentraciones de  $\beta$ -caroteno fueron significativamente mayores en los individuos con concentraciones adecuadas de zinc en comparación con los que tuvieron concentraciones inadecuadas en T1 ( $p = 0,024$ ) y T2 ( $p = 0,043$ ). Después de la cirugía, los niveles de malondialdehído fueron significativamente más altos en los individuos que tenían concentraciones inadecuadas de zinc, en comparación con aquellos que tenían concentraciones

adecuadas de zinc. Analizando la evolución clínica, la mortalidad operatoria, el tiempo de hospitalización y la duración de la estancia en la Unidad de Cuidados Intensivos (UCI) fueron menores en G2.

**Conclusión:** la vitamina A ha contribuido a la reducción del estrés oxidativo, particularmente en pacientes con concentraciones adecuadas de zinc y, de esta manera, puede ayudar a disminuir el riesgo de complicaciones postoperatorias.

**Palabras clave:** Vitamina A. Zinc. Estrés oxidativo. Malondialdehído. Cirugía cardíaca.

## INTRODUCTION

Currently, non-communicable diseases (NCD) kill 40 million people each year, equivalent to 70% of all deaths globally (1). Cardiovascular disease (CVD) accounts for most NCD deaths and is considered as a serious global public health problem (1,2).

Coronary artery bypass grafting (CABG) is one of the most common cardiac surgical procedures for CVD patients (3). CABG is accepted as the standard treatment for ischemic coronary disease in many patient subgroups. It is widely reported that oxidative stress increases in patients undergoing cardiovascular surgery and can cause postoperative complications (3-5).

Malondialdehyde (MDA) is an indicator that is commonly used to assess oxidative stress. MDA is a product of lipid peroxidation, a process induced by free radicals, and it can be estimated by measuring thiobarbituric acid reactive substances (TBARS; nmol/l). This method is one of the best and most frequently used in basic and clinic research (6).

In this context, nutrition serves an important function in the immunological system because it plays a role in fighting oxidative stress. Vitamin A and zinc are among the micronutrients that perform this function.

Studies have shown the important role that vitamin A plays in preventing and/or delaying the atherogenesis process; it prevents LDL oxidation and the oxidative lesion of endothelium and decreases cytotoxicity of high-dose oxidized low-density lipoprotein (7,8). In addition, Ochoa et al. (4) reported a significant decrease in vitamin A serum concentration after cardiovascular surgery, implying that the decreased antioxidant defenses in these patients may increase the incidence of postoperative complications and mortality.

Zinc deficiency is an important factor that can affect the blood concentrations of vitamin A because zinc plays a role in the metabolism of this vitamin, particularly in its transportation (9,10).

The principal mechanisms behind this dependence are related to the regulatory role of zinc in vitamin A transport, which is mediated by protein synthesis and the oxidative conversion of retinol to retinaldehyde. This conversion requires the action of the zinc-dependent retinol dehydrogenase enzyme (8). Thus, zinc deficiency may contribute to vitamin A deficiency (11), and interfere with the intestinal absorption of retinol even in the presence of adequate liver reserves of vitamin A (12). Zinc is required for the synthesis and hepatic secretion of retinol binding protein (RBP) (10). In cases of zinc deficiency, the RBP synthesis may be impaired, which triggers a secondary vitamin A deficiency (12).

Some studies of retinol supplementation in patients suffering from cardiopathy have presented conflicting results. However, such studies do not assess the relationship between vitamin A supplementation and zinc nutritional status (13,14). Considering that the use of vitamin A supplementation needs to be accompanied by the evaluation of the nutritional status of zinc, the present study was carried out.

**Nutrición  
Hospitalaria**

## MATERIALS AND METHODS

---

This study is a controlled trial that enrolled 90 patients with a primary diagnosis of *angina pectoris* (using the criteria of the Canadian Cardiovascular Society [15]) who underwent coronary artery bypass grafting in a Rio de Janeiro public hospital. The study was approved by the local ethics committee (CAAE 0005.0.185.185-07), and all of the participants signed informed consent forms.

To achieve 80% statistical power and a significance level of 5% ( $p < 0.05$ ), the sample size was calculated as 30 patients for each group. The patients were assigned to one of two groups (2 x 1). The control group patients (G1 = 60) received a standard diet in accordance with the usual practices of the hospital. For 21 days, the individuals in the vitamin A intervention group (G2 = 30) received a daily supplementation of 5,000 IU of vitamin A in the form of retinol palmitate. These two groups were divided into two additional subgroups: patients with adequate zinc concentrations and patients with inadequate zinc concentrations.

Supplementation, delivered in the form of a capsule, began immediately after the blood was collected at baseline (T0), before surgery. The patients in both groups were observed for 21 days, and additional blood was collected at the 3<sup>rd</sup> day after surgery (T1) and the 21<sup>st</sup> day after surgery (T2).

The inclusion criteria were as follows: a) admission for elective coronary artery bypass grafting surgery; b) age  $\geq 20$  years old (16); and c) formal authorization was given to participate in the study. Considering that hypoalbuminemia is associated with a worse postoperative prognosis, the patients who had serum albumin  $\leq 3.5$  g/dl were automatically excluded from the study.

The individuals with the following conditions were also excluded: malabsorption syndrome, acute and chronic infections, liver disease, kidney disease, alcoholism or acquired immunodeficiency syndrome (AIDS), as well as individuals who had taken medication or a vitamin supplement containing vitamin A or zinc in the last six months or who underwent prior revascularization surgery.

High performance liquid chromatography (HPLC-UV) was used to quantify the serum levels of retinol and  $\beta$ -carotene (17). In the current study, inadequate levels of serum retinol and  $\beta$ -carotene were defined as  $< 30$   $\mu\text{g}/\text{dl}$  and  $\leq 40$   $\mu\text{g}/\text{dl}$ , respectively (18,19). The zinc levels were evaluated using atomic absorption spectrometry, and inadequate zinc values were defined as  $< 0.7$  mg/l (20). The MDA levels were evaluated using spectrophotometry (21) and the HPLC-UV method (22). Body mass index (BMI) was assessed by dividing weight (kg) by height (m), squared,

and using the World Health Organization (WHO) (1995) cut-off points (23).

Hypertension, dyslipidemia and/or diabetes mellitus diagnoses were obtained from the patients' medical records. The lipid profile was classified according to the Brazilian Society of Cardiology's V Brazilian Guidelines on Dyslipidemias and Atherosclerosis Prevention (24). Serum glucose levels were classified according to the standards of the American Diabetes Association (25). Blood pressure was classified according to the VII Brazilian Guidelines on Hypertension (26).

All deaths occurring during the duration of hospitalization or the first month after surgery were counted as operative mortality. The hospitalization time and the period of ICU stay were noted through visits and checking the patients' medical records.

To analyze the data, the central tendency and dispersion measurements were calculated. To assess the association between the categorical variables, a Chi-squared test was employed. To compare the continuous variables, Student's t-test was employed, as well as Spearman's correlation coefficient. The statistical analysis was conducted with the Statistical Package for the Social Sciences (SPSS version 20).

**Nutrición  
Hospitalaria**

## RESULTS

The sample included 90 patients who were divided into two groups by a randomized table: 60 patients in G1 and 30 patients in G2.

There were no statistically significant differences observed between the groups in terms of the general patient characteristics (Table I).

There was no statistically significant difference in the vitamin A serum levels (retinol and  $\beta$ -carotene) between the groups at T0. At T1 and T2, statistically significant differences were found between the groups for both variables (Table II).

The serum retinol concentrations measured before and after the surgical procedure decreased significantly from T0 to T1 in G1 ( $p < 0.001$ ) and G2 ( $p < 0.001$ ). There was an increase at T2 in G1 and G2; however, these concentrations did not reach the levels initially observed at T0 for the individuals with adequate zinc concentrations in G1 or G2 ( $p < 0.001$ ). The same result was also observed in the individuals who had inadequate zinc concentrations.

For both G1 and G2, zinc-deficient patients had retinol serum concentrations that were significantly lower at T1 ( $p < 0.001$  for both groups) and T2 ( $p < 0.001$  for both groups), compared to the patients with adequate zinc concentrations in both groups.

The  $\beta$ -carotene concentrations were significantly higher in G2 than in G1 in the individuals with adequate and inadequate zinc concentrations at T1 (G1 and G2,  $p = 0.024$  and  $p < 0.001$ , respectively) and at T2 (G1 and G2,  $p < 0.001$  and  $p = 0.047$ , respectively) (Table II).

Comparing the zinc levels of the G2 patients, we found that the  $\beta$ -carotene concentrations were significantly higher in the individuals with appropriate zinc concentrations at T1 ( $p < 0.001$ ) and T2 ( $p = 0.043$ ), compared to the individuals with inadequate concentrations.

As shown in table III, the prevalence of inadequate retinol and  $\beta$ -carotene was lower in G2 than in G1 for the patients who had adequate and inadequate serum concentrations of zinc, although the difference was more pronounced in those with inadequate concentrations of zinc.

After supplementation, a statistically significant, strong, positive correlation was found between the retinol and  $\beta$ -carotene concentrations ( $r = 0.852$  and  $p = 0.014$ ) in the patients who had adequate serum zinc concentrations.

After surgery, the mean MDA levels were significantly higher in the individuals who had inadequate zinc concentrations, compared to the individuals who had adequate zinc concentrations (Table IV).

No statistically significant difference was found between the MDA results obtained through HPLC-

UV and the results obtained through spectrophotometry ( $p = 0.051$ ).

There was no statistically significant relationship found between MDA levels and vitamin A serum concentrations when we analyzed patients according to their zinc nutritional status. Note that among the patients with zinc inadequacy, 69% had both retinol and  $\beta$ -carotene deficiencies. The deficiencies were more common in the patients with higher circulating MDA concentrations (located at quartiles 50 and 75 immediately after the surgery), but the difference did not reach statistical significance.

Analyzing the clinical evolution, a shorter hospital stay in G2 ( $13.4 \pm 9.4$ ) compared to G1 ( $25.4 \pm 17.3$ ) was observed in patients with adequate levels of zinc ( $p = 0.001$ ).

Regarding the length of stay in ICU, this difference was observed in those with adequate levels of zinc ( $G1 = 8.5 \pm 9.9 / G2 = 4.6 \pm 6.4$ ,  $p = 0.047$ ) and inadequate ( $G1 = 8.7 \pm 10.5 / G2 = 2.7 \pm 0.4$ ,  $p = 0.039$ ). Operative mortality was found to be 8.33% in G1 and 3.33% in G2.

**Nutrición  
Hospitalaria**



## DISCUSSION

---

The antioxidant role that vitamin A plays in CVD patients has been documented in the literature (27,28). However, its relationship with zinc nutritional status in these individuals has not yet been described.

Patients undergoing coronary artery bypass grafting are exposed to high oxidative stress that increases the demand for antioxidants. In addition, they suffer a period of ischemia-reperfusion that exacerbates the production of free radicals and further depletes the nutrients serving an antioxidant function (4,5).

In the current study, a significant post-surgery reduction in the retinol and  $\beta$ -carotene serum concentrations was observed in G1 at T1 and T2 in the patients with adequate and inadequate serum zinc concentrations. This result corroborates the study by Schindler et al. (29), who found a significant reduction in retinol concentrations 48 hours after surgery, and the study by Gu et al. (30), who found significantly lower  $\beta$ -carotene concentrations in the patients undergoing coronary artery bypass grafting. These studies did not assess the patients' zinc concentrations.

Although the human body has antioxidant defense mechanisms that protect tissues from the cellular damage induced by oxidative stress, stress remained high in the assessed patients, and this stress level may result in the significant depletion of antioxidant nutrients (5). This depletion may explain the mean concentrations of MDA after surgery, which were significantly higher in the individuals who had inadequate serum zinc concentrations than in the individuals who had adequate concentrations in G1 ( $p < 0.001$ ) and G2 ( $p < 0.001$ ) at T1 and in G1 ( $p < 0.001$ ) and G2 ( $p = 0.002$ ) at T2. These findings correspond with those of Bao et al. (31), who found an inverse association between concentrations of zinc and malondialdehyde in a double-blinded study. Jain et al. (32) found a positive correlation between the total antioxidant capacity and low serum zinc concentrations.

Based on the serum zinc concentrations, no correlation was observed between the concentrations of MDA and vitamin A. Because of strict homeostatic control, zinc concentrations can fluctuate above real limits, even in individuals with low body stores of this nutrient. Maintaining a relatively constant zinc concentration is essential to ensure the normal functioning of an organism. Thus, serum and plasma concentrations are often the methods that are utilized to evaluate the nutritional state of zinc in the body (20).

The data from the current study demonstrate that inadequate serum retinol and  $\beta$ -carotene were more common in patients with higher circulating concentrations of MDA (located at quartiles 50

and 75). Although they did not reach statistical significance, these findings match the observation of Stahl et al. (33) that carotenoids are related to the inhibition of TBARS formation, and they also corroborate Hozawa et al. (34) who, in a longitudinal study that lasted 15 years, demonstrated that carotenoids are negatively correlated with F2-isoprostanes, an oxidative stress product.

Adequate vitamin A nutritional status reduces the biotransformation of carotenoids into retinol, which demonstrates the relationship between the nutritional status of retinol and the nutritional status of carotenoids (35).  $\beta$ -carotene is known as the most powerful retinol precursor. The current study corroborates the findings of Mecocci et al. (35), who demonstrated that a decrease in  $\beta$ -carotene serum concentration corresponds to a decrease in circulating retinol, suggesting the mobilization of  $\beta$ -carotene for retinol conversion. This finding deserves attention, because the reduction in  $\beta$ -carotene serum concentration follows the increase in oxidative stress, demonstrating that this powerful sweeper of free radicals could most likely be diverted to other important functions, such as fighting oxidative stress, to maintain an adequate retinol nutritional status.

Zinc nutritional status can also affect the blood concentration of vitamin A because it damages hepatic synthesis and RBP secretion (10,11). These findings show that significantly lower retinol and  $\beta$ -carotene serum concentrations after surgery in individuals with zinc deficiency may indicate a relationship between zinc nutritional status and vitamin A serum concentrations, as observed in patients undergoing coronary artery bypass grafting. However, no previous studies have shown that evaluating zinc nutritional status is truly a valid measure without considering the levels of circulating RBP. Thus, the results related to vitamin A supplementation in such patients may be masked. Studies that focus on the variables that are attributable to RBP levels and other factors, such as the use of fibrates and thiazolidinediones, thyroid function, iron levels and ferritin levels, should be performed.

Moreover, there are no available data concerning the patients with profiles such as those presented here, which makes it impossible to compare the results to other studies, although this relationship is well established in the classic risk group literature (36,37).

The results showed relevant improvement in the vitamin A levels in G2 patients who had adequate serum zinc concentrations and, consequently, less oxidative stress, which can contribute to a better prognosis among these patients.

Mortality is one of the possible complications after myocardial revascularization (38), and data about this outcome can vary widely depending on the center, country and other factors (39). In

our research operative mortality was found to be 8.33% in the control group and 3.33% in the supplemented group. Gomes et al. (40) evaluated Brazilian patients undergoing various types of cardiac surgeries, among them, CABG, and the overall postoperative mortality, up to seven days post-surgery, was 4%, and 2.6% for CABG, being similar to G2. The mortality rates documented in developed nations after cardiac surgeries in general are 2.1% and for CABG are 1.8% (41).

A study that assessed surgical patients found a connection between retinol deficiency and postoperative complications (42). However, this study did not address serum zinc concentrations. According to Barbosa, in a double-blind, placebo-controlled study, supplementation with antioxidants, including zinc, yielded greater protection against oxidative stress (43). We noticed a significantly shorter time in hospital among those from the supplemented group with adequate zinc levels, and with no statistically significant difference in the zinc deficient patients, suggesting that the dosage of vitamin A used was not effective in individuals with hypozincemia.

Although no study with the same characteristics than the present one was found, Shehzad et al. (44), in a randomized controlled trial, assessed 300 children hospitalized with pneumonia and observed a significantly shorter duration of hospital stay in the supplemented group (20 mg zinc/day q 12 hourly) than the placebo group ( $p < 0.001$ ). In contrast, Yuan et al. (45) also evaluated hospitalized children with pneumonia ( $n = 96$ ), and did not find significant reduction in the duration of hospital stay between the supplemented group and placebo.

Vitamin A supplementation was shown to have a positive effect regardless of zinc concentrations by analyzing length of stay in ICU. There was a significant difference between the groups both in patients with adequate and inadequate zinc concentrations. Visser et al. (46), in a systematic review and meta-analysis, observed that micronutrient supplementation, including zinc, was not associated with a significant difference in the length of ICU stay. Still in this context, Hasanzadeh Kiabi et al. (47) assessed 186 adults during their ICU stay and found that zinc supplementation may be associated with a significant reduction in the occurrence of ventilator-associated pneumonia in adult mechanically ventilated trauma patients.

The results of the current study showed that vitamin A supplementation had a positive impact on individuals with adequate zinc concentrations by combating oxidative stress in these patients, thus reducing the risk of postoperative complications. The findings presented here can be considered to be an important contribution to the issue addressed, and they may also generate subsidies to future studies of the postoperative management protocol for the group in question, and improve the outcomes for these patients.

## ACKNOWLEDGMENTS

We thank the National Council of Technological and Scientific Development and the National Institute of Cardiology of Laranjeiras for their research support for this study and the American Journal Experts for the review of the manuscript.

## REFERENCES

1. World Health Organization. Noncommunicable diseases (NCD). Geneva: WHO. Accessed July 9, 2017. Available from: <http://www.who.int/gho/ncd/en/>
2. Organização Pan-Americana de Saúde/WHO. Doenças cardiovasculares. Setembro de 2016. Available from: [http://www.paho.org/bra/index.php?option=com\\_content&view=article&id=5253:doencas-cardiovasculares&Itemid=8398](http://www.paho.org/bra/index.php?option=com_content&view=article&id=5253:doencas-cardiovasculares&Itemid=8398), accessed July 09, 2017.
3. Linda T, Alessandro DM, Benedetta P, Isabella S, Alice B, Chiara M. Assessing free-radical-mediated DNA damage during cardiac surgery: 8-Oxo-7,8-dihydro-2'-deoxyguanosine as a putative biomarker. *Ox Med Cell Longev* 2017;2017:1-8. DOI: 10.1155/2017/9715898.
4. Ochoa JJ, Vílchez MJ, Ibáñez S, Huertas JR, Palacio MA, Muñoz-Hoyos A. Oxidative stress is evident in erythrocytes as well as plasma in patients undergoing heart surgery involving cardiopulmonary bypass. *Free Radic Res* 2003;37(1):11-7.
5. Chambers DJ. Oxidative stress injury during cardiac surgery: how important is it? *Cardiovasc Res* 2007;73(4):626-8.
6. Tsikas D, Rothmann S, Schneider JY, Suchy MT, Trettin A, Modun D, et al. Development, validation and biomedical applications of stable-isotope dilution GC-MS and GC-MS/MS techniques for circulating malondialdehyde (MDA) after pentafluorobenzyl bromide derivatization: MDA as a biomarker of oxidative stress and its relation to 15(S)-8-iso-prostaglandin F2 $\alpha$  and nitric oxide (NO). *J Chromatogr B Analyt Technol Biomed Life Sci* 2016;1019:95-111. DOI: 10.1016/j.jchromb.2015.10.009
7. Delport R. Antioxidants and coronary artery disease risk in South African males. *Cl Chim Acta* 1998;278(1):55-60.
8. Kaliora AC, Dedoussis GV, Schmidt H. Dietary antioxidants in preventing atherogenesis. *Atheros* 2006;187(1):1-17.
9. Mafra D, Cozzolino S. The importance of zinc in human nutrition. *Rev Nutr* 2004;17(1):79-

87.

10. Silva L, Thiapó A, Souza G, Saunders C, Ramalho A. Micronutrients in pregnancy and lactation. *Rev Bras Saude Mater Infant* 2007;7(3).

11. Satre MA, Jessen KA, Clegg MS, Keen CL. Retinol binding protein expression is induced in HepG2 cells by zinc deficiency. *FEBS Lett* 2001;491(3):266-71.

12. Ferraz I, Daneluzzi J, Vannucchi H, Jordão Jr A, Ricco R, Del Ciampo L, et al. Nível sérico de zinco e sua associação com deficiência de vitamina A em crianças pré-escolares. *J Pediatr* 2007;83(6).

13. Riccioni G, Bucciarelli T, Mancini B, Ilio CD, Capra V, D'Orazio N. The role of the antioxidant vitamin supplementation in the prevention of cardiovascular diseases. *Expert Opin Invest Drugs* 2007;16(1):1-8. DOI: 10.1517/13543784.16.1.25.

14. Gale CR, Ashurs HE, Power HJ, Martyn CN. Antioxidant vitamin status and carotid atherosclerosis in the elderly. *Am J Clin Nutr* 2001;74:402-8.

15. Campeau L. Grading of angina pectoris. *Circulation* 1976;54:522-3.

16. World Health Organization. WHO discussion papers on adolescence - Issues in adolescent health and development. Geneva: WHO; 2004.

17. Clae-Uv. The United States Pharmacopeia (Usp 28). The National Formulary (NF 23) (2004). The United States Pharmacopeia Convention Inc. 12601 Twinbrook Parkway, Rockville, MD 20852, 2353-4.

18. World Health Organization. Indicators for assessing vitamin A deficiency and their application in monitoring and evaluating intervention programmes. Micronutrient Series, WHO/NUT 10. Geneva: WHO; 1996.

19. Sauberlich HE, Hodges RE, Wallace DL, Kolder H, Canham J, Hood J, et al. Vitamin A metabolism and requirements in the human studied with the use of labeled retinol. *Vitam Horm* 1974;32:251-75.

20. Hotz C, Pearson JM, Brown KH. Suggested lower cutoffs of serum zinc concentrations for assessing zinc status: reanalysis of the second National Health and Nutrition Examination Survey data (1976-1980). *A J Clin Nutri* 2003;78:756-64.

21. Ohkawa H, Oshishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* 1979;95:351-8.

22. Karatas F, Karatepe M, Baysar A. Determination of free malondialdehyde in human serum by high-performance liquid chromatography. *Anal Biochem* 2002;311:76-9.

23. World Health Organization. Physical status: the use and interpretation of anthropometry. Report of a WHO expert committee. Geneva: WHO; 1995.
24. Sociedade Brasileira de Cardiologia. Atualização da Diretriz Brasileira de Dislipidemias e Prevenção da Aterosclerose. Departamento de Aterosclerose da Sociedade Brasileira de Cardiologia. Arq Bras Cardiol 2017;109(1):1-90.
25. American Diabetes Association. Standards of medical care in diabetes. Diabetes Care 2017;40(Suppl 1):S4-5.
26. Sociedade Brasileira de Cardiologia. 7ª Diretriz Brasileira de Hipertensão Arterial. Arq Bras Cardiol 2016;107(3 Suppl 3):1-83.
27. Sesso HD, Buring JE, Norkus EP, Gaziano JM. Plasma lycopene, other carotenoids, and retinol and the risk of cardiovascular disease in women. Am J Clin Nutr 2004;79:47-53.
28. Matos AC, Souza GG, Moreira V, Ramalho A. Effect of vitamin A supplementation on clinical evolution in patients undergoing coronary artery bypass grafting, according to serum levels of zinc. Nutr Hosp 2012;27(6):1981-6. DOI: 10.3305/nh.2012.27.6.5891.
29. Schindler R, Berndt S, Schroeder P, Oster O, Rave G, Sievers HH. Plasma vitamin E and A changes during cardiopulmonary bypass and in the postoperative course. Langenbecks Arch Surg 2003;387(9-10):372-8. DOI: 10.1007/s00423-002-0336-4.
30. Gu M, Love H, Schofield D, Turkie W, Odom N, Braganza JM. A pilot study of blood antioxidant and free radical marker profiles in patients awaiting coronary artery bypass grafting. Clin Chim Acta 1996;252(2):181-95.
31. Bao B, Prasad AS, Beck FW, Fitzgerald JT, Snell D, Bao GW, et al. Zinc decreases C-reactive protein, lipid peroxidation, and inflammatory cytokines in elderly subjects: a potential implication of zinc as an atheroprotective agent. Am J Clin Nutr 2010;91(6):1634-41. DOI: 10.3945/ajcn.2009.28836
32. Jain A, Varma M, Agrawal BK, Jadhav AA. Serum zinc and malondialdehyde concentrations and their relation to total antioxidant capacity in protein energy malnutrition. J Nutr Sci Vitaminol 2008;54(5):392-5.
33. Stahl W. Lipid oxidation and antioxidants. Curr Opin Clin Nutr Metab Care 2000;3(2):121-6.
34. Hozawa A, Jacobs DR Jr, Steffes MW, Gross MD, Steffen LM, Lee DH. Relationships of circulating carotenoid concentrations with several markers of inflammation, oxidative stress, and endothelial dysfunction: the Coronary Artery Risk Development in Young Adults (CARDIA)/Young Adult Longitudinal Trends in Antioxidants (YALTA) study. Clin Chem 2007;53(3):447-55. DOI:

10.1373/clinchem.2006.074930

35. Mecocci P, Polidori C, Troiano L, Cherubini A, Cecchetti R. Plasma antioxidants and longevity: a study of health centenarians. *Free Radical Biol Med* 2000;28(8):1243-8.
36. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, et al; Maternal and Child Nutrition Study Group. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet* 2013;382(9890):427-51. DOI: [http://dx.doi.org/10.1016/S0140-6736\(13\)60937-X](http://dx.doi.org/10.1016/S0140-6736(13)60937-X)
37. Gebreselassie SG, Gase FE, Deressa MU. Prevalence and correlates of prenatal vitamin A deficiency in rural Sidama, Southern Ethiopia. *J Health Popul Nutr* 2013;31(2):185-94.
38. Ribeiro KRA. Postoperative complications of coronary artery bypass grafting: implications for the nursing practice. *Rev Enferm UFPI* 2017;6(3):59-64.
39. Monteiro GM, Moreira DM. Mortalidade em cirurgias cardíacas em hospital terciário do Sul do Brasil. *Int J Cardiovasc Sci* 2015;28(3):200-5.
40. Gomes WJ, Moreira RS, Zilli AC, Bettiati LC Jr, Figueira FAMDS, D'Azevedo SSP, et al. The Brazilian registry of adult patient undergoing cardiovascular surgery, the BYPASS Project: results of the first 1,722 patients. *Braz J Cardiovasc Surg* 2017;32(2):71-6.
41. Osnabrugge RL, Speir AM, Head SJ, Fonner CE, Fonner E, Kappetein AP, et al. Performance of EuroSCORE II in a large US database: implications for transcatheter aortic valve implantation. *Eur J Cardiothorac Surg* 2014;46(3):400-8.
42. Zago L, Dupraz H, Torino F, Río ME. Preoperative nutritional status and surgical risk: identification of promissory biochemical markers. *Nutr Hosp* 2010;25(1):91-8.
43. Barbosa E, Faintuch J, Moreira E. Supplementation of vitamin E, vitamin C, and zinc attenuates oxidative stress in burned children: a randomized, double-blind, placebo-controlled pilot study. *J Burn Care Res* 2009;30(5):859-66. DOI: 10.1097/BCR.0b013e3181b487a8
44. Shehzad N, Anwar MI, Muqaddas T. Zinc supplementation for the treatment of severe pneumonia in hospitalized children: a randomized controlled trial. *Sudan J Paediatr* 2015;15(1):37-41.
45. Yuan X, Qian SY, Li Z, Zhang ZZ. Effect of zinc supplementation on infants with severe pneumonia. *World J Pediatr* 2016;12(2):166-9.
46. Visser J, Labadarios D, Blaauw R. Micronutrient supplementation for critically ill adults: a systematic review and meta-analysis. *Nutrition* 2011;27(7-8):745-58.

47. Hasanzadeh Kiabi F, Alipour A, Darvishi-Khezri H, Aliasgharian A, Emami Zeydi A. Zinc supplementation in adult mechanically ventilated trauma patients is associated with decreased occurrence of ventilator-associated pneumonia: a secondary analysis of a prospective, observational study. Indian J Crit Care Med 2017;21(1):34-9.






**Table I. General characteristics of the study population**

<i>Characteristics</i>	<i>G1</i>		<i>G2</i>		<i>p</i>
	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	
<i>Age (years)</i>	65.06	9.81	63.6	9.3	0.68
<i>Male</i>	70		76.66		0.62
<i>Female</i>	30		23.34		
<i>BMI (kg/m<sup>2</sup>)</i>	28.25	3.75	27.09	2.76	0.7
<i>SH (%)</i>	100		100		*
<i>DLP (%)</i>	93.3		96.6		0.61
<i>DM (%)</i>	35		30		0.81

\*No possibility of statistical calculation. SH: systemic hypertension; DLP: dyslipidemia; DM: diabetes mellitus.

1



**Table II. Mean values of retinol and  $\beta$ -carotene serum concentrations, according to zinc nutritional status, before and after coronary artery bypass grafting**

		<i>Adequate zinc concentration</i>					
		<i>T0</i>	<i>SD</i>	<i>T1</i>	<i>SD</i>	<i>T2</i>	<i>SD</i>
<i>Retinol (<math>\mu\text{g}/\text{dl}</math>)</i>	<i>G1 (n = 50)</i>	50.4	14.6	29.3	17.7	38.7	17.1
	<i>G2 (n = 23)</i>	65.3	15.7	54.3	22.9	62.1	20.3
	<i>p</i>	0.051		< 0.001		< 0.001	
<i><math>\beta</math>-carotene (<math>\mu\text{g}/\text{dl}</math>)</i>	<i>G1 (n = 50)</i>	28.1	8.98	10	1.2	12.3	5.7
	<i>G2 (n = 23)</i>	38.5	13.1	46.5	14.2	53.5	20.9
	<i>p</i>	0.051		0.024		< 0.001	
		<i>Inadequate zinc concentration</i>					
		<i>T0</i>	<i>SD</i>	<i>T1</i>	<i>SD</i>	<i>T2</i>	<i>SD</i>
<i>Retinol (<math>\mu\text{g}/\text{dl}</math>)</i>	<i>G1 (n = 10)</i>	44.5	17.3	28.4	14.9	35.5	18.7
	<i>G2 (n = 7)</i>	52.1	17.7	47.2	26.5	51	28.5
	<i>p</i>	0.052		< 0.001		< 0.001	
<i><math>\beta</math>-carotene (<math>\mu\text{g}/\text{dl}</math>)</i>	<i>G1 (n = 10)</i>	25	6.3	10.9	6.1	8	4.2
	<i>G2 (n = 7)</i>	31.6	6.7	40.4	19.1	40.5	18.1
	<i>p</i>	0.06		< 0.001		0.047	

G1: control group; G2: intervention group; T0: before surgery; T1: 3<sup>rd</sup> day after surgery; T2: 21<sup>st</sup> day after surgery.

**Table III. Prevalence of inadequate retinol and  $\beta$ -carotene levels in G1 and G2 before and after coronary artery bypass grafting, according to zinc nutritional status**

		<i>Adequate zinc concentration</i>		
		<i>T0</i>	<i>T1</i>	<i>T2</i>
<i>Retinol (%)</i>	<i>G1</i>	23.3	90	53.3
	<i>G2</i>	30	18	30
<i><math>\beta</math>-carotene (%)</i>	<i>G1</i>	87.5	100	100
	<i>G2</i>	88.8	50	50.5
		<i>Inadequate zinc concentration</i>		
		<i>T0</i>	<i>T1</i>	<i>T2</i>
<i>Retinol (%)</i>	<i>G1</i>	85	87.3	85
	<i>G2</i>	76.6	56.6	70
<i><math>\beta</math>-carotene (%)</i>	<i>G1</i>	100	100	100
	<i>G2</i>	91.4	73.5	73.3

G1: control group; G2: intervention group; T0: before surgery; T1: 3<sup>rd</sup> day after surgery; T2: 21<sup>st</sup> day after surgery.

**Table IV. Mean concentrations of levels of malondialdehyde (nmol/l) according to the nutritional status of zinc in G1 and G2 before and after coronary artery bypass grafting**

	<i>T0</i>	<i>SD</i>	<i>T1</i>	<i>SD</i>	<i>T2</i>	<i>SD</i>
<i>G1</i>						
<i>Adequate zinc</i>	2.1	2.0	5.2	4.4	3.0	1.5
<i>Inadequate zinc</i>	2.6	3.6	7.0	4.9	3.5	2.8
<i>p</i>	0.48		< 0.001		< 0.001	
<i>G2</i>						
<i>Adequate zinc</i>	2.0	1.9	3.7	2.7	1.6	1.1
<i>Inadequate zinc</i>	2.3	3.6	6.2	2.2	2.6	1.6
<i>p</i>	0.053		< 0.001		0.002	

G1: control group; G2: intervention group; T0: before surgery; T1: 3<sup>rd</sup> day after surgery; T2: 21<sup>st</sup> day after surgery.

**Nutrición  
Hospitalaria**