

**REV 931**

**The nutritional limitations of plant-based beverages in infancy and childhood**

Isidro Vitoria

Unit of Nutrition and Metabolopathies. Hospital Universitario y Politécnico La Fe.  
Valencia, Spain

**Received:** 18/01/2017

**Accepted:** 20/02/2017

**Correspondence:** Isidro Vitoria Miñana. Unit of Nutrition and Metabolopathies.  
Hospital Universitario y Politécnico La Fe. Av. Fernando Abril Martorell, 106. 46026  
Valencia, Spain

e-mail: vitoria\_isi@gva.es

**DOI:** 10.20960/nh.931

**ABSTRACT**

Breastfeeding, infant formula and cow's milk are basic foods in infant nutrition. However, they are being increasingly replaced either totally or partially by plant-based beverages.

The composition of 164 plant-based beverages available in Spain was reviewed based on the nutritional labeling of the package and the manufacturers' webpages. This was compared to the composition of cow's milk and infant formula. In addition, the nutritional disease associated with consumption of plant-based beverages in infants and children was reviewed by means of a literature search in Medline and Embase since 1990 based on the key words "plant-based beverages" or "rice beverages" or "almond beverages" or "soy beverages" and "infant" or "child".

The nutritional composition of 54 soy beverages, 24 rice beverages, 22 almond beverages, 31 oat beverages, 6 coconut beverages, 12 miscellaneous beverages and 15 mixed beverages was described. At least 30 cases of nutritional disease in children associated with nearly exclusive consumption of plant-based beverages have been

published. A characteristic association has been observed between soy beverage and rickets, rice beverage and kwashiorkor, and almond-based beverage and metabolic alkalosis.

The nutritional quality of plant-based beverages is lower than that of cow's milk and infant formula, therefore they are not a nutritional alternative. Predominant or exclusive use of these beverages in infant feeding can lead to serious nutritional risks. In the case of nonexclusive feeding with these beverages, the pediatrician should be aware of the nutritional risks and limitations of these beverages in order to complement their deficiencies with other foods.

**Key words:** Milk substitutes. Beverages. Soy milk. Infant formula. Failure to thrive. Kwashiorkor. Metabolic alkalosis.

## INTRODUCTION

Breastfeeding and complementary feeding in the first year of life achieve adequate growth of the child (1). When breastfeeding is not possible or supplements are required, infant formula from cow's milk is recommended. The composition of these formulas must meet nutritional recommendations (2). However, when breastfeeding or formula is replaced by other beverages, serious nutritional consequences may result (3).

The intake of plant-based beverages (PBBs) in the early years of life has increased in recent years (4). The main reasons for this change are preference for plant foods, aversion to the use of cow's milk, and prevention or treatment of cow's milk allergy, as part of strict vegetarian diets or as a consequence of the advice of professionals from alternative medicines (5). Primary use in the early years of life of mainly soy, rice, almond or oat PBBs results in nutritional risks (rickets, failure to thrive, kwashiorkor or metabolic alkalosis, among others) (6). In addition, in our country we reported a case of scurvy with bone fractures in an infant fed almost exclusively with almond beverages (7).

The aim of this study was to review the composition of PBBs marketed in Spain in order to compare them to the nutritional recommendations for infant formulas and to

the composition of cow's milk. In addition, publications on nutritional disease associated with consumption of PBBs in children were reviewed in order to determine whether there was a specific type of nutritional disease associated with each type of PBB.

## **MATERIALS AND METHODS**

The composition of 164 brands of PBBs marketed in Spain was reviewed (54 soy beverages, 24 rice beverages, 22 almond beverages, 31 oat beverages, 6 coconut beverages, 12 beverages from other miscellaneous plants [barley, canary grass, hazelnut, hemp, macadamia nut, sesame or spelt] and 15 mixed PBBs [rice and almond, coconut, hazelnut or quinoa, and oat with coconut or almond]). Composition values were taken from the nutritional labels of the PBB packages purchased in stores and from the manufacturers webpages, where the content in kilocalories, carbohydrates, sugars, proteins, total fats and saturated, monounsaturated, and polyunsaturated fatty acids, fiber, salt and supplements, if any, both minerals and vitamins, were specified. The composition of PBBs was compared to the recommended composition of infant formula and soy infant formula (2) and to the composition of cow's milk (8).

In addition, the literature since 1990 on nutritional disease associated with primary intake of PBBs in children was reviewed by means of a search in Medline and Embase based on the key words "plant based beverages" or "rice beverages" or "almond beverages" or "soy beverages" and "infant" or "child".

## **RESULTS**

Table I shows the mean content in kilocalories, macronutrients, percentage of energy/protein, number of brands supplemented with calcium, vitamin D and other minerals and vitamins for each group of PBBs, as well as the recommended composition of soy infant formula, infant formula and composition of cow's milk.

The composition of 54 brands of soy beverages is shown in supplemental table I (<http://www.xxxxxxxx>). The mean calorie content was  $46.7 \pm 13.1$  kcal/100 ml. There was no uniformity in their composition as shown by the wide energy range (27-80.7). In 43 of the 54 brands, calorie provision was less than 60 kcal/100 ml. Protein content

was 2.1-3.8 g/100 ml. Forty-three brands were supplemented with calcium and 23 of these were also supplemented with vitamin D. The most commonly added amounts were 120 mg of calcium per 100 ml and 0.75 µg of vitamin D per 100 ml. Other vitamins were added in 25 soy beverages, especially B<sub>2</sub>, B<sub>12</sub> and A. Only two of the 55 soy beverages included added minerals, such as iron.

The composition of 24 brands of rice beverages marketed in our country is shown in supplemental table II (<http://www.xxxxxxxx>). These beverages had a mean calorie content of 56.8 ± 6.3 kcal/100 ml, with a range from 47 to 68 kcal/100 ml, a low mean protein content of 0.3 ± 0.2 g/100 ml and low fat levels (0.8-2 g/100 ml). Of the 24 brands, only eight specified the added amounts of calcium and only five of these, the added amounts of vitamin D. The percentage of energy provided by proteins was less than 3% in most cases.

The composition of 22 different brands of almond beverages is shown in supplemental table III (<http://www.xxxxxxxx>). Almond beverages are hypocaloric and hypoproteic beverages as compared to infant formula and cow's milk. Mean calorie provision was 40.2 ± 14.3 kcal/100 ml, and equal to or less than 60 kcal/100 ml in 19 of the 22 brands studied. Protein content was 0.3-1.6 g/100 ml. Carbohydrate content was intermediate between soy and rice beverages. Of the 22 brands, only five were supplemented with calcium and vitamin D.

Mean content of the rest of beverages studied is shown in supplemental tables IV and V. The group of oat, coconut and miscellaneous beverages comprised 49 brands. Mean calorie content was 44.9 ± 10.7 kcal/100 ml (range 15-65), mostly at the expense of carbohydrates (mean value 6.9 ± 2.5 g/100 ml, range 2-11) and to a lesser extent of fats (mean value 1.4 ± 0.8 g/100 ml, range 0.1-3.6). Mean protein content was low but not as low as for rice beverages (mean value 0.7 ± 0.2 g/100 ml, range 0.1-1.4). Only 13 of these 49 beverages were supplemented with calcium and vitamin D. As shown in table I, the group of six coconut beverages had the lowest calorie content of the PBBs (mean value 33.8 ± 15.1 kcal/100 ml) and a protein content similar to rice beverages (0.2 ± 0.2 g/100 ml). The mixed group of beverages included 15 brands (Supplemental Table VI [<http://www.xxxxxxxx>]). The composition of the 12 mixed beverages containing rice had a higher calorie content at the expense of carbohydrates and to a lesser extent of proteins than mixed oat beverages.

Following the literature review on nutritional disease related to primary intake of PBBs, 20 papers were found reporting 30 clinical cases associated with consumption of soy, rice or almond beverages (Tables II-IV).

## COMMENTS

### Composition of PBBs

Rice beverages were the PBBs with the highest energy content, although their mean value was lower than the minimum value of infant formula or cow's milk (60 kcal/100 ml). The rest of the PBB groups had a mean value lower than 50 kcal/ml. Thus, almond beverages and coconut beverages had a mean calorie content of 40 and 33 kcal/100 ml, respectively. Only mixed beverages, most of which contain rice, had a higher calorie content.

Overall analysis of all PBB groups showed that soy beverages had the highest protein content. However, the nutritional value of soy protein is limited by the content in methionine and cysteine, with a lowest digestible indispensable amino acid score (DIASS) value of 90.6%, based on the biological value and true ileal amino acid digestibility (9). The rice protein isolate also has a DIASS value of 37.1% (10)<sup>iError! Marcador no definido.</sup>. Regarding the rest of plants used in PBBs, no information was available about the DIASS of their proteins (4). However, the value of the protein digestibility-corrected amino acid score (PDCAAS) is known. The PDCAAS values of the raw materials used in some commercial PBBs are 67.7% (quinoa), 63-66% (hemp), 45-60% (oat), 54% (rice) and 30% (almond) (11). Consequently, infant formula, milk and other dairy products have higher value protein than PBBs.

With regard to carbohydrates, in most PBBs over 70% are sugars. According to the European regulation on nutritional labeling, sugars include monosaccharides and disaccharides but not polyols or starch (12). Formula intended for infants under 4-6 months should not contain fructose or sucrose (2) On the other hand, PBBs do not contain lactose. Lactose is considered to provide beneficial effects for gut physiology, including prebiotic effects, softening of stools, and enhancement of calcium absorption (2). In this regard, the 2014 EFSA proposal recommends that infant formula contains a minimum of 4.5 g/100 kcal of lactose (13). Fiber content was less than 0.5 g/100 ml in most cases (64 of 113 PBBs).

With regard to fats, only soy beverages had a profile with a clear predominance of polyunsaturated fatty acids, but their overall fat content was very low ( $1.8 \pm 0.4$  g/100 ml) as compared to the recommended total fat content of 2.8-3.9 g/100 ml in infant formulas, equivalent to about 40-54% of energy content, which is similar to values found in human milk. In almond beverages, the predominant fats were monounsaturated fatty acids, while in coconut beverages, saturated fatty acids were predominant. In all cases, mean fat content values were lower than for infant formula and cow's milk. Thus, fat content values were very low in rice and oat beverages (mean value 1 g/100 ml) and low (mean value 1.5-2 g/100 ml) in the rest of PBBs (Table I). Furthermore, no information was available about the minimum content in linoleic acid, erucic acid or the maximum values of trans fatty acids, among others (2).

With regard to minerals, divalent cations like zinc, magnesium and iron are bound by phytates present in all seeds, reducing their bioavailability (14). Processing treatments to prepare PBBs such as flaking, blanching, hot grinding and ultra-high temperature treatment could cause loss of vitamins (15). Therefore, the addition of minerals and vitamins after processing is important. Of the 164 PBBs in the present study, calcium and/or vitamin D were added in over half of the beverages, both other minerals and vitamins were added in only 43 cases.

### **Consumption of PBBs and nutritional disease**

The nutritional disorder most often associated with consumption of soy beverages in small children is rickets and failure to thrive, along with ferropenic anemia in some cases (6,16,17) (Table II). In the published cases, soy beverages were mostly given for suspected allergy to cow's milk proteins or due to the parents' belief that it was more suitable for their child. The age of the patients ranged from five to 17 months. The interval between the start of soy beverage consumption and diagnosis of rickets was from four to eight months, depending on the age at which consumption was started. The case reported by Imataka G et al. (10) began to take soy beverage at one month of age and developed hypocalcemic tetany.

According to ESPGHAN **Error! Marcador no definido.**, for soy protein infant formula, only protein isolates should be used, and the minimum protein content required by European legislation is higher than that of cow's milk protein infant formula (2.25

g/100 kcal vs 1.8 g/100 kcal) (1.5 to 1.2 g/100 ml) to account for potentially lower digestibility and, therefore, lower bioavailability of soy protein compared with intact cow's milk protein. According to the Committee on Nutrition of the American Academy of Pediatrics, the protein of the soy formula must be a soy isolate supplemented with L-methionine, L-carnitine and taurine to provide a protein content of 2.45 to 2.8 g per 100 kcal (1.65 to 1.9 g/100 ml), and phytases can be used (18). Mean protein content of soy beverages in this study was  $3.1 \pm 0.4$  g/100 ml, but the soy proteins of soy beverages were not supplemented with amino acids. For these reasons and despite being the PBBs with the highest mean protein content, their consumption in small children is probably associated with failure to thrive.

The reasons why a diet rich in non-supplemented soy beverage is a determining factor for rickets are related to calcium and vitamin D. Thus, regarding the calcium added, it depends on the type of salt used. The tricalcium phosphate present in many soy beverages supplemented with calcium is absorbed in a proportion of 75% to the calcium of cow's milk, whereas calcium carbonate has better absorption (19). In addition, heat treatment of commercial soy beverages precipitates the calcium (20), which is the reason for the large difference in calcium content depending on whether the sample is shaken or not (21). The absence of lactose and higher content in insoluble fiber also reduces calcium absorption (22). Moreover, when the type of vitamin D added is specified, it is vitamin D<sub>2</sub>, which has lower effectiveness than vitamin D<sub>3</sub> (23). Of the 54 brands studied, only 23 (42%) were supplemented with calcium and vitamin D.

Regarding the clinical manifestations secondary to the use of rice beverages, the information on 17 cases is given in supplemental table IV (<http://www.xxxxxxx>) (5,6,24-34). The principal nutritional consequence in infants of consumption of rice beverages instead of infant formula is protein malnutrition or kwashiorkor, reported in 14 of 17 cases, with clinical data of hypoalbuminemia, edema and rash. Kwashiorkor is a known case of failure to thrive and growth delay in developing countries. However, it is exceptional in developed countries. In the majority of published cases, rice beverage was given for suspected allergy to cow's milk protein. The age of patients at diagnosis ranges from four to 22 months. In ten of 17 cases, consumption of rice beverage was

started at four months or earlier. The interval between the start of consumption of rice beverage and diagnosis of kwashiorkor ranged from one to nine months.

The cause of kwashiorkor is the higher calorie content of rice beverages with a very low protein content (0.1-0.8 g/100 ml), which results in proteins accounting for  $2.4 \pm 1.4\%$  of energy, a significantly lower amount than the percentage of protein provided by breastmilk (5-6%) or infant formula (7-9%) (35). Of the different PBBs, the lowest value of the percentage of energy provided by proteins was that of rice beverages, followed by coconut beverages, which may have the same nutritional risk (Table I). In contrast with these data, children who develop marasmus have a deficient intake of both energy and proteins. The higher calorie provision of rice beverages was due the higher content in carbohydrates (9.4-14.2 g/100 ml). In addition, rice beverages contain no vitamins and are deficient in iron (0.07 mg/100 g) and calcium (0.9 mg/100 ml) (19), unless it is added. Only five of 24 brands of rice beverages were supplemented with calcium and vitamin D.

The clinical manifestations secondary to the use of almond beverages in small children are shown in table IV. Of the ten reported cases (5,7,33,36-39), metabolic alkalosis was noted in three. Since 1980, cases have been reported of similar conditions of hypochloremic and hypokalemic metabolic alkalosis in infants fed with milks lacking sodium chloride (40,41). In the case of almond beverages, the problem is the low chloride content. Thus, the French Food Safety Agency (ANSES) (42) determined that the cause of a case (33) was the low chloride content of the almond beverage (2.4 mg/100 ml). The lower amount of chloride anion due to the lack of intake leads to proximal tubular reabsorption of the bicarbonate anion together with the sodium anion, causing metabolic alkalosis (36). Although the composition stated on the packages is incomplete, according to Doron (38), they contain 0.4 mg sodium per 100 ml and 0.32 mg iron per 100 ml, much lower amounts than those recommended for infant formula.

Almond beverages may also be responsible for severe rickets (38), which may be accompanied by seizure-inducing hypocalcemia. Other authors have reported three cases of hyperoxaluria in children aged three to eight years who took more than 500 ml daily of almond beverages. One of them had kidney stones. The cause is the higher content in oxalates of almond beverages, particularly if they are obtained from



homemade almond milk (39). A case of scurvy has also been reported in an 11-month-old infant who took almond beverage and almond flour prescribed by a physician for dermatitis. The scurvy caused fractures of the femur, irritability and failure to thrive (7).

Aside from the cases reported from ingestion of a soy beverage, rice beverage or almond beverage, attention should be given to combinations of different seeds, such as rice, almonds, quinoa, oats, coconut, etc. (Supplemental Table VI [<http://www.xxxxxxxx>]). In fact, it seems there are now published cases of children who consumed these beverages exclusively in the first months of life. For instance, the case of a 2.5-month-old infant who consumed a beverage containing chestnuts, soy, almonds and nuts, and developed malnutrition with hypotonia and somnolence with severe hyponatremia and hypopotasemia (33).

This study has a number of limitations. Despite including 164 types of PBBs in our country, new products are continually appearing. The nutritional information extracted from the labels or manufacturers' webpages does not include information about nutrient bioavailability or information about nutrients covering less than 15% of daily recommended allowances.

## **SUMMARY**

Plant-based beverages are inappropriate alternatives to breast milk, infant formula or cow's milk in the first years of life as they are low in calories, protein, fat, lactose and vitamins. In the case of older children with nonexclusive feeding with PPBs, the pediatrician should be aware of the nutritional risks and limitations of these beverages in order to complement the deficiencies with other foods (43). According to the literature review, nearly exclusive consumption of any kind of PPB is associated with a specific type of disease (Table V). Thus, soy beverages non-supplemented with vitamins or minerals primarily cause rickets and failure to thrive. Rice beverages primarily cause kwashiorkor, associated with failure to thrive or anemia. Almond beverages can cause severe metabolic alkalosis, though cases of rickets, hyperoxaluria or scurvy have also been reported.

## **REFERENCES**

1. ESPGHAN Committee on Nutrition, Agostoni C, Braegger C, Decsi T, Kolacek S, Koletzko B, et al. Breast-feeding: A commentary by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr* 2009;49:112-25.
2. Koletzko B, Baker S, Cleghorn G, Neto UF, Gopalan S, Hernell O, et al. Global Standard for the Composition of Infant Formula: Recommendations of an ESPGHAN Coordinated International Expert Group. *J Pediatr Gastroenterol Nutr* 2005;41:58-599.
3. Kirby M, Danner E. Nutritional deficiencies in children on restricted diets. *Pediatr Clin North Am* 2009;56:1085-103.
4. Singhal S, Baker RD, Baker SS. A comparison of the nutritional value of cow's milk and non-dairy beverages. *J Pediatr Gastroenterol Nutr* 2017;64(5):799-805.
5. Le Louer B, Lemale J, Garcette K, Orzechowski C, Chalvon A, Girardet JP, et al. Severe nutritional deficiencies in young infants with inappropriate plant milk consumption. *Arch Pediatr* 2014;21:483-8.
6. Carvalho NF, Kenney RD, Carrington PH, Hall DE. Severe nutritional deficiencies in toddlers resulting from health food milk alternatives. *Pediatrics* 2001;107:E46.
7. Vitoria I, López B, Gómez J, Torres C, Guasp M, Calvo I, et al. Improper use of a plant-based vitamin C-deficient beverage causes scurvy in an infant. *Pediatrics* 2016;137:e20152781.
8. Jensen RHG. *Handbook of milk composition*. New York: Academic Press; 1995.
9. Dietary protein quality evaluation in human nutrition: Report of an FAO Expert Consultation. *FAO Food and Nutrition*, 31 March-2 April, 2011.
10. Rutherford SM, Fanning AC, Miller BJ, Moughan PJ. Protein digestibility-corrected amino acid scores and digestible indispensable amino acid scores differentially describe protein quality in growing male rats. *J Nutr* 2015;145:372-9
11. Mäkinen OE, Wanhalinna V, Zannini E, Arendt EK. Foods for special dietary needs: Non-dairy plant-based milk substitutes and fermented dairy-type products. *Crit Rev Food Sci Nutr* 2016;56:339-49.
12. Regulation (EU) No. 1169/2011 of the European Parliament and the Council of 25 October 2011 on the provision of food information to consumers, amending Regulations (EC) No. 1924/2006 and (EC) No. 1925/2006 of the European Parliament and of the Council, and repealing Commission Directive 87/250/EEC, Council Directive 90/496/EEC, Commission Directive 1999/10/EC, Directive 2000/13/EC of the European

Parliament and of the Council, Commission Directives 2002/67/EC and 2008/5/EC and Commission Regulation (EC) No 608/2004. Available at: <http://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:32011R1169&from=ES>.

13. EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies). Scientific opinion on the essential composition of infant and follow-on formulae. *EFSA J* 2014;12:3760(106). DOI: 10.2903/j.efsa.2014.3760.
14. Sandberg AS, Carlsson NG, Svanberg U. Effects of inositol tri-, tetra-, penta-, and hexaphosphates on in vitro estimation of iron availability. *J Food Sci* 2006;54:159-61.
15. Sethi S, Tyagi SK, Anurag RK. Plant-based milk alternatives an emerging segment of functional beverages: A review. *J Food Sci Technol* 2016;53:3408-23.
16. Fox AT, Du Toit G, Lang A, Lack G. Food allergy as a risk factor for nutritional rickets. *Pediatr Allergy Immunol* 2004;15:566-9.
17. Imataka G, Mikami T, Yamanouchi H, Kano K, Eguchi M. Vitamin D deficiency rickets due to soybean milk. *J Paediatr Child Health* 2004;40:154-5.
18. Bhatia J, Greer F, Committee on Nutrition. American Academy of Pediatrics. Use of soy protein-based formulas in infant feeding. *Pediatrics* 2008;121:1062-8.
19. Zhao Y, Martin BR, Weaver CM. Calcium bioavailability of calcium carbonate fortified soy milk is equivalent to cow's milk in young women. *J Nutr* 2005;135:2379-82.
20. Pathomrungsinyounggul P, Grandison AS, Lewis MJ. Effects of calcium chloride and sodium hexametaphosphate on certain chemical and physical properties of soymilk. *J Food Sci* 2007;72:E428-34.
21. Heaney RP, Rafferty K. The settling problem in calcium-fortified soy bean drinks. *J Am Diet Assoc* 2006;106:1753.
22. Dagnelie PC, Vergote FJ, Van Staveren WA, Van den Berg H, Dingjan PG, Hautvast JG. High prevalence of rickets in infants on macrobiotic diets. *Am J Clin Nutr* 1990;51:202-8.
23. Armas LA, Hollis BW, Heaney RP. Vitamin D<sub>2</sub> is much less effective than vitamin D<sub>3</sub> in humans. *J Clin Endocrinol Metab* 2004;89:5387-91.
24. Massa G, Vanoppen A, Gillis P, Aerssens P, Alliet P, Raes M. Protein malnutrition due to replacement of milk by rice drink. *Eur J Pediatr* 2001;160:382-4.

25. Liu T, Howard RM, Mancini AJ, Weston WL, Paller AS, Drolet BA, et al. Kwashiorkor in the United States: Fad diets, perceived and true milk allergy, and nutritional ignorance. *Arch Dermatol* 2001;137:630-6.
26. Novembre E, Leo G, Cianferoni A, Bernardini R, Pucci N, Vierucci A. Severe hypoproteinemia in infant with AD. *Allergy* 2003;58:88-9.
27. Kuhl J, Davis MD, Kalaaji AN, Kamath PS, Hand JL, Peine CJ. Skin signs as the presenting manifestation of severe nutritional deficiency: Report of 2 cases. *Arch Dermatol* 2004;140:521-4.
28. Katz KA, Mahlberg MJ, Honig PJ, Yan AC. Rice nightmare: Kwashiorkor in 2 Philadelphia-area infants fed Rice Dream beverage. *J Am Acad Dermatol* 2005;52(5 Suppl 1):S69-72.
29. Barreto-Chang OL, Pearson D, Shepard WE, Longhurst CA, Greene A. Vitamin D-deficient rickets in a child with cow's milk allergy. *Nutr Clin Pract* 2010;25:394-8.
30. Tierney EP, Sage RJ, Shwayder T. Kwashiorkor from a severe dietary restriction in an 8-month infant in suburban Detroit, Michigan: Case report and review of the literature. *Int J Dermatol* 2010;49(5):500-6.
31. Diamanti A, Pedicelli S, D'Argenio P, Panetta F, Alterio A, Torre G. Iatrogenic Kwashiorkor in three infants on a diet of rice beverages. *Pediatr Allergy Immunol* 2011;22(8):878-9.
32. Keller MD, Shuker M, Heimall J, Cianferoni A. Severe malnutrition resulting from use of rice milk in food elimination diets for atopic dermatitis. *Isr Med Assoc J* 2012;14:40-2.
33. Fourreau D, Peretti N, Hengy B, Gillet Y, Courtil-Teysedre S, Hess L, et al. Pediatric nutrition: Severe deficiency complications by using vegetable beverages, four cases report. *Presse Med* 2013;42:e37-43.
34. Mori F, Serranti D, Barni S, Pucci N, Rossi ME, De Martino M, et al. A kwashiorkor case due to the use of an exclusive rice milk diet to treat atopic dermatitis. *Nutr J* 2015;14:83.
35. Michaelsen KF, Greer FR. Protein needs early in life and long-term health. *Am J Clin Nutr* 2014;99:718S-22S.
36. Kanaka C, Schütz B, Zuppinger KA. Risks of alternative nutrition in infancy: A case report of severe iodine and carnitine deficiency. *Eur J Pediatr* 1992;151:786-8.

37. Mesa O, González JL, García Nieto V, Romero S, Marrero C. Alcalosis metabólica de origen dietético en un lactante. *An Pediatr (Barc)* 2009;70:370-3.
38. Doron D, Hershkop K, Granot E. Nutritional deficits resulting from an almond-based infant diet. *Clin Nutr* 2001;20:259-61.
39. Ellis D, Lieb J. Hyperoxaluria and genitourinary disorders in children ingesting almond milk products. *J Pediatr* 2015;167:1155-8.
40. Reznik VM, Griswold WR, Mendoza SA, McNeal RM. Neo-Mull-Soy metabolic alkalosis: A model of Bartter's syndrome? *Pediatrics* 1980;66:784-6.
41. Rodríguez-Soriano J, Valo A, Castillo G, Oliveros R, Cea JM, Balzategui MJ. Biochemical features of dietary chloride deficiency syndrome: A comparative study of 30 cases. *J Pediatr* 1983;103:209-14.
42. Avis de l'ANSES relatif a l'adaptation d'une boisson instantanée aux amandes a l'alimentation d'un enfant de douze mois, en termes de composition et de conditions d'emploi. Available at: <http://www.anses.fr/Documents/NUT2011sa0073.pdf>.
43. Thorning TK, Raben A, Tholstrup T, Soedamah-Muthu SS, Givens I, Astrup A. Milk and dairy products: Good or bad for human health? An assessment of the totality of scientific evidence. *Food Nutr Res* 2016;60:32527.

**Nutrición  
Hospitalaria**

**Table I. Nutritional composition of plant-based beverages, cow's milk and recommended composition of soy formula and infant formula**

Plant-based	Number of brands	Kcal/100 ml (range)	Carbohydrates g/100 ml (range)	Fats g/100 ml (range)	Proteins g/100 ml (range)	Protein energy/Total energy (%)	Supplemented (n)			
							Calcium	Vitamin D	Other minerals	Other vitamins
Soy	54	46.7 ± 13.1 (27-80.7)	4.3 ± 2.9 (0.1-11.8)	1.8 ± 0.4 (0.9-2.9)	3.1 ± 0.4 (2.1-3.8)	28.3 ± 6.9 (20.0-42.0)	43	23	2	25
Rice	24	56.8 ± 6.3 (47-68)	11.5 ± 1.5 (9.4-14.2)	0.9 ± 0.1 (0.8-2)	0.3 ± 0.2 (0.1-0.8)	2.4 ± 1.4 (0.8-6.4)	8	5	0	2
Almond	22	40.2 ± 14.3 (25-74)	4.4 ± 2.5 (0.1-10.5)	2.0 ± 0.6 (1.1-2.8)	0.8 ± 0.3 (0.3-1.6)	8.0 ± 2.5 (4.3-12.3)	11	5	1	5
Oat	31	45.3 ± 8.3 (30-60)	7.5 ± 1.7 (4.4-11)	1.1 ± 0.4 (0.5-1.8)	0.9 ± 0.3 (0.3-1.4)	8.3 ± 3.1 (3.3-13.7)	16	9	0	5
Coconut	6	33.8 ± 15.1 (15-53)	4.3 ± 2.5 (2-9.1)	1.8 ± 1.1 (0.1-3.3)	0.2 ± 0.2 (0.1-0.5)	3.0 ± 1.9 (1.2-6.3)	2	2	0	2
Miscellaneous	12	48.1 ± 10.2 (29-65)	6.0 ± 3.0 (2.2-10.5)	2.2 ± 0.8 (1-3.6)	0.7 ± 0.3 (0.4-1.1)	6.4 ± 2.6 (5.5-7.1)	1	2	0	1
Mixed	15	61.3 ± 13.0	10.7 ± 2.7	1.6 ± 0.7	0.6 ± 0.4	4.5 ± 3.3	1	0	0	0

		(36-90)	(5.2-14.5)	(0.8-3.1)	(0.3-1.8)	(2.5-10.4)				
--	--	---------	------------	-----------	-----------	------------	--	--	--	--

	Kcal/100 ml	Carbohydrates g/100 ml	Fats g/100 ml	Proteins g/100 ml	Protein energy/Total energy (%)
Cow's milk (8)	60-70	4.5-5.0	3.5-4.0	3.2	18.2-21.3
Soy infant formula (2)	60-70	5.8-9.1	2.8-3.9	1.5-1.9	7.4-10.6
Infant formula (2)	60-70	5.8-9.1	2.8-3.9	1.2-1.9	7.4-10.6

**Table II. Published clinical cases of nutritional problems associated with soy beverages consumed by infants and toddlers**

Authors Year	Reasons for introduction of soy beverage	Age of introduction of soy beverage (age of diagnosis)	Characteristics of feeding	Daily intake	Laboratory findings	Diagnosis
Carvalho NF et al (6)	Taste preference Breastfeeding without	10 months (17 months)	Soy beverage, vegetables, fruits	900 ml	Ca 2.22 mmol/l P 0.55 mmol/l AP 1879 U/l	Rickets Failure to thrive

2001	vitamin D supplement				VitD 19.2 nmol/l PTH 12.1 pmol/l	
Fox AT et al. (16) 2004	Breastfeeding without vitamin D supplement Urticaria with infant formula at 6 months	6 months (14 months)	Breastfeeding, soy beverage, vegetables, fruits	--	Ca 1.71 mmol/l P 1.06 mmol/l AP 2054 U/l VitD 15 nmol/l PTH 44.1 pmol/l	Rickets Failure to thrive Ferropenic anemia
Imataka G et al. (17) 2004	Eczema at 3 weeks Parental decision	1 month (5 months)	Soy beverage Calcium: 28.9 mg/l No vitamin D	--	Ca 1.32 mmol/l P 1.6 mmol/l AP 2303 U/l VitD 19.9 nmol/l PTH 254 pmol/l	Hypocalcemic tetany Rickets Failure to thrive

AP: Alkaline phosphatase; Ca: Calcium; P: Phosphorus; PTH: Parathyroid hormone; VitD: 25-OH-vitamin D<sub>3</sub>.



**Table III. Published clinical cases of nutritional problems associated with rice beverages consumed by infants and toddlers**

Authors Year	Reasons for introduction of rice beverage	Age of introduction of rice beverage (age of diagnosis)	Characteristics of feeding	Daily intake	Laboratory findings	Diagnosis
Massa G et al. (24) 2001	Dermatitis unimproved with a soy formula (homeopathic physician)	16 weeks (33 weeks)	Rice beverage Fruits, vegetables	RB: 1.0-1.38 l	Alb 26 g/l	Kwashiorkor
Carvalho NF et al. (6) 2001	Eczema and perceived milk intolerance	13-15 months? (22 months)	Rice beverage Vegetables	RB: 1.5 l 0.3 g prot/kg/d 79 kcal/kg/d	Alb 10 g/l Zinc 32.2 µg/dl	Kwashiorkor
Liu T (25) 2001	Perceived intolerance of formula	? (4 months)	Rice beverage Vitamins	--	Alb 14 g/l TProt 29 g/l Zinc 22 µg/dl	Kwashiorkor

Novembre E et al. (26) 2003	Atopic dermatitis (naturopathic doctor)	5 months (6 months)	Rice beverage, rice cream, vegetables, fruits	RB: 660 ml 0.5 g prot/kg/d 86 kcal/kg/d	Alb 14 g/dl TProt 28 g/l	Kwashiorkor
Kuhl J et al. (27) 2004	Atopic dermatitis positive RAST to multiple foods	14 months (17 months)	Rice beverage, 1-2 tablespoons of baby food	5 g prot/d 600 kcal/d	Alb 12 g/l TProt 35 g/l Zinc 27 µg/dl	Failure to thrive Kwashiorkor
Katz K et al. (28) 2005	Breastfed 8 m Rejection of infant formula	8 months (14 months)	Rice beverage, meat, vegetables	--	Alb 14 g/l TProt 36 g/l Zinc 28 µg/dl	Kwashiorkor
Katz K et al. (28) 2005	Rejection of infant formula	2 months (7 months)	Rice beverage, baby food, iron supplementation	--	Alb 15 g/l TProt 34 g/l Zinc 31 µg/dl	Failure to thrive Kwashiorkor
Barreto-Chang OL et al. (29) 2010	Cow's milk allergy	13 months (16 months)	Rice milk (0.4 g proteins/100 ml)	--	VitD 9 nmol/l PTH 20.4 pmol/l	Failure to thrive Rickets
Tierney E et al. (30)	Scalp rash	4 months (8 months)	Rice milk, bananas, sweet potatoes	--	Alb 20 g/l TProt 37 g/l	Kwashiorkor

2010					Zinc 91.5 µg/dl	
Diamanti A et al. (31) 2011	Cow's milk allergy (3 cases)	3 months (4 months) 1.5 months (4 months) 3 months (5 months)	Rice beverage	--	Alb < 20 g/l TProt < 40 g/l	Kwashiorkor
Keller MD et al. (32) 2012	Eczema. Allergy to cow's milk, soy, egg, peanut, etc.	13 months (19 months)	Rice beverage, rice, potatoes, carrots	--	Alb 16 g/l TProt 33 g/l	Kwashiorkor
Keller MD et al. (32) 2012	Suspected cow's milk allergy (eczema, vomiting)	12 months (16 months)	Rice beverage Lentils, chick-peas, olives	--	Alb 12 g/l Hb 7 g/dl	Kwashiorkor Anemia
Fourreau D et al. (33) 2013	Suspected cow's milk allergy (naturopathic doctor)	7 months (9 months)	Rice beverage (0.1 g prot/100 ml), fruits, vegetables	RB:800-900 ml	Alb 7 g/l Hb 10 g/dl	Kwashiorkor Anemia
Fourreau D	Suspected cow's	13 months	Rice beverage	RB: 300 ml	Alb 7 g/l	Failure to

et al. (33) 2013	milk allergy (parental decision)	(14.5 months)			Hb 3.5 g/dl Vit B <sub>12</sub> 143 ng/l	thrive anemia
Le Louer B et al. (5) 2014	Vomiting	2 months (4.5 months)	Rice beverage	--	Hb 5.7 g/dl Alb 1.8 g/dl Zinc 3.5 µmol/l	Failure to thrive anemia
Le Louer B et al. (5) 2014	Eczema	1 months (7 months)	Rice beverage	--	Hb 8.7 g/dl Alb 1.98 g/dl Zinc 3.9 µmol/l	Failure to thrive Kwashiorkor Anemia
Mori et al. (34) 2015	Atopic dermatitis (naturopathic doctor)	4 months (6 months)	Rice milk, fruits, rice poultry and vegetable broth.	--	Alb 13 g/l TProt 30 g/l Hb 5.7 g/dl	Kwashiorkor Anemia

Alb: Albumin; Hb: Hemoglobin; PTH: Parathyroid hormone; RB: rice beverage; TProt: total protein; VitD: 25-OH-vitamin D<sub>3</sub>.

**Table IV. Published clinical cases of nutritional problems associated with almond beverages consumed by children**

Authors Year	Reasons for introduction of almond beverage	Age of introduction of almond beverage (Age of diagnosis)	Characteristics of feeding	Daily intake	Laboratory findings	Diagnosis
Kanaka C et al. (36) 1992	Eczematous reaction to cow's milk formula (maternal decision)	2.5 months (7.5 months)	Self-prepared extract of almonds Cereals Fruits	98% DRI proteins 54% DRI energy	TSH 378 $\mu$ IU/ml Iodine 47 nmol/l Free carnitine 12 $\mu$ mol/l	Failure to thrive Iodine and carnitine deficiency
Mesa O et al. (37) 2009	--	Birth (31 days)	Almond beverage	--	Cl <sup>-</sup> 94 mmol/l Na <sup>+</sup> 136 mmol/l K <sup>+</sup> 3 mmol/l CO <sub>3</sub> H <sup>-</sup> 40.3 mmol/l	Dehydration Metabolic alkalosis
Mesa O et al. (37) 2009	--	Birth (4 months)	Almond beverage	--	Cl <sup>-</sup> 74 mmol/l Na <sup>+</sup> 124 mmol/l K <sup>+</sup> 2.2 mmol/l CO <sub>3</sub> H <sup>-</sup> 49.8 mmol/l	Metabolic alkalosis

Fourreau D et al. (33) 2013	Suspected gastro-esophageal reflux	12 months (13 months)	Almond beverage (17 mg sodium/100 ml; 24 mg chloride/100 ml) Yogurt Vegetables	840 ml	Cl <sup>-</sup> 69 mmol/l Na <sup>+</sup> 127 mmol/l K <sup>+</sup> 1.9 mmol/l CO <sub>3</sub> H <sup>-</sup> 48 mmol/l	Metabolic alkalosis
Doron D et al. (38) 2013	Diarrhea and vomiting attributed by the mother to cow's milk protein allergy	4 months (6 months)	Almond-based home made "formula" (Almond 10 g/water 100 ml)	1,000 ml	Ca 1.4 mmol/l P 1.2 mmol/l AP 818 U/l vitD < 12 nmol/l PTH 30.3 pmol/l Hb 7.7 g/dl	Failure to thrive Rickets Anemia
Doron D et al. (38) 2013	Rash	4-5 months (8 months)	Almond-based and honey home made "formula" (20 gr almonds / 100 ml water)	600 mL	Alb 20 g/l TProt 36 g/l	Kwashiorkor
Le Louer B et al. (5)	Gastro-esophageal reflux, eczema	3.5 months (5 months)	Almond and chestnut beverage	-	Alb 19.5 g/l Ca 0.64 mmol/l Zinc 7 μmol/l	Hypocalcemic tetany Malnutrition

2014						
Le Louer B et al. (5) 2014	Parental decision	8.5 months (16.5 months)	Almond and walnuts beverage	--	VitD < 12.5 nmol/l Ca 2.32 mmol/l P 1.71 mmol/l PTH 8.8 pmol/l	Rickets
Ellis D et al. (39) 2015	Tourette syndrome (1 case) Lactose intolerance (2 cases)	3 years 9 years 10 years	Almond milk and varied diet	700-1,000 ml	Urine oxalate 53.5, 81.5 and 97.9 mg/1.73 m <sup>2</sup> /d (27.6- 35.4)	Hyperoxaluria Hematuria (2 cases) Kidney stones (1 case)
Vitoria I et al. (7) 2016	Medical indication (atopic dermatitis)	2.5 months (11 months)	Almond milk Almond flour Cereals	840 ml/d	Ascorbic acid < 10 µmol/l VitD 31 nmol/l	Scurvy

Alb: Albumin; Ca: Calcium; Hb: Hemoglobin; P: Phosphorus; TProt: total protein; VitD: 25-OH-vitamin D<sub>3</sub>.

**Table V. Types of plant-based beverages in infants and risk of nutritional disease**

Plant-based beverage	Primary associated nutritional disease	Other associated disease
Soy	Rickets	Failure to thrive
Rice	Kwashiorkor	Failure to thrive Anemia
Almond	Metabolic alkalosis	Rickets Hyperoxaluria Scurvy

**Supplemental Table I. Content in energy, carbohydrates, sugars, fats, proteins, calcium and vitamin D of 54 soy beverages**



Brand	Characteristics	Kcal	Carbohydrates (g/100 ml)	Sugars (g/100 ml)	Fats (g/100 ml)	Saturated/Monounsaturated/Polyunsaturated (g/100 ml)	Proteins (g/100 ml)	Fiber (g/100 ml)	Salt (g/100 ml)	Calcium (mg/100 ml)	Vitamin D (µg/100 ml)	Others minerals	Others vitamins
Alpro		39	2.8	2.8	1.8	0.3/-/-	3	0	0.08	160	0.75		B <sub>2</sub> : 0.21** B <sub>12</sub> : 0.38*
Alpro	Light	27	1.8	1.6	1.1	0.2/-/-	2.1	0.6	0.08	160	0.75		B <sub>2</sub> : 0.21** B <sub>12</sub> : 0.38*
Alpro	Chocolate	61	7.8	7.5	1.7	0.4/-/-	3.1	0.9	0.14	120	0.75		B <sub>2</sub> : 0.21** B <sub>12</sub> : 0.38*
Alpro	Vanilla	54	6.5	6.4	1.7	0.3/-/-	3	0.5	0.14	120	0.75		B <sub>2</sub> : 0.21**

													B <sub>12</sub> : 0.38*
Biocesta	Calcium	42	2.5	2.4	1.9	0.3/0.4/1.2	3.5	0.5	0.07	120	0.75		B <sub>2</sub> : 0.21** B <sub>12</sub> : 0.38*
Bio Gerblé		42	1.6	0.7	2.2	0.3/0.4/1.2	3.8	0.5	0.03				
Bio Gerblé	Calcium	45	2.9	2.1	1.9	0.4/0.4/1.1	3.7		0.10	125	0.9		
Bio Gerblé	Calcium hazelnut flavor	47	4.3	4	1.8	0.3/0.4/1.0	3.5		0.12	125	1	Fe: 2.5** Zn: 2.3**	B <sub>12</sub> : 0.40*
Bjorg	Calcium	52	3.5	3	2	0.4/-/-	3.8	0.5	0.08	120			
Carrefour	Calcium	43	3.7	3	1.7	0.3/-/-	3.1	0.1	0.10	120	0.75		A: 120*
Carrefour	Calcium light	32	3.5	2.9	1.1	0.2/-/-	2.1		0.10	120			
Carrefour	Chocolate	70	10	9.4	1.8	0.4/-/-	3.3		0.10	120	0.75		A: 120*
Carrefour	Vanilla	62	8.2	7.6	1.8	0.3/-/-	3.3		0.10	120			B <sub>2</sub> : 0.21**

														B <sub>12</sub> : 0.40*
Consum		32.3	1	0.7	1.7	0.3/-/-	3.1	0.2	0.06					
Consum	calcium	43.1	3.7	3	1.7	0.3/-/-	3.1	0.3	0.10					A: 120*
Consum	cappuccino	54	6.2	5.8	1.7	0.3/0.4/1.0	3.1	0.9	0.10	120	0.75			A: 120*
Consum	chocolate	80.7	11.8	11	1.9	0.3/-/-	3.8	0.6	0.10					
Consum	Vanilla	65.3	9.2	7.7	1.7	0.3/-/-	3.3	< 0.5	0.13	120				B <sub>2</sub> : 0.21** B <sub>12</sub> : 0.38*
Dia		43	3.7	3	1.7	0.3/-/-	3.1		0.10	120	0.75			A: 120*
Dia	Light	32	3	2.8	1.2	0.2/-/-	2.1		0.13	120	0.75			A: 120*
Don Simon		48	4.4	4	1.9	0.3/0.4/1.2	3.2	0.3	0.01	120	0.75			B <sub>2</sub> : 0.21** B <sub>12</sub> : 0.38*
Don Simon	Light	28	2.1	1.8	1.1	0.2/0.3/0.7	2.1	0.6	0.15	160	0.75			
Frias	Calcium	33	3.5	2.9	1.2	0.2/-/-	2.1	0.2	0.10	120	0.75			
Frias		32	3.5	2.9	1.7	0.3/-/-	3.1	0.2	0.06					

Granovita		36	0.6	0.1	2.1	0.4/-/-	3.6	1	0.04				
Hacendado		30	0.6	0.6	1.7	0.3/0.3/1.0	3.1	< 0.5	0.10				
Hacendado	Calcium	44	2.8	2.8	2	0.3/0.4/1.2	3.4	0.5	0.15	120			B <sub>2</sub> : 0.21** B <sub>12</sub> : 0.37*
Hacendado	Coffee	72	11	9.5	1.7	0.3/0.3/1.0	3	0.5	0.19	120			B <sub>2</sub> : 0.21** B <sub>12</sub> : 0.37*
Hacendado	Chocolate	69	9	9	2	0.3/0.4/1.2	3.4	0.8	0.15	120			B <sub>2</sub> : 0.21** B <sub>12</sub> : 0.37*
Hacendado	Vanilla	65	9.5	9.5	1.7	0.3/0.3/1.0	3	< 0.5	0.06	120			
Kaiku Sojavit		40	2.8	2.8	1.7	0.3/0.3/1.0	3.4	0.6	< 0.10	120			
Milbona	Calcium	43	3.7	3	1.7	0.3/-/-	3.1	0.3	0.10	120	0.75		A: 120*
Naturgreen Nature		45	0.9	0.8	2.9	0.5/-/-	3.6		0.10				

Naturgreen	Cinnamon and lemon	61	7.3	6.8	1.9	0.3/0.4/1.2	3.4	1.2	0.05				
Naturgreen	Calcium	49	4.1	3	2.9	0.4/-/-	3.4		0.22	120			
Provamel Calcimel		45	2.5	2.5	2.2	0.4/0.5/1.3	3.7	0.6	0.15	140			
Provamel Calcimel Bio		45	2.4	2.4	2.1	0.4/0.4/1.3	3.7	0.6	0.15	120			
Provamel Natural		35	0.1	0.1	2.1	0.4/0.4/1.3	3.7	0.6	0.04				
Provamel	Chocolate	67	8.1	7.9	2.2	0.6/-/-	3.4	1.1	0.14				
Shoyce		50	4.5	2.7	2.0	0.3/-/-	2.4	0.3	0.20	120	0.75		A: 120**
Shoyce	Light	36	3.3	1.8	1.4	0.2/-/-	2.4	0.2	0.18	120		Fe: 2.1**	A: 120* B <sub>12</sub> : 0.38*
Shoyce junior		53	4.7	2.6	2.2	0.3/-/-	3	0.9	0.02	120			
Sojade		37	0.45	0.45	2.1	0.3/0.5/1.3	3.8	0.5	0.08				
Sojade	Calcium	45	2.5	2.5	2.1	0.4/0.5/1.2	3.7	0.5	0.10	120			
Sojasun	Calcium	43	2.3	2.3	2	0.3/0.5/1.2	3.6		0.10	144	0.75		

Sojasun	Chocolate	71	9.4	9	2	0.4/0.5/1.1	3.4		0.15	120			
Sojasun	Vanilla	49	4.2	4.2	2	0.3/0.5/1.2	3.4		0.17	120			
Sorianatural	Calcium	33.7	2.7	1.5	0.9	0.1/0.1/0.4	3.4	0.8	0.20	120			
Special Line	Calcium	49	4	3.2	2	0.3/-/-	3.6	0.3	0.10	120			A: 120*
Vivesoy		41	2.9	2.6	1.7	0.3/0.4/1.0	3.1	0.7	0.10	120	0.75		
Vivesoy	Chocolate	53	6.1	5.8	1.6	0.3/0.4/0.9	3	1.1	0.10	120	0.75		A: 120*
Vivesoy	Light	27	1.8	1.7	1.1	0.2/0.3/0.7	2.2	0.5	0.09	120	0.75		A: 120*
Vivesoy	Vanilla	37	2.3	2	1.6	0.3/0.4/1.0	3	0.7	0.08	120	0.75		A: 120*
Yo soy	Calcium	38	4	1.9	1	0.2/0.2/0.6	3	0.3	0.05	120	0.75		

\*: µg/100 ml; \*\*: mg/100 ml.

**Nutrición  
Hospitalaria**

**Supplemental Table II. Content in energy, carbohydrates, sugars, fats, proteins, calcium and vitamin D of 24 rice beverages**

Brand	Characteristics	Kcal	Carbohydrates (g/100 ml)	Sugars (g/100 ml)	Fats (g/100 ml)	Saturated/Monounsaturated/Polyunsaturated (g/100 ml)	Proteins (g/100 ml)	Fiber (g/100 ml)	Salt (g/100 ml)	Calcium (mg/100 ml)	Vitamin D (µg/100 ml)	Others vitamins
Alpro		49	9.8	6.7	1	0.1/-/-	0.1	0	0.08	120	0.75	B <sub>12</sub> : 0.38*
Amandín Organic		62	13	5.3	0.8	0.1/0.6/0.1	0.6	0.5	0.06			
Bio Gerblé		63	13	7.1	1	0.1/0.7/0.1	0.4	0.1	0.10			
Carrefour Rice	Calcium	54	11	6	1	< 0.1/-/-	0.4		0.10	120		
Finestra sul cielo		63	13.3	6.6	1	0.1/0.3/0.6	0.2	0.2	0.10			
Joya		57	12	2.8	0.8	0.1/0.2/0.5	0.2	0.3	0.10			
Lima		52	10.8	7.9	0.9	0.1/0.7/0.1	0.1	0.1	0.10			
Milbona		47	9.7	6.5	0.9	0.1/-/-	0.1		0.06	120	0.75	
Monsoy		55	11	8	1	0.1/0.3/0.6	0.3	0.2	0.07			

Probio Rice	Vanilla	55	10.5	7	1.1	0.1/-/-	0.4	0.7	0.10			
Rice Dream		47	9.4	4	1	0.1/-/-	0.1	0.1	0.03			
Rice Dream	Calcium	63	13.3	6.6	1	0.1/-/-	0.2	0.2	0.12	120	0.75	
Sorianatural		50	9.4	4.3	1.0	0.1/0.6/0.2	0.8	0.12	0.25			
Scotti		61	13.1	8.5	0.9	0.2/-/-	0.2		0.10			
Scotti	Cocoa	68	14.2	10.4	0.8	0.3/-/-	0.5		0.10			
Scotti	Calcium	61	13.1	8.5	0.9	0.2/-/-	0.2		0.10	120		
Special Line Rice		67	14	11	0.8	0.0/-/-	0.7	0.12	0.09	120	0.75	A: 120*
Special Line Rice	Cocoa	55	12	4.6	0.8	0.1/-/-	0.5		0.12			
Vegetalia		54	11	7.1	1	0.1/0.3/0.6	0.3	0.2	< 0.10			
Vitariz Alinor		55	10.5	7	1.1	0.1/-/-	0.7	0.4	0.10			
Vitariz Alinor	Cocoa	67	11	8	2.0	0.7/-/-	0.7	1	0.10			
Vitariz Alinor	Calcium	55	10.5	7	1.1	0.1/-/-	0.4	0.7	0.10	120		
Vivesoy		49	9.7	7.1	1	0.1/-/-	0.2	0.06	0.06	60	0.38	
Yosoy		55	11.2	4.7	1	0.1/0.3/0.6	0.3	0.1	0.07			



# Nutrición Hospitalaria

\*:  $\mu\text{g}/100\text{ ml}$ .

**Supplemental Table III. Content in energy, carbohydrates, sugars, lipids, proteins, calcium and vitamin D of 22 almond beverages**

Brand	Characteristics	Kcal	Carbohydrates (g/100 ml)	Sugars (g/100 ml)	Fats (g/100 ml)	Saturated/Monounsaturated/Polyunsaturated (g/100 ml)	Proteins (g/100 ml)	Fiber (g/100 ml)	Salt (g/100 ml) (& Na)	Calcium (mg/100 ml)	Vitamin D (µg/100 ml)	Others vitamins and minerals
Almendrino		74	10.5	10.5	2.8	0.4/1.6/0.6	1.6	0.31	< 1.0	14		A: 0.96* B <sub>12</sub> : 88*
Almendrola		58	8.3	8	2.1	0.2/1.4/0.5	0.8	0.3	0.04			
Almendrola	Agave syrup	42	4	3	2.4	0.2/1.7/0.5	1	0.1	0.05			
Almond Breeze		24	2.9	2.8	1.1	< 0.1/-/-	0.5		0.13	120		
Almond Dream original	Calcium	43	4	3.5	1.4	0.1/-/-	0.8	0.9	0.04 <sup>&amp;</sup>	120	0.75	B <sub>12</sub> : 0.38*
Alpro	Sugar free	13	0.1	0.1	1.1	0.1/-/-	0.4	0.4	0.13	120	0.75	B <sub>12</sub> : 0.38* B <sub>2</sub> : 0.21** E: 18**
Carrefour	Calcium	23	3	3	1.1	< 0.1/-/-	< 0.5		0.12	120		
Diemilk		50	6.3	3.6	2.2	0.6/1.2/0.4	1.1	0.4	0.05 <sup>&amp;</sup>			
DiemilkNature		39	2.7	0.1	2.6	0.3/1.8/0.5	1.1	0.4	0.02 <sup>&amp;</sup>			

EcoMil Nature		37	1.9	< 0.3	2.7	0.8/-/-	1		0.10			
EcoMil Nature	Calcium	37	1.9	< 0.3	2.7	0.8/-/-	1		0.10	120		
EcoMil Original	Calcium	46	5.4	3.8	2.1	0.2/1.4/0.5	0.9	0.8	0.05	120		
EcoMil Original	Cocoa	46	6.1	4.9	2.1	0.6/-/-	0.5		0.10			
EcoMil Original	Vanilla	35	4.5	3.4	1.7	0.4/-/-	0.4		0.10			
EcoMil Mandorla Bio		46	5.4	3.8	2.1	0.2/1.4/0.5	0.9	0.8	0.25			
Ecomil	Muesli	46	5.9	3.7	2.2	0.6/1.2/0.4	0.6	0.3	0.10	120		
Gerblé	Light	25	2.3	2	1.6	0.2/0.8/0.6	0.3	0.3	0.10	120	15	A: 37.5* E: 3.1**
Milbona		29	3.5	3.3	1.4	0.3/-/-	0.5	< 0.5	0.12	120	0.75	B <sub>12</sub> : 0.38* A: 120*
Monsoy		60	9.5	8.5	2.1	0.1/1.5/0.5	0.8	0.2	0.02			
Santiveri		60	9.5	8.5	2.1	0.1/-/-	0.8	0.2	0.10			
Special Line		42	5.3	4.1	1.8	0.3/-/-	0.8	0.6	0.20			
Vivesoy		27	3.4	3.3	1.2	0.1/-/-	0.5	0.3	0.12	60	0.38	Fe: 1.1**

\*: µg/100 ml; \*\*: mg/100 l.


Supplemental Table IV. Content in energy, carbohydrates, sugars, fats, proteins, calcium and vitamin D of 31 oat beverages

Brand	Characteristics	Kcal	Carbohydrates (g/100 ml)	Sugars (g/100 ml)	Fats (g/100 ml)	Saturated/Monounsaturated/Polyunsaturated (g/100 ml)	Proteins (g/100 ml)	Fiber (g/100 ml)	Salt (g/100 ml) & sodium	Calcium (mg/100 ml)	Vitamin D (µg/100 ml)	Others minerals	Others vitamins
Alpro		46	8	6	0.8	0.1/-/-	1.2	0.8	0.01	120	0.75		
Amandin	Calcium	58	10.5	8.6	1.6	0.2/1.0/0.3	0.53	0.33	0.12	120			
Amandin	Cocoa	58	10	8.7	1.4	0.2/0.8/0.3	0.7	0.7	0.13				
Biocesta		46	8	5.5	0.9	0.2/0.3/0.4	1.2	0.8	0.05				
Cereal Bio		59	10.5	8.6	1.6	0.2/-/-	0.5	0.3	0.10				
Finestra sul cielo		38.5	6.1	4	0.95	0.1/0.3/0.5	0.99	0.86					
Finestra sul cielo	Calcium	33	4.8	2.4	1.2	0.2/0.3/0.7	0.6	0.4	0.10	120			
Gerblé	Calcium	43	7	5	0.9	0.2/0.3/0.4	1.3	0.6	0.10	120	0.8		
Gerblé	Light	30	5.5	3.7	0.5	0/-/-	0.5	0.9	0.09	120	0.75		

Granovita		44	7.6	4.5	1.1	0.2/0.3/0.6	0.7	0.5	0.10				
Joya		42	7.7	4.5	0.8	0.1/0.2/0.4	0.7	0.7	0.09				
Milbona		46	8.3	6.3	0.8	0.1/-/-	1.4	0.2	0.07	120	0.75		
Naturgreen		38	7	3.5	0.7	0.1/0.3/0.4	0.8	0.4	0.04 <sup>&amp;</sup>				
Naturgreen	Calcium	45	7	3.5	1.5	0.2/0.5/0.8	0.7	0.4	0.10	120			
Naturgreen	Calcium Choco	60	9.4	7.1	1.8	0.3/0.6/0.9	0.9	1.3	0.03	120			
Oat Dream	Calcium	51	8.6	4.5	1.3	0.2/-/-	0.6	1	0.04	120	0.75		B <sub>12</sub> : 0.37**
Oat Ly		50	6.6	4.1	1.5	0.2/-/-	1	0.8	0.11	120	1.5		B <sub>2</sub> : 0.21** B <sub>12</sub> : 0.38*
Oat Ly	Orange	55	11	8.5	0.5	0.1/-/-	1	0.8	0				
Oat Ly	Calcium	45	6.5	4	1.5	0.2/-/-	1	0.8	0.1	120			
Oat Ly	Vanilla	40	8	5.5	0.5	0.1/-/-	1	0.8	0.1				
Oat Ly	Chocolate	55	9.5	7	1.5	0.2/-/-	1	1	0.1	120	1.5		B <sub>2</sub> : 0.21** B <sub>12</sub> :

													0.38*
Santiveri Lactavena		52	8.9	4.8	1.5	0.2/0.5/0.8	0.7	0.1	0.05				
Santiveri Calciavena	Calcium	38	6	4.5	0.9	0.2/0.3/0.4	1.3	0.5	< 0.05	120	0.75		
Shoyce		36	4.4	2.5	1.5	0.2/-/-	0.3	1.5	0.11	120			B <sub>2</sub> : 0.21** B <sub>12</sub> : 0.38*
Sorianatural	Fiber	38.5	6.07	0.7	0.95	0.1/0.3/0.4	0.9	0.86	0.32				
Sorianatural	Calcium	39.7	6.27	0.7	0.95	0.1/0.3/0.4	1.0	0.98	0.32	120			
Special Line		36	6.5	4	0.5	0.1/-/-	1		0.10				
Special Line	Cocoa	58	10	8.7	1.4	0.2/-/-	0.7		0.13				
Vegetalia		46	8		0.9	0.2/0.3/0.4	1.2	0.8	0.05				
Vivesoy		40	6.3	4.7	0.9	0.2/-/-	1.3	0.5	0.10	60	0.39		Biotin: 3.8*
Yosoy		38	6	4.5	0.9	0.2/0.3/0.4	1.3	0.5	0.03				

Brand	Kcal	Carbohydrates (g/100 ml)	Sugars (g/100 ml)	Fats (g/100 ml)	Saturated/Mono unsaturated/Poly unsaturated (g/100 ml)	Proteins (g/100 ml)	Fiber (g/100 ml)	Salt (g/100 ml)	Calcium (mg/100 ml)	Vitamin D ( $\mu\text{g}/100\text{ ml}$ )	Others minerals	Others vitamins
-------	------	--------------------------	-------------------	-----------------	--	---------------------	------------------	-----------------	---------------------	---	-----------------	-----------------



\*:  $\mu\text{g}/100\text{ ml}$ ; \*\*:  $\text{mg}/100\text{ ml}$ .

**Supplemental Table V. Content in energy, carbohydrates, sugars, fats, proteins, calcium and vitamin D of 6 coconut beverages and 12 miscellaneous beverages**

<i>Coconut beverages</i>												
Alpro	20	2.7	1.9	0.9	0.9/-/-	0.1		0.13	120	0.75		B <sub>12</sub> : 0.38*
Bjorg	32	3.5	3.5	2	1.7/-/-	0.5	0.5	0.10				
Dr Goerg	15	3.7	3.6	0.1	< 0.1/-/-	0.1		0.06				
Ecomil	34	2	< 0.3	2.7	2.5/-/-	0.2		0.10				
Ecomil calcium	49	4.6	3.2	3.3	3.0/-/-	0.1		0.04	120	0.75		B <sub>12</sub> : 0.38*
Koko chocolat	53	9.1	7.4	1.9	1.7/0.1/0.0	0.5	0.4	0.10				
<i>Barley</i> SoriaNatural	55	9.3	6.9	1.6	0.2/0.5/0.9	0.6	0.1	0.14				
<i>Canary grass</i> Soriantural	45	7.1	3.2	1	0.1/0.4/0.4	1.1	0.3	0.12				
<i>Hazelnut</i>	29	3.2	3.2	1.6	0.2/-/-	0.4	0.3	0.14	120	0.75		B <sub>2</sub> : 0.21**




Alpro													B <sub>12</sub> : 0.38* E: 1.8**
<i>Hazelnut</i> DieMilk	55.7	6.8	5.1	2.9	0.6/2.0/0.3	0.6	0.4	0.10				0.14	
<i>Hemp</i> Ecomil	43	3.1	2.3	2.9	0.3/-/-	1		0.07					
<i>Hemp</i> Ecomil nature	40	2.2	< 0.3	2.9	0.3/-/-	1		0.07					
<i>Macadamia nut</i> Amandin	65	8.7	4.8	3.6	0.8/2.5/0.1	0.4	2.2	0.18					
<i>Millet</i> Isola-Bio	55	10.5	5.5	1.1	0.1/0.3/0.7	0.7	0.4	0.08					
<i>Quinoa</i>	43	3	2.4	2.4	0.4/-/-	1		0.11					

Ecomil												
<i>Sesame</i> Ecomil	51	6.7	3.4	2.4	0.5/1.4/0.5	0.6	0.2	0.02				
<i>Spelt</i> Special Line	47.6	9.3	8.7	1	0.12/-/-	0.6	0.3	0.05				
<i>Spelt</i> Isola-Bio	57	10.5	5.5	1.1	0.1/0.3/0.7	0.8	0.9	0.08				

\*: µg/100 ml; \*\*: mg/100 ml.

Supplemental Table VI. Content in energy, carbohydrates, sugars, fats, proteins, calcium and vitamin D of 15 mixed beverages



Brand	Kcal	Carbohydrates (g/100 ml)	Sugars (g/100 ml)	Fats (g/100 ml)	Saturated/Monounsaturated/Polyunsaturated (g/100 ml)	Proteins (g/100 ml)	Fiber (g/100 ml)	Salt (g/100 ml)	Calcium (mg/100 ml)	Vitamin D (µg/100 ml)	Others minerals and vitamins
-------	------	-----------------------------	----------------------	--------------------	---	------------------------	---------------------	--------------------	------------------------	--------------------------	------------------------------------

**Nutrición  
Hospitalaria**

<i>Rice and almond</i>											
Finestra sul cielo	90	14.5	6.7	3.1	0.3/1.6/1.2	0.8	0.6	0.07			
Joya	36	5.2	3.8	1.3	0.1/0.8/0.3	0.6	0.6	0.12			
<i>Rice and coconut</i>											
BioGerblé	58	11	5.8	1.1	0.8/0.4/0.1	0.5	0.5	0.14	120		
IsolaBio	60	13	6.5	0.8	0.6/0.1/0.1	0.5	0.5	0.08			
Special Line	62	12.3	8.9	0.9	0.3/-/-	0.3	0.1	0.08	120		
Yosoy	60	11.7	5.6	1.3	1.2/0.1/0.0	0.3	0.4	0.07			
Joya	64	11.4	5.9	1.8	1.6/0/0.1	0.3	0.1	0.09			
<i>Rice and hazelnut</i>											
Monsoy	62	12	6.3	1.3	0.1/1.0/0.2	0.7	0.7	0.03			
Finestra sul cielo	85	13.6	7	3	0.6/1.8/0.6	0.6	0.5	0.07			
Yosoy	65	12	5.6	1.5	0.1/1.1/0.3	0.5	0.5	0.06			
<i>Rice and quinoa</i>											

QuinoaReal	63	12.4	5.2	1.2	0.2/0.4/0.6	0.6	0.5	0.09			
Sorianatural	49	9.9	6.4	0.8	0.1/0.2/0.4	0.42	0.1	0.10			
<i>Oat and almond</i>											
Almendrola	58	7	5	2.4	0.3/1.4/0.7	1.8	0.7	0.05			
Gerblé	59	8	5.2	2.1	0.2/1.4/0.5	1.8	0.3	0.06			
<i>Oat and coconut</i>											
Amandín	49	7.3	4.5	2	0.8/-/-	0.5	0.6	0.12			

Nutrición  
 Hospitalaria

# **Nutrición Hospitalaria**