

# Nutrición Hospitalaria



**Relación entre índices  
antropométricos e indicadores de  
composición corporal por  
antropometría de brazo en  
pacientes pediátricos  
hospitalizados**  
**Relationship of anthropometric  
indexes and indicators of body  
composition by arm  
anthropometry on hospitalized  
pediatric patients**

10.20960/nh.2309

## **OR 2309 EPIDEMIOLOGÍA**

### **Relationship of anthropometric indexes and indicators of body composition by arm anthropometry on hospitalized pediatric patients**

*Relación entre índices antropométricos e indicadores de composición corporal por antropometría de brazo en pacientes pediátricos hospitalizados*

Nelly C. Muñoz-Esparza<sup>1</sup>, Edgar M. Vásquez-Garibay<sup>1,2</sup>, Larrosa-Haro Alfredo<sup>1</sup> and Enrique Romero-Velarde<sup>1,2</sup>

<sup>1</sup>Instituto de Nutrición Humana. Universidad de Guadalajara. Guadalajara, México. <sup>2</sup>Nuevo Hospital Civil de Guadalajara “Dr. Juan I. Menchaca”. Guadalajara, Jalisco. México

**Received:** 12/09/2018

**Accepted:** 26/11/2018

**Correspondence:** Edgar M. Vásquez-Garibay. Instituto de Nutrición Humana. Centro Universitario de Ciencias de la Salud. Universidad de Guadalajara. Nuevo Hospital Civil de Guadalajara. Salvador Quevedo y Zubieta, 750. Sector Libertad. 44240 Guadalajara, Jalisco. México  
e-mail: vasquez.garibay@gmail.com

#### **ABSTRACT**

**Introduction:** the purpose of this study was to evaluate the relationship of arm anthropometric indicators with direct indicators of nutritional status in hospitalized pediatric patients.

**Methods:** an analytical cross-sectional study with 760 patients hospitalized in the Pediatric Division of the Nuevo Hospital Civil de Guadalajara during 2014 was used. The anthropometric indices were

weight/length, weight/height, weight/age, length/age, height/age, head circumference/age and body mass index (BMI)/age. The arm indicators were mid-upper arm circumference (MUAC), total arm area (TAA), arm muscle area (AMA), arm fat area (AFA) and fat percentage (FP). The ANOVA, Kruskal-Wallis, Mann-Whitney U and Pearson's correlation tests and also odds ratios were used to identify the probability of nutritional status impairment.

**Results:** the prevalence of acute and chronic malnutrition was higher in infants (31% and 30%, respectively). With arm areas (TAA, AMA, AFA), the risk of deficit ( $\leq -2DE$ ) was higher in infants and early preschoolers ( $p < 0.001$ ). The correlation between the anthropometric indexes and the arm areas was direct and significant ( $p < 0.001$ ). The BMI variability was explained in 68% by the AMA, AFA, and FP ( $p < 0.001$ ); the variability of the height/age index was also explained in 34% by the AMA and AFA ( $p < 0.001$ ).

**Conclusion:** it is possible to diagnose both a chronic and acute deficit using the indirect indicators of the arm, while the body mass index only reflects an acute deficit. Therefore, arm areas would be more useful indicators in the assessment of nutritional status and the diagnosis of chronic-acute malnutrition in hospitalized pediatric patients.

**Key words:** Malnutrition. Body composition. Anthropometric indexes. Arm indicators. Pediatrics.

## RESUMEN

**Introducción:** el objetivo de este estudio fue evaluar la relación de los indicadores antropométricos de brazo con los indicadores directos del estado de nutrición en pacientes pediátricos hospitalizados.

**Métodos:** se utilizó un estudio transversal analítico con 760 pacientes ingresados en la División de Pediatría del Nuevo Hospital Civil de Guadalajara durante 2014. Los índices antropométricos fueron

peso/longitud, peso/altura, peso/edad, longitud/edad, altura/edad, circunferencia cefálica e IMC. Los indicadores del brazo fueron circunferencia media del brazo (CMB), área total del brazo (ATB), área muscular del brazo (AMB), área grasa del brazo (AGB) y porcentaje de grasa. Se utilizaron las pruebas de ANOVA, Kruskal-Wallis, U de Mann-Whitney, correlación de Pearson y razón de momios para identificar la probabilidad de deterioro del estado nutricional.

**Resultados:** la prevalencia de desnutrición aguda y crónica fue mayor en lactantes (31% y 30%, respectivamente). Con las áreas del brazo (ATB, AMB, AFA), el riesgo de déficit ( $\leq -2$  DE) fue mayor en lactantes y preescolares tempranos ( $p < 0,001$ ). La correlación entre los índices antropométricos y las áreas del brazo fue directa y significativa ( $p < 0,001$ ). La variabilidad del IMC fue explicada en un 68% por AMB, AGB y porcentaje de grasa ( $p < 0,001$ ); la variabilidad del índice de talla/edad también fue explicada en un 34% por AMB y AGB ( $p < 0,001$ ).

**Conclusión:** es posible diagnosticar el déficit crónico y agudo utilizando los indicadores indirectos del brazo, mientras que el IMC solo refleja un déficit agudo. Las áreas de brazo serían indicadores más útiles en el diagnóstico de desnutrición crónica-aguda en pacientes pediátricos hospitalizados.

**Palabras clave:** Desnutrición. Composición corporal. Índices antropométricos. Indicadores de brazo.

## INTRODUCTION

Protein energy malnutrition (PEM) continues to be a public health problem worldwide, mainly in children under five years of age (1,2). In hospitalized pediatric patients, the prevalence of malnutrition ranges between 6% and 50%; this wide range of discrepant reported frequencies seems a consequence of the degree of development of the country analyzed (1,3-6). Some authors have described that, according

to its severity, PEM can increase the morbidity and mortality of the hospitalized child (7). Several researchers have emphasized the importance of carrying out an adequate nutritional assessment of any child entering the hospital to identify any pediatric patients who are admitted with some degree of malnutrition (4,7-9) and those who are at risk of developing malnutrition during hospitalization (6,10). Several studies have shown that if an adequate assessment of a child's nutritional status is not carried out, no timely dietary management results and, as a consequence, the risk of complications and the length of the hospital stay increases (4,11,12).

Although there is some controversy about the usefulness of anthropometric indicators in the diagnosis of malnutrition (13), in general, there is a consensus that anthropometric indicators are adequate and accurate to assess nutritional status. In addition, they are noninvasive, low-cost, accessible, simple and practical (14,15). Particularly, the arm anthropometric indicators at the mid-upper-arm circumference (MUAC) and skin folds have been widely used in the hospital setting because they inform about body composition through the estimation of the total arm area (TAA), arm muscle area (AMA) and arm fat area (AFA) (15-19). Therefore, the purpose of this study was to estimate the indirect arm anthropometric indicators as an anthropometric expression of body composition and to explore its relationship with direct anthropometric indicators in hospitalized pediatric patients.

## **MATERIAL AND METHODS**

An analytical cross-sectional study was carried out with 750 patients hospitalized in the Pediatric Division of the Nuevo Hospital Civil de Guadalajara during 2014. All patients admitted in the previous day in weekdays to any of the services of the Pediatric Division were included. Those who remained less than 72 hours in the Emergency Room, who

were hospitalized in a clinical ward, and whose parents or legally responsible person signed the informed consent were included. We did not include patients in whom the anthropometric measurements could not be made and those who were re-hospitalized in a period of less than three months. Patients with incomplete records and/or data were excluded. For each participant (or relatives), a questionnaire was applied to the family member or legally responsible person that included general identification and sociodemographic data.

### **Anthropometric measurements**

Previous standardization of two observers was performed using the Habitch method (20); the measurements of weight, length, and cephalic circumference with the techniques previously described were made upon admission (3). With the measurements taken, the Z-scores of the weight/length, weight/height, weight/age, length/age, height/age, head circumference/age, and BMI/age were estimated using the WHO Anthro version 3.2.2 and WHO Anthro Plus version 1.0.4 programs.

The World Health Organization (WHO) normal limits ( $\pm 2SD$ ) were taken as a reference. The left mid-upper-arm circumference (MUAC) was obtained with a metallic metric tape (Rosscraft, USA). The triceps skinfold (TSF) was taken on the internal posterior surface of the previously marked middle part of the arm. The subscapular skin fold (SSF) was taken at the lower edge of the scapula. Both measurements were made with a Lange skinfold caliper (Michigan, USA). With MUAC and TSF arm areas were estimated with the Frisancho equations (21): total arm area (TAA):  $TAA (cm^2) = MUAC (cm^2)/(4 \times \pi)$ ; arm muscle area (AMA):  $AMA (cm^2) = [MUAC - (TSF (mm) \times \pi)]^2 / (4 \times \pi)$ ; arm fat area (AFA):  $AFA = TAA - AMA$ . Z-scores of these areas were estimated with the Sann reference values (22) for infants under 12 months and with the Frisancho reference (21) for children from one to 18 years. With the sum of the TSF and SSF, the percentage of body fat was calculated using the Slaughter

equation (23) in a differentiated way by gender. For purpose of anthropometrical and statistical analyses, the total population was stratified into age groups according to the WHO: infants, toddlers, preschoolers, schoolchildren and adolescents (24).

### **Statistical analysis**

One-way ANOVA with post-hoc tests (Bonferroni, Dunne's T3) were used. The odds ratio was estimated to identify the probability of alteration of the nutritional status. Correlation matrices were made between the anthropometric indexes, arm areas and percentage of body fat. Finally, multiple regression models were designed with the anthropometric indices as dependent variables and the arm areas and fat percentage as independent variables. The outlier values that were considered due to measurement error or capture were excluded from the analysis. The software SPSS version 20 was used.

### **Ethical considerations**

The protocol of investigation was evaluated and approved by the Bioethics and Research Committees of the Nuevo Hospital Civil de Guadalajara with the registration number in the Secretary of Health Jalisco: 1342/14.

### **RESULTS**

Of the 760 children included, 27% were infants (one to eleven months), 9% toddlers (12 to 23 months), 25% preschoolers (24 to 71 months), 21% schoolchildren (72 to 143 months) and 18% adolescents (144 to 216 months). The age average in months was  $4.6 \pm 3$  in infants,  $17.6 \pm 3$  in toddlers,  $45 \pm 14$  in preschoolers,  $104 \pm 21$  in schoolchildren and  $172 \pm 16$  in adolescents.

Table I shows the raw data and Z-scores of the anthropometric indicators; significant differences were observed in the anthropometric

indexes (Z) between age groups. Infants had a greater deficit compared to toddlers, preschoolers, schoolchildren and adolescents in almost all anthropometric indexes.

When analyzing the arm anthropometry, it was observed that the infants had a greater deficit in TAA (Z), AMA (Z) and AFA (Z) in comparison with the other age groups, and there was also a greater deficit in preschoolers *versus* schoolchildren and adolescents in TAA (Z) and AFA (Z) (Table II).

There was a higher prevalence of acute malnutrition (BMI < -2 SD) in infants (31%) and in adolescents (13%); likewise, a higher prevalence of chronic malnutrition (height/age < -2 SD) was observed in infants (30%) and in toddlers (18%). It should be noted that the presence of overweight/obesity increased gradually with age, until it was noticeable in schoolchildren (11%) and adolescents (8%). With respect to arm areas (TAA, AMA, AFA), a higher deficit ( $\leq -2$  SD) was observed in infants and toddlers. It should also be noted that there are differences between the indicators evaluated; for example, between the TAA and AMA against BMI, there were around 10% points of difference, with the TAA and AMA being more sensitive in the identification of acute malnutrition, especially in infants, toddlers and preschoolers (Table III). There was an increased risk of deficit in TAA in infants, toddlers and preschoolers. In relation to the AMA and AFA, the risk of deficit was significantly higher in infants, and the same happened with the BMI. The risk of deficit of height/age index was higher in infants, toddlers and late preschoolers (Table IV).

Table V shows the proportional direct correlations between the anthropometric indexes and the arm areas, which are directly proportional. When performing the linear regression, it was observed that the variability of the BMI is explained in 50% by the TAA, in 47% by the AMA, in 40% by the AFA, and in 46% by the percentage of fat. The variability of the weight/age index is explained in 53% by the TAA, in



51% by the AMA, and in 40% by the percentage of fat. The variability of the weight/height index is explained in 38% by the TAA, in 32% by the AMA, and in 42% by the percentage of fat. It should be noted that between the height/age index and the cephalic circumference, a positive correlation is maintained, where the height/age index predicts 46% of its variability.

Table VI shows the multiple linear regression models; it is observed that in children under 36 months of age, the variability of the BMI is explained in 67% by the AMA, AFA and percentage of fat; 31% of the variability of the cephalic circumference is explained by the AMA, AFA and percentage of fat. Likewise, the variability of the height/age index is explained in 35% by the AMA, AFA and percentage of fat. In patients older than 36 months, BMI variability is explained in 73% by the AFA and AMA; the variability of the height/age index is explained in 27% by the TAA, AMA and percentage of fat.

## **DISCUSSION**

In the studied pediatric sample in hospitalized patients, it was observed that the prevalence of acute malnutrition (deficit in BMI) and chronic malnutrition (deficit in height/age) was higher in infants than in preschoolers, schoolchildren and adolescents. These findings do not differ from those observed by other researchers (4-6). The probability of deficit in arm areas was significantly higher in children under 24 months (infants and toddlers) than in the other age groups.

The frequency of deficit observed with these indirect anthropometric indicators of the arm coincided with the frequency of malnutrition described above with the anthropometric indexes of BMI and height/age. It is known that when there is an impairment in nutritional status, the reserves of fat and muscle (reflected in the areas of the arm) are significantly affected, particularly at early ages when growth and development are accelerated and there is greater metabolic activity.

Therefore, any moderate or severe nutritional insult has a significant effect on nutritional status and body composition (3,15). These findings corroborate the hypothesis that arm areas are a useful tool in the diagnosis of malnutrition, especially acute malnutrition, regardless of the age group; it is also important to note that with the TAA and AMA, the identification of acute malnutrition increased by 10% points.

Another interesting finding refers to the positive correlations that occurred between the anthropometric indexes and body composition indicators. The correlation between BMI and TAA is noteworthy, since both indicators include fat and muscle mass; also, the correlation between BMI with the AMA and the AFA was observed previously (25,26). Hurtado-López et al. (17) mention that the anthropometry of the arm has a positive correlation with the indicators of body composition and, in turn, with linear growth. Their findings coincide with those observed in this study. As it has been observed, when the pediatric patient is undergoing a nutritional insult, the fat and muscle reserves are affected in the first instance; if the nutritional insult continues, linear growth is affected.

In the hospital setting, the prevalence of malnutrition is high, mainly in intermediate or intensive therapies (3,5,7). It should be noted that in these units of care for critically ill pediatric patients, a complete assessment of the nutritional status of the patient is not usually undertaken, due to the severity of the condition that prevents the patient from moving and/or the lack of adequate equipment to perform the proper anthropometric evaluation. Under these conditions, the evaluation of arm anthropometry is a good option, either as part of a comprehensive evaluation or as a specific alternative way of assessing nutritional status. This suggestion is based on the analysis of the multiple linear regression models, where it was observed that the anthropometric indexes are largely explained by the muscle and fat areas of the arm (16,25). The arm anthropometry has been commonly

used in the evaluation of patients with chronic kidney disease, chronic liver disease, and cystic fibrosis because the clinical conditions presented by these patients (visceromegaly, generalized edema, etc.) make it difficult to interpret weight/age, weight/height and BMI indexes (16,17,19).

There are several methods to evaluate body composition, such as dual energy X-ray absorptiometry (DXA) and bioelectrical impedance (IBE), among others (1). Among its advantages, the accuracy of the evaluation stands out; however, they are costly methods and are not always accessible in the hospital units that care for children, especially in less industrialized countries. In addition, the usefulness of these options in the hospitalized patient can be limited by the patient's clinical conditions, as is the case with IBE, which is affected by the patient's hydration conditions. Therefore, anthropometry of the arm would be an optimal, adequate, accessible and simple method for evaluating the body composition of the hospitalized patient, an opinion shared by other researchers (17,19,27-29).

One strength of the study is that the size of the sample was large and that different age groups with different pathologies were included. In addition, the length of the study period was a full year. One possible limitation was the outliers that had to be discarded in the statistical analysis.

In conclusion, the evaluation of arm anthropometry is a good option, either as part of a comprehensive evaluation or as a specific alternative way of assessing nutritional status. The measurement of the arm areas is a useful tool in the diagnosis of chronic-acute malnutrition, while the BMI only reflects an acute deficit in hospitalized pediatric patients.

## **REFERENCES**

1. Pileggi VN, Monteiro JP, Margutti AVB, Camelo-Jr JS. Prevalence of child malnutrition at a university hospital using the World Health

- Organization criteria and bioelectrical impedance data. *Braz J Med Biol Res* 2016;49:1-8. DOI: 10.1590/1414-431X20155012.
2. World Health Organization (WHO). Children: reducing mortality; WHO: Geneva. Updated October 2017. Accessed on November 2017. Available from: <http://www.who.int/mediacentre/factsheets/fs178/en/>
  3. Muñoz-Esparza NC, Vásquez-Garibay EM, Romero-Velarde E, Troyo-Sanromán R. Risk of malnutrition of hospitalized children in a university public hospital. *Nutr Hosp* 2017;34:41-50. DOI: 10.20960/nh.974.
  4. Kapci N, Akcam M, Koca T, Dereci S, Kapci M. The nutritional status of hospitalized children: has this subject been overlooked? *Turk J Gastroenterol* 2015;26:351-5. DOI: 10.5152/tjg.2015.0013
  5. Toussaint-Martínez de Castro G, Kaufer-Horwitz M, Carrillo-López HA, Klünder-Klünder M, Jarillo-Quijada A, García-Hernández HR. Estado nutricional de niños en condiciones críticas de ingreso a las unidades de terapia intensiva pediátrica. *Bol Med Hosp Infant Mex* 2013;70:216-21.
  6. Baxter JA, Al-Madhaki FI, Zlotkin SZ. Prevalence of malnutrition at the time of admission among patients admitted to a Canadian tertiary-care pediatric hospital. *Paediatr Child Health* 2014;19:413-7. PMID: PMC4220524
  7. Velandia S, Hodgson MI, Le Roy C. Evaluación nutricional en niños hospitalizados en un Servicio de Pediatría. *Rev Chil Pediatr* 2016;87:359-65. DOI: 10.1016/j.rchipe.2016.05.001
  8. Fonseca-Teixeira A, Araújo-Lourenco-Viana KD. Nutritional screening in hospitalized pediatric patients: a systematic review. *J Pediatr (Río J)* 2016;92:343-52. DOI: 10.1016/j.jpdp.2016.02.009
  9. Shaughnessy EE, Kirkland LL. Malnutrition in hospitalized children: a responsibility and opportunity for pediatric hospitalists. *Hosp Pediatr* 2016;6:37-41. DOI: 10.1542/hpeds.2015-0144
  10. Lama-More RA, Moráis-López A, Herrero-Álvarez M, Caraballo-Chicano S, Galera-Martínez R, López-Ruzafa E, et al., grupo GETNI. Validación de una herramienta de cribado nutricional para pacientes pediátricos hospitalizados. *Nutr Hosp* 2012;27:1429-36. DOI: 10.3305/nh.2012.27.5.5467

11. Moreno-Villares JM, Varea-Calderón V, Bousoño-García C, Lama-More R, Redecillas-Ferreiro S, Peña-Quintana L. Evaluación del estado nutricional de niños ingresados en el hospital en España; estudio DHOSPE (Desnutrición Hospitalaria en el Paciente Pediátrico en España). *Nutr Hosp* 2013;28:709-18. DOI: 10.3305/nh.2013.28.3.6356
12. Santafé-Sánchez LR, Sánchez-Rodríguez DA, Villegas-Galarza AL, González-Correa CH. Nutritional status among hospitalized children with mixed diagnoses at a referral teaching hospital in Manizales, Colombia. *Nutr Hosp* 2012;27:1451-9. DOI: 10.3305/nh.2012.27.5.5875
13. Molina Soria JB, Lobo Támer G, Pérez de la Cruz AJ, Ruiz-López MD. Prevalencia de desnutrición al ingreso en un hospital general básico. *Nutr Hosp* 2017;34:1390-8. DOI: 10.20960/nh.1133
14. Rodríguez PN, Bermúdez EF, Rodríguez GS, Spina MA, Zeni SN, Friedman SM, et al. Composición corporal en niños preescolares: comparación entre métodos antropométricos simples, bioimpedancia y absorciometría de doble haz de rayos X. *Arch Argent Pediatr* 2008;106:102-9. DOI: 10.1590/S0325-00752008000200003
15. Velásquez M, Salazar G, Vio del RF, Díaz N, Anziani A. Validación de ecuaciones antropométricas para evaluar composición corporal en niños preescolares chilenos. *Rev Méd Chile* 2008;136:433-41. DOI: S0034-98872008000400003.
16. Hurtado-López EF, Larrosa-Haro A, Vásquez-Garibay EM, Macías-Rosales R, Troyo-Sanromán R, Bojórquez-Ramos MC. Liver function test results predict nutritional status evaluated by arm anthropometric indicators. *Pediatr Gastroenterol Nutr* 2007;45:451-7. DOI: 10.1097/MPG.0b013e3180f60b9e
17. Hurtado-López EF, Vásquez-Garibay EM, Trujillo X, Larrosa-Haro A. Body composition predicts growth in infants and toddlers with chronic liver disease. *J Pediatr Gastroenterol Nutr* 2017;65:e117-9. DOI: [10.1097/MPG.0000000000001746](https://doi.org/10.1097/MPG.0000000000001746)
18. Berkley J, Mwangi I, Griffiths K, Ahmed I, Mithwani S, English M, et al. Assessment of severe malnutrition among hospitalized children in

- rural Kenya - Comparison of weight for height and mid upper arm circumference. JAMA 2005;294:591-7. DOI: 10.1001/jama.294.5.591
19. Chomtho S, Fewtrell MS, Jaffe A, Williams JE, Wells JCK. Evaluation of arm anthropometry for assessing pediatric body composition: evidence from healthy and sick children. *Pediatr Res* 2006;59:860-5. DOI: [10.1203/01.pdr.0000219395.83159.91](https://doi.org/10.1203/01.pdr.0000219395.83159.91)
20. Habitch JP. Estandarización de métodos epidemiológicos cuantitativos sobre el terreno. *Bol Oficina Sanit Panam* 1974;76:375-84.
21. Frisancho AR. Anthropometric standards for the assessment of growth and nutritional status. Michigan: The University of Michigan Press; 1990. pp. 9-30.
22. Sann L, Durand M, Picard J. Arm fat and muscle areas in infancy. *Arch Dis Child* 1988;63:256-60.
23. Slaughter MH, Lohman TG, Boileau RE, Horswill CA, Stillman RJ, Van Loan MD, et al. Skinfold equation for estimation of body fatness in children and youth. *Hum Biol* 1988;60:709-23.
24. World Health Organization (WHO). Ciclo de vida. Geneva; WHO. Updated November 2018. Accessed on November 2018. Available from: [http://www.who.int/elena/life\\_course/es/](http://www.who.int/elena/life_course/es/)
25. Grijalva-Eternod CS, Wells JCK, Girma T, Kaestel P, Admassu B, Friis H, et al. Midupper arm circumference and weight-for-length z-scores have different associations with body composition: evidence from a cohort of Ethiopian infants. *Am J Clin Nutr* 2015;102:593-9. DOI: 10.3945/ajcn.114.106419
26. Gharib NM, Rasheed P. Anthropometry and body composition of school children in Bahrain. *Ann Saudi Med* 2009;29:258-69. DOI: [10.4103/0256-4947.55309](https://doi.org/10.4103/0256-4947.55309)
27. Vásquez-Garibay EM, Larios-del-Toro YE, Larrosa-Haro A, Troyo-Sanromán R. Anthropometric indicators of nutritional status and growth in very low birth-weight premature infants hospitalized in a neonatal Intensive Care Unit. *Nutr Hosp* 2014;30:410-6. DOI: 10.3305/nh.2014.30.2.7373
28. Ortiz-Hernández L, Vega-López V, Ramos-Ibáñez N, Cazares-Lara LJ, Medina-Gómez J, Pérez-Salgado D. Equations based on anthropometry to

predict body fat measured by absorptiometry in schoolchildren and adolescents. J Pediatr 2017;93:365-73. DOI: DOI: 10.1016/j.jpeds.2016.08.008

29. Macías-Rosales R, Larrosa-Haro A, Ortiz-Gabriel G, Trujillo-Hernández. Effectiveness of enteral versus oral nutrition with a medium-chain triglyceride formula to prevent malnutrition and growth impairment in infants with biliary atresia. J Pediatr Gastroenterol Nutr 2016;62:101-9. ISSN: 0277-2116. DOI: 10.1097/MPG.0000000000000909

Nutrición  
Hospitalaria

**Table I. Raw data and Z-scores according to the age groups (n = 750). Comparison of values between groups**

Indicator		Infant	Toddler	Preschool	Schoolchildr	Adolescen
		s	s	ers	en	ts
Weight (kg)	n	198	70	189	160	133
	Mean	5.78	10.10	15.13	28.99	50.42
	SD	2.05	1.69	3.19	10.69	12.88
Lenght/height (cm)	N	198	70	189	160	133
	Mean	60.99	79.49	99.20	130.1	158.29
	SD	7.37	5.30	9.14	11.76	9.19
BMI (kg/m <sup>2</sup> )	n	198	70	189	160	133
	Mean	14.95	15.91	15.34	16.63	19.96
	SD	2.38	1.62	1.48	3.54	4.28
Cephalic C (cm)	n	192	68	52	-	-
	Mean	39.8	46.25	48.21	-	-
	SD	3.6	1.81	1.63	-	-
Weight/age (Z)*	n	198	70	189	-	-
	Mean	-1.68	-0.53	-0.43	-	-
	SD	1.77	1.35	1.11	-	-
Weight/height (Z)*	n	197	70	146	-	-
	Mean	-0.84	-0.27	-0.20	-	-
	SD	1.46	1.34	1.10	-	-
Cephalic C (Z)*	n	192	68	52	-	-
	Mean	-1.36	-0.38	-0.15	-	-
	SD	1.60	1.21	1.22	-	-
BMI/age (Z)*	n	198	70	189	160	133
	Mean	-1.26	-0.15	-0.15	-0.12	-0.20
	SD	1.65	1.31	1.11	1.53	1.54
Height/age (Z)*	n	198	70	189	160	133
	Mean	-1.45	-0.73	-0.54	-0.12	-0.40
	DE	1.62	1.33	1.17	1.01	1.08

BMI: body mass index; Cephalic C: cephalic circumference. \*ANOVA differences between groups,  $p < 0.001$ ; *post-hoc* tests; T3 by Dunnett and Bonferroni. Weight/age: infants vs toddlers  $p < 0.001$ ; infants vs preschoolers  $p < 0.001$ . Height/age: infants vs toddlers  $p < 0.001$ ; infants vs preschoolers  $p < 0.001$ ; infants vs schoolchildren  $p < 0.001$ ; infants vs adolescents  $p < 0.001$ ; toddlers vs schoolchildren  $p = 0.001$ ; preschoolers vs schoolchildren  $p = 0.001$ . Weight/height: infants vs toddlers  $p = 0.001$ ; infants vs preschoolers  $p = 0.001$ . BMI/age: infants vs



toddlers  $p < 0.001$ ; infants vs preschoolers  $p < 0.001$ ; infants vs schoolchildren  $p < 0.001$ ; infants vs adolescents  $p < 0.001$ . Cephalic circumference: infants vs toddlers  $p < 0.001$ ; infants vs preschoolers  $p < 0.001$ .



**Table II. Skinfolds and arm areas according to the age groups (n = 749). Comparison of arm areas between groups (Z-score)<sup>†</sup>**

Age group		MUAC (cm)	TSF (mm)	SSF (mm)	TAA (cm <sup>2</sup> )	TAA (Z)	AMA (cm <sup>2</sup> )	AMA (Z)	AFA (cm <sup>2</sup> )	AFA (Z)	Body fat (%)*
Infants	n	198	195	195	198	198	195	195	195	195	197
	Mea	11.47	7.27	5.8	10.77	-1.69	6.86	-1.68	3.90	-0.79	12.71
	SD	1.96	2.65	2.1	3.63	1.54	2.17	1.24	1.83	1.93	4.62
Toddlers	n	69	69	69	69	69	69	69	69	69	69
	Mea	13.62	7.26	5.6	14.92	-0.94	10.32	-1.05	4.60	-1.24	12.60
	SD	1.37	1.76	1.5	2.85	1.09	2.01	0.88	1.30	0.58	3.23
Preschoolers	n	189	189	187	189	189	189	189	189	189	185
	Mea	15.10	8.31	5.8	18.36	-0.41	12.53	-0.95	5.84	-0.74	13.56
	SD	1.64	2.88	2.1	3.99	1.18	2.33	0.79	2.41	0.94	4.31
Schoolchildren	n	160	160	160	160	160	160	160	160	160	159
	Mea	19.00	12.55	8.7	29.81	0.15	18.35	-0.86	11.46	0.05	18.65
	SD	3.73	6.61	2.1	12.11	1.10	5.10	0.79	7.84	1.06	8.69
Adolescents	n	133	133	131	133	133	133	133	133	133	131
	Mea	23.66	16.00	12.4	45.99	0.05	28.32	-0.97	17.66	0.06	23.87
	SD	4.27	7.68	6.0	16.27	1.09	8.47	1.01	10.49	0.95	8.92

MUAC: mid-upper-arm circumference; TSF: tricipital skin fold; SSF: subscapular skin fold; TAA: total arm area; AMA: arm muscle area; AFA: arm fat area. \*Percentage of fat (Slaughter, 1988). <sup>†</sup>ANOVA, differences between

groups,  $p < 0.001$ ; *post-hoc* tests; Dunnett's T3. TAA (Z): infants vs toddlers, preschoolers, schoolchildren and adolescents,  $p < 0.001$ ; preschoolers vs schoolchildren and adolescents,  $p < 0.005$ . AMA (Z): infants vs toddlers, preschoolers, schoolchildren and adolescents,  $p < 0.001$ . AFA (Z): infants vs schoolchildren and adolescents,  $p < 0.001$ ; toddlers vs preschoolers, schoolchildren and adolescents,  $p < 0.001$ ; preschoolers vs schoolchildren and adolescents,  $p < 0.001$ .

Nutrición  
Hospitalaria

**Table III. Distribution of the frequency (%) of the anthropometric indexes and arm areas by age group in Z-score (n = 760)**

Age group		TAA	AMA	AFA	BMI/age	Height/age
		n (%)	n (%)	n (%)	n (%)	n (%)
Infants	n	202	199	199	202	202
	< -3 a ≤ -2	95 (47)	86 (43.2)	62 (31.2)	63 (31.2)	60 (29.7)
	> -2 a ≤ 2	102 (50.5)	111 (55.8)	122 (61.3)	138 (68.3)	141 (69.8)
	> 2	5 (2.5)	2 (1)	15 (7.5)	1 (0.5)	1 (0.5)
Toddlers	n	70	70	70	71	71
	< -3 a ≤ -2	13 (18.6)	7 (10)	6 (8.6)	7 (9.9)	13 (18.3)
	> -2 a ≤ 2	57 (81.4)	63 (90)	64 (91.4)	62 (87.3)	58 (81.7)
	> 2	-	-	-	2 (2.8)	-
Preschoolers	n	191	191	191	191	191
	< -3 a ≤ -2	19 (9.9)	13 (6.8)	9 (4.7)	9 (4.7)	20 (10.5)
	> -2 a ≤ 2	166 (86.9)	177 (92.7)	178 (93.2)	176 (92.2)	169 (88.5)
	> 2	6 (3.1)	1 (0.5)	4 (2.1)	6 (3.1)	2 (1)
Schoolchildre	n	161	161	161	161	161
	< -3 a ≤ -2	2 (1.2)	9 (5.6)	-	15 (9.3)	7 (4.3)
	> -2 a ≤ 2	147 (91.3)	151 (93.8)	154 (95.7)	129 (80.1)	150 (93.2)
	> 2	12 (7.5)	1 (0.6)	7 (4.3)	17 (10.6)	4 (2.5)
Adolescents	n	135	135	135	135	135
	< -3 a ≤ -2	2 (1.5)	13 (9.6)	-	17 (12.6)	9 (6.7)
	> -2 a ≤ 2	124 (91.9)	117 (86.7)	130 (96.3)	107 (79.3)	124 (91.9)
	> 2	9 (6.7)	5 (3.7)	5 (3.7)	11 (8.1)	2 (1.5)

TAA: total arm area; AMA: arm muscle area; AFA: arm fat area; BMI: body mass index.

**Table IV. Probability of deficit (OR) in indicators of body composition by age group (< -2 Z) (n = 760)**

<i>Indicators</i>	<i>OR</i>	<i>95% CI</i>	<i>p</i>
<b>Total arm area</b>			
Infants vs toddlers	4	2.01-7.55	< 0.001
Infants vs preschoolers	8	4.64-13.9	< 0.001
Infants vs schoolchildren	71	17.0-292	< 0.001
Infants vs adolescents	59	14.2-245	< 0.001
Toddlers vs schoolchildren	18	3.97-82.8	< 0.001
Toddlers vs adolescents	15	3.32-69.4	< 0.001
Preschoolers vs schoolchildren	9	2.01-38.3	0.001
Preschoolers vs adolescents	7	1.68-32.1	0.004
<b>Arm muscular area</b>			
Infants vs preschoolers	7	2.99-15.70	< 0.001
Infants vs preschoolers	10	5.56-19.55	< 0.001
Infants vs schoolchildren	13	6.20-26.63	< 0.001
Infants vs adolescents	7	3.78-13.50	< 0.001
<b>Arm fat area</b>			
Infants vs toddlers	5	1.98-11.74	< 0.001
Infants vs preschoolers	9	4.46-19.34	< 0.001
<b>BMI</b>			
Infants vs toddlers	4	1.80-9.55	< 0.001
Infants vs preschoolers	9	4.41-19.10	< 0.001
Infants vs schoolchildren	4	2.40-8-11	< 0.001
Infants vs adolescents	3	1.75-5.67	< 0.001
Adolescents vs preschoolers	3	1.26-6.75	0.017
<b>Height/age</b>			
Infants vs preschoolers	4	2.05-6.21	< 0.001
Infants vs schoolchildren	9	4.11-21.01	< 0.001
Infants vs adolescents	6	2.82-12.41	< 0.001
Toddlers vs schoolchildren	5	1.88-12.97	0.001
Toddlers vs adolescents	3	1.27-7.76	0.020
Preschoolers vs school children	3	1.06-6.25	0.051

BMI: body mass index.

**Table V. Correlation and determination coefficients of Z-scores of indirect arm anthropometric indicators (independent variable) with Z-scores nutritional status and growth indicators (dependent variable) obtained in a sample of hospitalized pediatric patients**

Dependent variable	Independent variable	n	r	R2	$\beta$	p
BMI	Total arm area	748	0.669	0.489	0.477	< 0.001
	Arm fat area	753	0.657	0.432	0.453	< 0.001
	Arm muscle area	753	0.567	0.432	0.453	< 0.001
Weight for height	Total arm area	408	0.594	0.353	0.528	< 0.001
	Arm fat area	409	0.521	0.271	0.469	< 0.001
	Arm muscle area	409	0.522	0.305	0.670	< 0.001
Weight for age	Total arm area	568	0.698	0.473	0.721	< 0.001
	Arm fat area	567	0.566	0.320	0.625	< 0.001
	Arm muscle area	569	0.683	0.466	1.042	< 0.001
Height for age	Total arm area	733	0.507	0.257	0.441	< 0.001
	Arm fat area	734	0.410	0.168	0.377	< 0.001
	Arm muscle area	735	0.518	0.269	0.639	< 0.001
Cephalic circumference for age	Total arm area	308	0.469	0.220	0.467	< 0.001
	Arm fat area	312	0.348	0.121	0.345	< 0.001
	Arm muscle area	312	0.515	0.265	0.690	< 0.001

BMI: body mass index.

**Table VI. Multiple Linear Regression Models\*. Relationship between indirect anthropometric indicators (Z-score) and percentage of body fat<sup>†</sup> with direct indicators (Z score) (n = 749)**

Dependent variable	Independent variable	n	Mean	SD	r	R <sup>2</sup>	p
<b>&lt; 36 months</b>							
BMI	AMA	330	-1.40	1.18	0.82	0.67	< 0.001
	AFA		-0.98	1.60			
	% Body fat		12.6	4.27			
Cephalic C	AMA	311	-1.41	1.20	0.55	0.30	0.001
	AFA		-0.99	1.63			
	% Body fat		12.6	2.28			
Height/age	AMA	330	-1.40	1.18	0.59	0.34	0.010
	AFA		-0.98	1.60			
	% Body fat		12.6	4.27			
<b>≥ 36 months</b>							
BMI	AFA	425	-0.09	1.24	0.85	0.72	< 0.001
	AMA		-0.91	0.97			
Height/age	TAA	425	0.02	1.35	0.52	0.27	< 0.001
	AMA		-0.90	0.96			
	% body fat		19.2	9.1			
<b>Total population</b>							
BMI	AMA	748	-1.13	1.09	0.82	0.67	< 0.001
	AFA		-0.47	1.48			
Height/age	% body fat	748	16.3	8.04	0.58	0.33	< 0.001
	AMA		-1.13	1.09			
	AFA		-0.47	1.48			

TAA: total arm area; AMA: arm muscle area; AFA: arm fat area; Cephalic C: cephalic circumference; BMI: body mass index. \*Stepwise method. <sup>†</sup>Estimated fat percentage with Slaughter's equation (1988).