

**El polimorfismo del gen INSIG2
se asocia con una mayor presión
sanguínea y niveles de
triglicéridos en sujetos obesos
brasileños
INSIG2 gene polymorphism is
associated with higher blood
pressure and triglyceride levels
in Brazilian obese subjects**

OR 2359

INSIG2 gene polymorphism is associated with higher blood pressure and triglyceride levels in Brazilian obese subjects

El polimorfismo del gen INSIG2 se asocia con una mayor presión sanguínea y niveles de triglicéridos en sujetos obesos brasileños

Carolina F. Nicoletti, Raquel G. Azevedo, Marcela A.S. Pinhel, Heitor B.P. Delfino and Carla B. Nonino

Department of Internal Medicine. Ribeirão Preto Medical School. University of São Paulo. Ribeirão Preto, São Paulo

Received: 22/10/2018

Accepted: 15/11/2018

Correspondence: Carla Barbosa Nonino. Laboratory of Nutrigenomic Studies. Ribeirão Preto Medical School (FMRP/USP). Av. Bandeirantes, 3900. 14049-900 Monte Alegre, Ribeirão Preto. São Paulo, Brazil
e-mail: carla@fmrp.usp.br

ABSTRACT

Objective: this study aimed to evaluate the association between polymorphisms of INSIG, PCSK9 and FTO genes with anthropometric, biochemical characteristics and presence of metabolic syndrome in patients with severe obesity.

Material and methods: the present study enrolled 150 patients with grade II or III obesity, who were submitted to nutritional assessment, blood pressure measurement and peripheral blood collection. INSIG2 (rs75666605), PCSK9 (rs505151), and FTO (rs9939609) polymorphisms were genotyped using TaqMan Pre-Designed SNP Genotyping Assays probes in real time polymerase chain reaction

(PCR). The experimental data are processed in SPSS Statistics 22.0 ($p < 0.05$).

Results: in this study, 72.2% of obese subjects had metabolic syndrome (MS). There was a higher prevalence of AA (86.9%), CG (51.1%) and AT (46.2%) genotypes for the PCSK9, INSIG2 and FTO polymorphisms, respectively. There was no association of these polymorphisms with the prevalence of MS ($p > 0.05$). On the other hand, individuals with at least one variant allele (G) for the INSIG2 gene had higher triglycerides levels, systolic and diastolic blood pressure ($p < 0.05$).

Conclusions: the polymorphism rs7566605 of the INSIG2 gene is associated with higher triglycerides levels and blood pressure values, which are also considered as risk factors for the development of MS.

Key words: Obesity. Metabolic syndrome. Polymorphism. INSIG2. PCSK9. FTO.

RESUMEN

Objetivo: este estudio tuvo como objetivo evaluar la asociación entre polimorfismos de los genes INSIG, PCSK9 y FTO con las características antropométricas, bioquímicas y la presencia de síndrome metabólico (SM) en pacientes con obesidad grave.

Material y métodos: el presente estudio incluyó 150 pacientes con obesidad de grado II o III, que fueron sometidos a evaluación nutricional, medición de la presión arterial y extracción de sangre periférica. Los polimorfismos INSIG2 (rs75666605), PCSK9 (rs505151) y FTO (rs9939609) fueron genotipados utilizando sondas TaqMan Pre-Designed SNP Genotyping Assays en la reacción en cadena de la polimerasa en tiempo real (PCR). Los datos experimentales se procesan en SPSS Statistics 22.0 ($p < 0,05$).

Resultados: en este estudio, el 72,2% de los sujetos obesos tenían síndrome metabólico (EM). Hubo una mayor prevalencia de genotipos AA (86,9%), CG (51,1%) y AT (46,2%) para los polimorfismos PCSK9,

INSIG2 y FTO, respectivamente. No hubo asociación de estos polimorfismos con la prevalencia de SM ($p > 0,05$). Por otro lado, los individuos con al menos una variante de alelo (G) para el gen INSIG2 tenían niveles más altos de triglicéridos, presión arterial sistólica y diastólica ($p < 0,05$).

Conclusiones: el polimorfismo rs7566605 del gen INSIG2 se asocia con niveles más altos de triglicéridos y valores de presión arterial, que también se consideran factores de riesgo para el desarrollo del síndrome metabólico.

Palabras clave: Obesidad. Síndrome metabólico. Polimorfismo. INSIG2. PCSK9. FTO.

INTRODUCTION

Metabolic syndrome (MS) is a term used to characterize cardiovascular risk factors and is characterized by the association of various abnormalities, such as central obesity, dyslipidemia, type 2 diabetes mellitus and systemic arterial hypertension (SAH). There are different definitions for diagnostic criteria, all using combinations of risk factors (18,25). MS frequency is significantly associated with favorable socioeconomic conditions, sedentary lifestyle, hypercaloric diets and high body mass index (BMI) (20). The global prevalence of MS varies depending on the region, urban or rural environment and characteristics of the population studied, like sex, age, race and ethnicity (16).

Considering that obesity is one of the risk factors for MS, the literature has pointed out the important role of genetic susceptibility for this disease development (23), including gene interactions and gene-environment interactions (19). Recent studies show the existence of several single nucleotide polymorphisms (SNPs) located in different genes involved in lipid metabolism and insulin resistance, which play a role in obesity development (22).

In this context, the proprotein convertase subtilisin/kexin 9 (PCSK9) gene encodes the glycoprotein PCSK9, which plays an important role in the cholesterol metabolism. Its expression modulates the number of low-density lipoprotein (LDLR) receptors (28) and affects the circulating levels of low-density lipoprotein cholesterol (LDL-cholesterol), the main carrier of cholesterol in humans (11). The polymorphism E670G (rs505151) of the PCSK9 gene promotes increase of PCSK9 activity (gain of function), leading to reduction of LDLR and consequently, raise in plasma LDL-cholesterol, contributing to hypercholesterolemia (11,28).

Another gene that regulates cholesterol synthesis is the insulin-induced gene 2 (INSIG2), which encodes the INSIG2 membrane protein (8). INSIG2 inhibits sterol synthesis in the liver when the cells have an adequate supply of these (29). Deregulation of this process may affect lipid metabolism and increase insulin resistance (3), which are closely related to MS (4). Some studies have shown an association between the polymorphism rs7566605 of INSIG2 gene and obesity (13,29).

Furthermore, the fat mass and obesity-associated gene (FTO) encode an important protein whose exact physiological function remains unknown. Studies indicate that this gene plays a role in the nervous and cardiovascular systems and has a strong association with BMI, obesity and type 2 diabetes development (30,31). The FTO gene has several known SNPs, many of them related to BMI, with emphasis in rs9939609. In this case, the A allele is directly related to a greater accumulation of body fat, especially in homozygous form (AA).

In line with this, the objective of the present study was to evaluate the association between polymorphisms of INSIG2, PCSK9 and FTO genes with anthropometric, biochemical characteristics and presence of metabolic syndrome in patients with severe obesity.

MATERIALS AND METHODS

Subjects

This is a cross-sectional study, which enrolled individuals of an ethnic mixed population (27), aged between 18 and 60 years, with grade II or III obesity. The sampling procedure is fully described elsewhere and anthropometric and biochemical data of the studied population were already published by our group (21). Pregnant and lactating individuals, patients on hypocaloric diets and who underwent bariatric surgery were excluded. The study was conducted after approval by the Ethics Committee of the institution and according to the Declaration of Helsinki. All participants provided a written informed consent.

Questions about personal history of chronic diseases such as diabetes mellitus, systemic arterial hypertension, cardiovascular diseases and obesity were collected from medical records.

Anthropometric and blood pressure (BP) measurements

For anthropometric evaluation, the following indicators were used: weight (kg), height (m), BMI (kg/m^2) and abdominal circumference (AC). According to Nicoletti et al. (2016), patients were weighed in a Filizola® scale-type digital scale, with a capacity of 300 kg and an accuracy of 0.2 kg. For height measurement, a vertical rod with a graduation of 0.5 cm was used. BMI was obtained using the formula: $\text{BMI} = P/A^2$, where P is the weight in kilograms and A is the height in meters. AC was measured by passing an inextensible metric tape with graduation of 0.1 mm on the largest circumference.

The systolic and diastolic blood pressure (BP) was measured using the indirect method with an auscultatory technique using a mercury column sphygmomanometer, with cuffs appropriate for obese individuals. The procedure was performed with the patient after five minutes of rest in a quiet environment, without having practiced physical exercises for at least 60 minutes and without having smoked in the 30 minutes before the procedure.

Biochemical evaluation

Plasma total cholesterol (TC), low-density lipoprotein (LDL-cholesterol) and high-density lipoprotein (HDL-cholesterol), triglycerides (TG) and glycemia were analyzed in blood samples after 12 hours of fasting. As a protocol of the service, the dosages of TC, HDL-cholesterol and TG were performed by automated colorimetric method. LDL-cholesterol values were calculated by the Friedwald formula: $\text{LDL-cholesterol} = \text{TC} - (\text{HDL-cholesterol} + \text{TG}/5)$, with TG values lower than 400 mg/dl (9).

Metabolic syndrome identification

MS was defined according to the criteria of the I Brazilian Guidelines for the Diagnosis and Treatment of Metabolic Syndrome, which uses diagnostic standards similar to the National Cholesterol Education Program-Adult Treatment Panel III (NCEP ATP III). In this case, MS was defined by a group of cardiovascular risk factors, characterized by at least three of the following components: a) abdominal circumference > 88 cm; b) fasting glycemia > 100 mg/dl; c) triglycerides > 150 mg/dl; d) HDL cholesterol < 50 mg/dl; and e) increased blood pressure: systolic blood pressure (SBP) \geq 130 mmHg or diastolic blood pressure (DBP) \geq 85 mmHg.

DNA extraction and analysis of genetic polymorphisms

DNA was extracted from peripheral blood samples using a GE Healthcare kit according to manufacturer's protocol. The genotypic determination was performed by the allele discrimination method in real time PCR (polymerase chain reaction), which allows the analysis of the variants in a specific segment of DNA. The single nucleotide polymorphisms rs505151, rs75666605 and rs9939609 were genotyped using TaqMan™ Pre-Designed SNP Genotyping Assays probes (Applied Biosystems, Foster City, CA), as specified by the manufacturer.

Statistical analysis

Data were presented in mean and standard deviation. The Kolmogorov-Smirnov test was used to verify the data normality. The Hardy-Weinberg equilibrium calculations for the polymorphisms were performed by applying the Chi-square test. The t-test for independent samples was used to compare the phenotypic variables among the genotypes of each polymorphism. To analyze the dominant model (DM), individuals with at least one variant allele were grouped and compared with those of the reference genotype. For the recessive model (RM) analysis, individuals with at least one wild-type allele were grouped and compared with those with both variant alleles. The associations of continuous variables with genotypes were analyzed with general linear models controlling for sex and age. Statistical significance was set at 5%, all analyzes being performed in Statistical Package for Social Science software (SPSS version 22.0; Inc. Chicago, IL).

RESULTS

For this study, 150 severe obese subjects (80% females) with mean age of 47.2 ± 10.4 years were selected. The means of systolic and diastolic pressure were 122.2 ± 15.4 mmHg and 78.1 ± 10.0 mmHg, respectively. It was identified that 94.7% of patients presented grade III obesity, 73.3% systemic arterial hypertension, 25% high glycemic levels and 26.4% hypertriglyceridemia. Indeed, abdominal obesity was present in all patients. Thus, taking into account the criteria established by the NCEP, 72.7% of the patients selected for this study present metabolic syndrome.

Genotypic distributions of the polymorphisms rs505151 ($X^2 = 12.33$, $p = 0.92$), rs7566605 ($X^2 = 1.72$, $p = 0.64$) and rs9939609 ($X^2 = 0.82$, $p = 0.49$) exhibited the Hardy-Weinberg equilibrium. The allelic and genotype frequencies of the polymorphisms are presented in table I. The genotypes AA homozygous for PCSK9 gene, CG heterozygote for INSIG2 gene prevailed. On the other hand, there was a similar distribution between the AA and TT genotypes for FTO gene.

Tables II, III and IV show the anthropometric and metabolic variables for DM and RM for polymorphisms of PCSK9, FTO and INSIG2 gene, respectively. No difference was observed between the possible genotypes for PCSK9 and FTO gene. However, there was a difference between systolic and diastolic BP and triglycerides levels among the possible genotypes for INSIG2 gene, showing that individuals with at least one variant allele (G) had higher BP and triglycerides values. Linear regression analyses confirm the association between the polymorphism of INSIG2 gene and this variables, even when adjusted for gender and age (Table V). However, none of the polymorphisms studied was directly related to the risk of MS developing in patients with severe obesity (Fig. 1).

DISCUSSION

No direct relationship was found between the presence of polymorphisms rs505151 of PCSK9 gene, rs7566605 of INSIG2 gene and rs9939609 of FTO gene and the presence of MS. However, it was observed that rs7566605 of INSIG2 gene showed a significant relationship with three parameters analyzed: systolic blood pressure, diastolic blood pressure and triglyceride levels.

Although our findings do not show a direct relationship between PCSK9 polymorphism and BMI or abdominal obesity, some cases have already been well described in the literature (28,32). Moreover, a Chinese study showed an association between the variant allele and HDL-cholesterol and LDL-cholesterol concentrations (2). Evidence supports that in patients with the polymorphism, the expression of the protein was higher, and consequently the circulating LDL-cholesterol concentrations were increased (15). Furthermore, a study investigating the effect of this polymorphism on the risk of cardiovascular diseases (CVD) and stroke showed that in patients with a higher risk of CVD who had the variant allele, LDL-cholesterol concentrations were significantly higher (28).

Regarding the polymorphism of the INSIG2 gene, the literature is still controversial. Our findings corroborate previous studies that did not evidence associations of polymorphism with BMI or waist circumference (1). In contrast, Oki et al. (2009) evidenced that CC genotype is a protective genetic factor against the progression of hypercholesterolemia. The polymorphism of INSIG2 gene does not inhibit fatty acid synthesis, leading to an exaggerated production of cholesterol, fatty acids, triglycerides and phospholipids (3). Other studies found no association between polymorphism and severe obesity, or with phenotypes associated with hypertension (5,13). Thus, the present study is a pioneer to show association between blood pressure levels and this polymorphism. A possible mechanism of increased systolic and diastolic blood pressure would be the fact that the polymorphism increases the production of cholesterol and may contribute for atherosclerosis development, consequently raising the individual's blood pressure. As for the biochemical and metabolic parameters, the wild homozygous form of the INSIG2 (CC) gene was related to the reduction of waist/hip ratio and glycated hemoglobin, which may lead to an improvement in glucose intolerance (6).

The association between the FTO gene and obesity in humans is already well described (7,13). However, the evidence showing the role of this polymorphism in the development of type 2 diabetes mellitus is weak. Li et al. (2008) did not find associations between the gene and waist circumference and fasting glycemia. On the other hand, a recent study by Guclu-Geyik et al. (2016) found an association between polymorphism and MS in men. The mechanism by which the FTO gene leads to the development of obesity is not fully understood. The polymorphism of FTO gene was related to obesity, but not to MS; nonetheless, a predisposition to an increased risk of obesity may trigger other variables such as increased AC, altered fasting glycemia and dyslipidemia.

Thus, as a main finding, despite not being directly associated with the prevalence of MS, the presence of the variant allele of INSIG2

polymorphism could increase the chances of developing the disease. It should be emphasized that our sample size is limited and our analysis was performed in a population of mixed ethnic origin. Low power may have been responsible for lack of significant associations among severe obese individuals, and reaching significance for some of the potential interactions. Moreover, although adjustment by sex and age was performed, residual confounding such as physical active and dietary intake cannot be totally ruled out.

CONCLUSION

Polymorphisms of PCSK9, INSG2 and FTO genes are not associated with the MS prevalence; however, the polymorphism of INSIG2 gene is associated with higher concentrations of triglycerides and blood pressure values, which are also considered as risk factors for the development of the disease.

ACKNOWLEDGMENTS

This work was supported by grants #[2015/20513-8](#) from the São Paulo Research Foundation (FAPESP).

ETHICAL APPROVAL

All procedures performed in this study were in accordance with the ethical standards of the institutional research committee and with the Declaration of Helsinki of 1964 and its later amendments or comparable ethical standards.

REFERENCES

1. Apal Sammy YD, Moy FM, Rampal S, Bulgiba A, Mohamed Z. Genetic associations of the INSIG2 rs7566605 polymorphism with obesity-related metabolic traits in Malaysian Malays. *Genet Mol Res* 2014;13(3):4904-10. DOI: 10.4238/2014.July.4.4
2. Aung LHH, Yin RX, Miao L, Hu XJ, Yan TT, Cao XL, et al. The proprotein convertase subtilisin/kexin type 9 gene E670G polymorphism and serum lipid levels in the Guangxi Bai Ku Yao and

Han populations. *Lipids Health Dis* 2011;10:5. DOI: 10.1186/1476-511X-10-5

3. Baylin A, Deka R, Tuitele J, Viali S, Weeks DE, McGarvey ST. INSIG2 variants, dietary patterns and metabolic risk in Samoa. *Eur J Clin Nutr* 2013;67(1):101-7. DOI:10.1038/ejcn.2012.124

4. Bhowmik B, Afsana F, Siddiquee T, Munir SB, Sheikh F, Wright E, et al. Comparison of the prevalence of metabolic syndrome and its association with diabetes and cardiovascular disease in the rural population of Bangladesh using the modified National Cholesterol Education Program Expert Panel Adult Treatment Panel III and International Diabetes Federation definitions. *J Diabetes Investig* 2015;6(3):280-8. DOI: 10.1111/jdi.12268

5. Bressler J, Fornage M, Hanis CL, Kao WH, Lewis CE, McPherson R, et al. The INSIG2 rs7566605 genetic variant does not play a major role in obesity in a sample of 24,722 individuals from four cohorts. *BMC Med Genet* 2009;10:56. DOI: 10.1186/1471-2350-10-56

6. Burgdörfer E, Korenkov M, Jonas D, Weise D, Haaf T, Zechner U, et al. FTO and INSIG2 genotyping combined with metabolic and anthropometric phenotyping of morbidly obese patients. *Mol Syndromol* 2013;4(6):273-9. DOI: 10.1159/000353563

7. Fang H, Li Y, Du S, Hu X, Zhang Q, Liu A, et al. Variant rs9939609 in the FTO gene is associated with BMI among Chinese children. *BMC Med Genet* 2010;11:136. DOI: 10.1186/1471-2350-11-136

8. Fornage M, Papanicolaou G, Lewis CE, Boerwinkle E, Siscovick DS. Common INSIG2 polymorphisms are associated with age-related changes in body size and high-density lipoprotein cholesterol from young adulthood to middle age. *Metabolism* 2010;59(8):1084-91. DOI: 10.1016/j.metabol.2009.11.005

9. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972;18(6):499-502.

10. Guclu-Geyik F, Onat A, Yuzbasiogullari AB, Coban N, Can G, Lehtimäki T, et al. Risk of obesity and metabolic syndrome associated with FTO gene variants discloses clinically relevant gender

- difference among Turks. *Mol Biol Rep* 2016;43(6):485-94. DOI: 10.1007/s11033-016-3992-0
11. Guo YL, Zhang W, Li JJ. PCSK9 and lipid lowering drugs. *Atheroscler Suppl* 2015;18:21-7. DOI: 10.1016/j.cca.2014.07.008
12. [Hinney A](#), [Herrfurth N](#), [Schonnop L](#), [Volckmar AL](#). Genetic and epigenetic mechanisms in obesity. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2015;58(2):154-8. DOI: 10.1007/s00103-014-2094-1
13. Hotta [K](#), [Nakamura M](#), Nakata Y, Matsuo T, Kamohara S, Kotani K, et al. INSIG2 gene rs7566605 polymorphism is associated with severe obesity in Japanese. *J Hum Genet* 2008;53(9):857-62. DOI: 10.1007/s10038-008-0317-8
14. Hubáček JA, Kuthanova L, Bohuslavova R, Adámková V, Lánská V, Meitinger T, et al. INSIG2 promoter variant, obesity markers and lipid parameters - No association in a large Slavonic Caucasian population Sample. *Folia Biol (Praha)* 2010;56(3):131-4.
15. Ibarretxe D, Girona J, Plana N, Cabré A, Ferré R, Amigó N, et al. Circulating PCSK9 in patients with type 2 diabetes and related metabolic disorders. *Clin Investig Arterioscler* 2016;28(2):71-8. DOI: 10.1016/j.arteri.2015.11.001
16. Kaur J. A comprehensive review on metabolic syndrome. *Cardiol Res Pract* 2014;2014:943162. DOI: 10.1155/2014/943162
17. Li HX, Wu Y, Loos RJF, Hu FB, Liu Y, Wang J, et al. Variants in the fat mass- and obesity-associated (FTO) gene are not associated with obesity in a Chinese Han population. *Diabetes* 2008;57(1):264-8. DOI: 10.2337/db07-1130
18. [Martínez-Hernández A](#), [Córdova EJ](#), [Rosillo-Salazar O](#), [García-Ortiz H](#), [Contreras-Cubas C](#), [Islas-Andrade S](#), et al. Association of HMOX1 and NQO1 polymorphisms with metabolic syndrome components. *PLoS One* 2015;10(5):e0123313. DOI: 10.1371/journal.pone.0123313
19. [Moleres A](#), [Milagro FI](#), [Marcos A](#), [González-Zorzano E](#), [Campoy C](#), [Garagorri JM](#), et al. Common variants in genes related to lipid and energy metabolism are associated with weight loss after an intervention in overweight/obese adolescents. *Nutr Hosp* 2014;30(1):75-83. DOI: 10.3305/nh.2014.30.1.7542

20. Moreira GC, Cipullo JP, Ciorlia LAS, Cesarino CB, Vilela-Martin JS. Prevalence of metabolic syndrome: association with risk factors and cardiovascular complications in an urban population. *PLoS One* 2014;9(9):e105056. DOI: 10.1371/journal.pone.0105056
21. Nicoletti CF, De-Oliveira APRP, Brochado MJF, Oliveira BAP, Pinhel MA, Marchini JS, et al. UCP1 -3826 A>G polymorphism affects weight, fat mass, and risk of type 2 diabetes mellitus in grade III obese patients. *Nutrition* 2016;32:83-7. DOI: 10.1016/j.nut.2015.07.016
22. [Nilsson E](#), [Jansson PA](#), [Perfilyev A](#), [Volkov P](#), [Pedersen M](#), [Svensson MK](#), et al. Altered DNA methylation and differential expression of genes influencing metabolism and inflammation in adipose tissue from subjects with type 2 diabetes. *Diabetes* 2014;63(9):2962-76. DOI: 10.2337/db13-1459
23. Oliveira JS, Boery RNSO. An integrative review of associations between polymorphic variants and the metabolic syndrome. *J Vasc Bras* 2018; 17(2):141-147. DOI: 10.1590/1677-5449.007917
24. Oki K, Yamane K, Kamei N, Asao T, Awaya T, Kohno N, et al. The single nucleotide polymorphism upstream of insuline-induced gene 2 (INSIG2) is associated with the prevalence of hypercholesterolemia, but not with obesity, in Japanese American women. *Br J Nutr* 2009;101(3):322-7. DOI: 10.1017/S0007114508006557
25. O'Neill S, O'Driscoll L. Metabolic syndrome: a closer look at the growing epidemic and its associated pathologies. *Obes Rev* 2015;16(1):1-12. DOI: 10.1111/obr.12229
26. Oki K, Yamane K, Kamei N, Asao T, Awaya T, Kohno N, et al. The single nucleotide polymorphism upstream of insuline-induced gene 2 (INSIG2) is associated with the prevalence of hypercholesterolemia, but not with obesity, in Japanese American women. *Br J Nutr* 2009;101(3):322-7. DOI: 10.1017/S0007114508006557
27. Pena SD, Bastos-Rodrigues L, Pimenta JR, Bydlowski SP. DNA tests probe the genomic ancestry of Brazilians. *Braz J Med Biol Res* 2009;42(10):870-6. DOI: 10.1590/S0100-879X2009005000026
28. [Slimani A](#), [Harira Y](#), [Trabelsi I](#), [Joma W](#), [Maatouk F](#), [Hamda KB](#), et al. Effect of E670G Polymorphism in PCSK9 gene on the risk and

severity of coronary heart disease and ischemic stroke in a Tunisian cohort. *J Mol Neurosci* 2014;53(2):150-7. DOI: 10.1007/s12031-014-0238-2

29. Xiao-Ying D, Qiu TS. Insulin-induced gene: a new regulator in lipid metabolism. *J Peptides* 2010;31(11):2145-50. DOI: 10.1016/j.peptides.2010.07.020

30. Yang Q, Xiao T, Guo J, Su Z. Complex relationship between obesity and the fat mass and obesity locus. *Int J Biol Sci* 2017;15;13(5):615-29. DOI: 10.7150/ijbs.17051

31. Yang Y, Liu B, Xia W, Yan J, Liu H, Hu L, et al. [FTO](#) genotype and type 2 diabetes mellitus: spatial analysis and meta-analysis of 62 case-control studies from different regions. *Genes (Basel)* 2017;8(2):70. DOI: 10.3390/genes8020070

32. Yin RX, Wu DF, Miao L, Htet-Aung LH, Cao XL, Yan TT, et al. Interactions of several single nucleotide polymorphisms and high body mass index on serum lipid traits. *Biofactors* 2013;39(3):315-25. DOI: 10.1002/biof.1073

Nutrición
Hospitalaria

Table I. Genotypic and allelic frequency of polymorphisms assessed in severe obese patients evaluated in the study (n = 150)

<i>Polymorphism</i>	<i>Gene</i>	<i>Genotype frequency</i>		<i>Allele frequency</i>		
		<i>Genotype</i>	<i>n</i>	<i>%</i>	<i>Allele</i>	
rs 505151	<i>PCSK9</i>	AA	126	86.9%	Allele A	0.92
		AG	15	10.3%	Allele G	0.08
		GG	4	2.8%		
rs 7566605	<i>INSIG2</i>	CC	53	38.7%	Allele C	0.64
		CG	70	51.1%	Allele G	0.36
		GG	14	10.2%		
rs 9939609	<i>FTO</i>	AA	37	25.5%	Allele A	0.49
		AT	67	46.2%	Allele T	0.51
		TT	41	28.3%		

n: number of individuals.

Table II. Anthropometric and metabolic variables for genotypes of the rs505151 polymorphism in the PCSK9 gene in subjects with severe obesity (n = 150)

<i>Variables</i>	<i>Dominant model</i>		<i>Recessive model</i>	
	AA	AG + GG	AA + AG	GG
Weight (kg)	137.5 ± 23.5	140.9 ± 23.3	138.0 ± 23.2	147.8 ± 31.2
BMI (kg/m ²)	51.4 ± 7.5	50.7 ± 6.1	51.5 ± 7.4	49.1 ± 5.2
AC (cm)	144.7 ± 16.8	137.3 ± 21.4	144.6 ± 17.1	121.0 ± 5.6
Systolic BP (mmHg)	121.5 ± 15.1	122.6 ± 14.1	122.1 ± 15.6	117.5 ± 9.6
Diastolic BP (mmHg)	77.9 ± 10.2	78.7 ± 9.1	78.2 ± 10.1	75.0 ± 5.8
Total cholesterol (mg/dl)	185.9 ± 35.6	177.8 ± 28.2	185.3 ± 35.0	160.0 ± 15.5
LDL-cholesterol (mg/dl)	115.8 ± 32.1	110.1 ± 22.9	115.2 ± 31.3	104.7 ± 14.8
HDL-cholesterol (mg/dl)	40.6 ± 10.0	40.2 ± 6.3	40.6 ± 9.5	34.5 ± 4.1
Triglycerides (mg/dl)	151.0 ± 89.8	136.4 ± 59.0	150.6 ± 87.0	102.5 ± 17.5
Glycemia (mg/dl)	101.6 ± 31.1	111.6 ± 56.5	103.6 ± 35.6	89.7 ± 17.3

BMI: body mass index; AC: abdominal circumference; BP: blood pressure; LDL-cholesterol: low density lipoprotein; HDL-cholesterol: high density lipoprotein.

Table III. Anthropometric and metabolic variables for genotypes of the rs9939609 polymorphism of FTO gene in severe obese subjects evaluated in the study (n = 150)

<i>Variables</i>	<i>Dominant model</i>		<i>Recessive model</i>	
	AA	AT + TT	AA + AT	TT
Weight (kg)	137.4 ± 24.5	138.3 ± 23.3	136.7 ± 22.9	142.1 ± 24.7
BMI (kg/ m ²)	51.8 ± 7.2	51.3 ± 7.5	50.8 ± 7.1	52.9 ± 7.9
AC (cm)	143.9 ± 18.6	144.0 ± 16.9	142.8 ± 16.2	146.5 ± 19.6
Systolic BP (mmHg)	123.3 ± 18.2	121.3 ± 14.0	121.6 ± 15.0	122.5 ± 15.6
Diastolic BP (mmHg)	79.4 ± 12.4	77.6 ± 9.1	77.6 ± 9.9	79.2 ± 10.5
Total cholesterol (mg/dl)	186.7 ± 28.5	184.7 ± 36.4	186.1 ± 33.9	182.7 ± 36.2
LDL-cholesterol (mg/dl)	117.4 ± 25.5	114.6 ± 32.7	117.2 ± 29.0	110.5 ± 35.3
HDL-cholesterol (mg/dl)	40.0 ± 9.8	40.7 ± 9.6	40.9 ± 9.4	39.7 ± 10.2
Triglycerides (mg/dl)	165.9 ± 110.5	143.7 ± 75.5	150.8 ± 92.8	145.9 ± 66.4
Glycemia (mg/dl)	109.0 ± 42.2	100.9 ± 32.7	101.3 ± 30.9	107.4 ± 45.2

BMI: body mass index; AC: abdominal circumference; BP: blood pressure; LDL-cholesterol: low density lipoprotein; HDL-cholesterol: high density lipoprotein

Table IV. Anthropometric and metabolic variables for genotypes of polymorphism rs7566605 of INSIG2 gene in severe obese evaluated in this study (n = 150)

<i>Variables</i>	<i>Dominant model</i>		<i>Recessive model</i>	
	CC	CG + GG	CC + CG	GG
Weight (kg)	133.0 ± 22.8	141.0 ± 23.7	138.2 ± 23.4	134.7 ± 25.8
BMI (kg/m ²)	51.4 ± 7.4	51.3 ± 7.4	51.5 ± 7.3	50.2 ± 8.4
AC (cm)	143.6 ± 17.8	143.6 ± 16.0	144.1 ± 16.7	135.5 ± 17.2
Systolic BP (mmHg)	116.4 ± 14.7	125.6 ± 15.2*	121.2 ± 15.2	129.2 ± 17.8
Diastolic BP (mmHg)	75.0 ± 9.8	79.9 ± 9.9*	77.8 ± 10.0	80.0 ± 11.3
Total cholesterol (mg/dl)	179.9 ± 35.9	186.9 ± 33.4	183.6 ± 35.2	188.3 ± 27.9
LDL-cholesterol (mg/dl)	109.1 ± 32.2	117.7 ± 28.9	113.1 ± 30.6	124.8 ± 27.2
HDL-cholesterol (mg/dl)	42.6 ± 9.8	39.5 ± 9.3	40.9 ± 9.9	39.3 ± 8.2
Triglycerides (mg/dl)	132.2 ± 54.9	160.1 ± 102.4*	148.9 ± 85.9	150.6 ± 106.1
Glycemia (mg/dl)	98.1 ± 34.2	106.1 ± 36.2	102.4 ± 36.2	102.4 ± 36.2

BMI: body mass index; AC: abdominal circumference; BP: blood pressure; LDL-cholesterol: low density lipoprotein; HDL-cholesterol: high density lipoprotein. *: $p < 0.05$

Table V. Multiple linear regression analysis, adjusted for sex and anthropometric variables, showing the contribution of the polymorphism rs 7566605 in the INSIG2 gene in the concentrations of triglycerides, systolic and diastolic pressure in subjects with severe obesity (n = 150)

Variables	β	r^2	p	IC 95%
Triglycerides				
		0.068		
rs 7566605	0.275		0.044	(1.186; 79.430)
Sex	-0.015		0.915	(-67.751; 60.915)
Weight (kg)	- 0.353		0.050	(-2.431; -0.002)
AC (cm)	0.144		0.371	(-0.774; 2.036)
Systolic pressure				
		0.078		
rs 7566605	0.263		0.002	(2.999; 13.702)
Sex	-0.034		0.724	(-8.917; 6.211)
Age	0.101		0.239	(-0.101; 0.400)
Weight (kg)	0.145		0.144	(-0.033; 0.226)
Diastolic pressure				
		0.087		
rs 7566605	0.196		0.022	(0.589; 7.505)
Sex	-0.011		0.907	(-5.177; 4.598)
Age	0.211		0.015	(0.041; 0.364)
Weight (kg)	0.173		0.080	(-0.009; 0.159)

AC: abdominal circumference; CI: confidence interval.




Fig. 1. Odds ratio (95% CI) of the association between polymorphisms rs505151 of PCSK9 gene, rs7566605 of INSG2 gene and rs9939609 of FTO gene and the metabolic syndrome. MD: dominant model; MR: recessive model.

Nutrición
Hospitalaria