

**Comparación de métodos de
estimación del peso en niños y
adolescentes brasileños
hospitalizados**

**Comparison of weight estimation
methods in hospitalized Brazilian
children and adolescents**

OR 2594

Comparison of weight estimation methods in hospitalized Brazilian children and adolescents

Comparación de métodos de estimación del peso en niños y adolescentes brasileños hospitalizados

Catarina Tenório de Cerqueira, Maria Júlia Gonçalves de Mello, Larissa de Andrade Viana, Derberson José do Nascimento Macêdo, and José Natal Figueiroa

Nutrition Department. Instituto de Medicina Integral Prof. Fernando Figueira. Boa Vista, Recife. Pernambuco, Brazil

Received: 22/03/2019

Accepted: 23/07/2019

Correspondence: Catarina Tenório de Cerqueira. Departamento de Nutrição. Instituto de Medicina Integral Prof. Fernando Figueira. Rua dos Coelhos 300. Boa Vista. 50070-550 Recife, PE, Brazil

e-mail: catarinatcq@hotmail.com

ABSTRACT

Introduction: in emergencies or in situations involving critically ill patients, an accurate calculation of body weight is essential to ensure adequate medical care. Generally, simple techniques are used to determine weight.

Aim: to evaluate the weight estimation methods Advanced Paediatric Life Support (APLS) Update, Luscombe and Owens, Best Guess, and the Broselow tape, comparing measured weight with the weight estimated by each method in hospitalized Brazilian children and adolescents.

Methods: an observational study in which anthropometric data were collected from patients of 0-14 years of age between August 2016 and

January 2017. The paired t-test was used to compare the patients' measured weight with their estimated weight. The accuracy of each method was analyzed using Bland-Altman plots and Lin's concordance correlation coefficient.

Results: the sample consisted of 446 patients. Most of the patients were male (58.1%), of 1-5 years of age (43%) and well nourished (85%). For those under 12 months of age, the APLS Update and Best Guess methods performed best, with the difference in means of measured weight and estimated weight being 0.4 kg ($p = 0.183$) and -0.2 kg ($p = 0.140$), respectively. In the 1-5 years of age group only the APLS Update yielded satisfactory results (0.2 kg; $p = 0.200$). The best agreement with measured weight, according to the Bland-Altman plots and Lin's coefficient, was found for the Broselow tape ($CC = 0.96$).

Conclusion: of the estimation methods evaluated the Broselow tape was the most accurate one. Further studies are required to adapt this method for use in the Brazilian population, thus ensuring its appropriate application in this country.

Keywords: Paediatric. Body weight. Measurements. Methods and theories.

RESUMEN

Introducción: en las emergencias o en las situaciones que involucran a pacientes críticamente enfermos, el cálculo preciso del peso corporal es esencial para garantizar una atención médica adecuada. En general se utilizan técnicas simples para determinar el peso.

Objetivo: evaluar los métodos de estimación del peso Advanced Paediatric Life Support (APLS) Update, Luscombe y Owens, Best Guess y la cinta Broselow, comparando el peso medido con el peso estimado por cada método en los pacientes hospitalizados.

Métodos: estudio observacional en el que se recopilaron datos antropométricos de pacientes de 0 a 14 años de edad entre agosto de 2016 y enero de 2017. Se utilizó la prueba t pareada para comparar el

peso medido de los pacientes con su peso estimado. La precisión de cada método se analizó mediante gráficos de Bland-Altman y el coeficiente de correlación de concordancia de Lin.

Resultados: la muestra estuvo constituida por 446 pacientes. La mayoría eran varones (58,1%) de 1 a 5 años (43%) y bien nutridos (85%). Para los menores de 12 meses de edad, los métodos APLS Update y Best Guess obtuvieron los mejores resultados, siendo la diferencia entre las medias de peso medido y peso estimado de 0,4 kg ($p = 0,183$) y -0,2 kg ($p = 0,140$), respectivamente. En el grupo de 1 a 5 años, solo el método APLS Update arrojó resultados satisfactorios (0,2 kg; $p = 0,200$). El mejor acuerdo con el peso medido, según las gráficas de Bland-Altman y el coeficiente de Lin, se encontró al utilizar la cinta Broselow ($CC = 0,96$).

Conclusión: de los métodos de estimación evaluados, la cinta Broselow fue la más precisa. Se requieren estudios adicionales para adaptar este método para su uso en la población brasileña.

Palabras clave: Pediatría. Peso corporal. Mediciones. Métodos y teorías.

INTRODUCTION

In emergencies or in situations involving critically ill patients, the accurate calculation of body weight is essential to ensure adequate medical care (1). Indeed, weight is not only associated with the patient's nutritional status, but also forms a basis on which to calculate intravenous fluid volumes and drug doses, as well as on which to determine the dimensions of the equipment used in resuscitation (1,2). In no other patient population is there a greater need to calculate and manipulate these doses than in the paediatric population. Therefore, recording the patient's correct weight plays a crucial role in minimizing drug dosage errors (3).

Simple techniques are generally used to determine weight (3). In some circumstances, however, an anthropometric evaluation is impossible and other people able to supply the necessary information are not always present. In critical and/or bedridden patients, these measurements are rarely taken in view of the difficulty involved in applying the conventional

methods of determining the correct weight. Attempting to estimate weight solely by visual observation is a practice that is not recommended and that may result in serious errors in drug administration (4).

Various tools aimed at enabling weight to be estimated quickly have been proposed in the literature. The most commonly used methods are based on the age or the length of the patient. Based on the patient's age are methods such as APLS (Advanced Paediatric Life Support), Best Guess, and those adapted by Luscombe and Owens. The Broselow tape proposed in the Pediatric Advanced Life Support (PALS) manual (5-8) is an example of a method based on length. In a review of the literature on this subject, Young and Korotzer argued that the ideal method of estimating weight should be precise, reliable, quickly obtainable, simple to use, and readily available to healthcare professionals (9).

Several studies comparing weight estimation methods have been published, with results varying as a function of the different populations and ethnic groups evaluated (2,4,10,11). A study conducted in hospitalized paediatric patients in Australia compared measured weight with the weight estimated using different equations. The methods that proved most accurate in that study population were the APLS Update for patients under 12 months of age, and the Best Guess for those over a year old (10).

A review of the literature showed that few studies have been conducted to evaluate the precision of these weight estimation methods in the Brazilian paediatric population. The objective of the present study was to evaluate these weight estimation methods, comparing estimated weight with measured weight in children and adolescents receiving care at a Brazilian reference hospital.

MATERIALS AND METHODS

Study design and site

This observational study was performed in a paediatric clinic at the *Instituto de Medicina Integral Professor Fernando Figueira* (IMIP) between August 2016 and January 2017. IMIP, situated in the Brazilian state of Pernambuco, is a philanthropic quaternary hospital dedicated exclusively to the care of patients within the public healthcare system. With 1,066

beds, the hospital includes 214 paediatric beds and 6 intensive care units (ICUs), 2 of which are paediatric. During the data collection period of this study, 900 paediatric patients on average were admitted to this hospital each month.

Population & study sample

Data were collected from hospitalized patients of both sexes under 14 years of age, who were able to walk and/or on whom it was possible to take their anthropometric measurements. Patients with encephalopathy, amputees or patients whose limbs were disproportional, those with genetic syndromes, oedematous patients, and any who had been included in the study previously (readmissions to hospital) were excluded from the study.

Data collection

Following an interview and after the informed consent had been obtained from the parents/guardians or from the child itself, anthropometric measurements were performed in accordance with the techniques recommended by the World Health Organization (WHO) (12,13). Measurements were taken in duplicate, sequentially, by the same investigator, using standardized procedures. In the analysis, in order to maximize quality control, thus improving reliability and accuracy, the mean value of the two measurements was used as the measured weight. Weight was measured using a digital electronic scale (Welmy, Brazil) and length/height was measured using an infantometer (Welmy, Brazil) for children two years of age or under, and a stadiometer (Tonelli, Brazil) for those over two years of age.

Ethnicity was self-reported by each individual and the overall sample was then dichotomized into white or non-white.

Weight was estimated using the following formulae:

APLS Update (5)

- 0-12 months: weight (kg) = (0.5 x age in months) + 4
- 1-5 years: weight (kg) = (2 x age in years) + 8
- 6-12 years weight (kg) = (3 x age in years) + 7

Luscombe and Owens (6)

- 1-14 years: weight (kg) = (3 x age in years) + 7

Best Guess (7)

- 1-11 months: weight (kg) = (age in months + 9)/2
- 1-4 years: weight (kg) = 2 x (age in years + 5)
- 5-14 years: weight (kg) = 4 x age in years

Broselow Tape (8)

A version of the Broselow tape produced by PediaTape was used. This version is based on the combined data obtained from the databases of the 2007 National Health and Nutrition Examination Survey (NHANES) in the United States. The tape is divided into colour zones that estimate the patient's weight in accordance with his/her length. To perform the estimation, the patient lies down, barefoot, with his/her arms extended along their body and with their legs stretched out. According to the manufacturer's recommendations, the tape was applied to paediatric patients of up to 144 cm in length and/or 37 kg. In the case of obese individuals, it is recommended that the estimated weight should correspond to the next higher colour zone. Estimated weight for the later analysis of agreement was defined as the mean point on each color-coded range of weight.

Data analysis

The STATA software program, version SE12.0, and Excel 2010 were used to store, process and analyse the data. The Anthro and the Anthro Plus software program was used to calculate the z-scores in accordance with the criteria proposed by the WHO (12,13). Patients whose BMI-for-age z-score was between -2 and +2 were considered to be well nourished.

Student's t-test for paired samples was used to compare the mean measured weight with the mean weight estimated by each method. A significance level of 5% was established for the tests applied.

The Bland-Altman method (14) and Lin's concordance correlation coefficient (15) were used to analyse concordance. The Bland-Altman

method generates graphs that show the average value of measured and estimated weight on the horizontal axis, and the differences between them on the vertical axis. This technique provides the bias, precision (standard deviation [SD] of the bias) and the limits of concordance according to the 95% confidence interval (95% CI) ($\text{bias} \pm 1.96 \text{ SD}$). A positive bias indicates underestimation by the method, while a negative bias indicates overestimation.

As proposed by McBride, Lin's concordance correlation coefficient should be interpreted according to the position of the lower limit of the 95% CI for this coefficient as follows: < 0.90 = poor; $0.90-0.95$ = moderate; $0.95-0.99$ = good; and > 0.99 = excellent (15).

The concordance tests were initially applied to the entire sample and subsequently to groups classified according to nutritional status, bearing in mind that one of the methods, the Broselow tape, requires a different estimation to be used for obese patients. However, considering the small percentage of dystrophic patients in the sample, and in order to preserve the pre-established criteria equally for each method, it was decided to present the results obtained only from the group of well-nourished patients.

Ethical issues

The internal review board of IMIP approved the study protocol under reference CAAE 51877115.0.0000.5201. Patients were admitted to the study following authorization from their parents or guardians, who were required to sign an informed consent form. In addition, participants of 8 years of age or more were asked to sign an assent form.

RESULTS

Participants

During the data collection period, 465 hospitalized patients fulfilled the eligibility criteria. Of these, 19 (4.1%) refused to participate or did not agree to have their anthropometric measurements taken for the study. Therefore, the final sample consisted of the 446 individuals whose clinical and sociodemographic data are described in table I. Most of the included

children and adolescents (58.1%) were male, of 1-5 years of age (43%), and well nourished (85%).

Measured and estimated weight

The mean measured weight of all the patients evaluated was 21.8 kg (\pm 13.8 kg). The means and standard deviations for each weight estimation method according to age group, sex, and ethnicity are described in table II. For children under 12 months of age, there were no statistically significant differences between the means of measured weight and estimated weight according to the APLS Update and Best Guess ($p = 0.183$ and $p = 0.140$, respectively). For patients of 1-5 years of age, there were no statistically significant differences between the means of measured and estimated weight with APLS Update ($p = 0.200$); however, the differences found with the Luscombe and Owens and Best Guess methods were statistically significant ($p < 0.001$). In the case of the children over five years of age, statistically significant differences were found between measured and estimated weight ($p < 0.001$) for all of the methods evaluated. The Broselow tape was not included in this analysis because it does not take age into account when estimating weight. The Broselow tape was applied to 383 patients whose length was within the limit of its extension (144 cm), and in 61% of cases the measured weight was in agreement with the weight range estimated by the tape.

Figures 1, 2 and 3 are graphic representations of the Bland-Altman plot analysis. In the group of patients under 12 months of age, the bias was minimal both with the APLS Update and with the Best Guess, although the concordance limits indicated a considerable difference between measured weight and the weight estimated by each method for this age group (95% CI: -1.9 to 2.1; 95% CI: -2.3 to 1.5, respectively), as shown in table III.

Lin's concordance correlation coefficient was also applied to the sample and the values found are shown in table III. Concordance was best with the Broselow tape, with a coefficient of 0.96 and a lower limit of 0.94, considered a moderate correlation.

DISCUSSION

Although weight estimation methods are widely used worldwide in paediatric emergency departments, to the best of our knowledge no studies have been published to validate their use in the Brazilian population. The agreement between the age-based weight estimation method and the measured weight of the children and adolescents in the present study was not good. In general, agreement was poorest when the method proposed by Luscombe and Owens was used. On the other hand, the Broselow tape, which is based on patient length, was the method in which the estimated weight was closest to the patient's measured weight. The performance of the methods deteriorated as the age of the participants in the study increased.

In the case of patients under 12 months of age, the results obtained with the method proposed by the APLS Update and by Best Guess were satisfactory in that the mean measured weight was similar to the mean estimated weight. Similar findings were reported from a study involving around 30,000 patients at the Children's Hospital at Westmead, a tertiary hospital in Australia (10), with the APLS Update proving even more specific than the Best Guess (a difference between means of 0.5 kg and 0.9 kg) (10). Flannigan et al. retrospectively evaluated around 10,000 patients in the United Kingdom and also found that the APLS Update gave a more accurate estimation of weight for patients under 12 months of age ($p = 0.189$) (16).

For the group of patients of 1-5 years of age in the present study, agreement was best with the APLS Update, with only 0.2 kg of difference between measured and estimated mean weight. The APLS Update introduced three new methods for three different age groups instead of using one single method for all ages, as previously recommended (5). Results have shown that agreement is more satisfactory with the new version compared to the previous one, particularly in patients under five years of age (9,10).

As age increased, statistically significant differences began to appear between measured and estimated weight when using these mathematical methods (17). In a study conducted by Chavez et al. to evaluate the effectiveness of four weight estimation methods in 324 patients at a paediatric emergency department in Florida, USA, the difference between

mean weights was also found to be directly proportional to the age of the individual (18). This finding agrees with the results from other studies showing that the methods that are based exclusively on patient age do not appear to be sufficiently accurate to enable their widespread use in paediatric emergency departments (1,4,9,16,19).

The percentage of participants in the present study whose weight was correctly estimated using the Broselow tape is close to that reported by Mishra et al., who evaluated around 500 children of up to ten years of age receiving outpatient care at a hospital in India (20). Agreement was found between measured and estimated weight according to the colour zone proposed by the tape in 63.2% of the sample (20). These results are more satisfactory than those obtained by other studies, in which the mean percentage of accuracy was below 50% (10,16,18,21).

The analyses of concordance evaluated using the Bland-Altman plot technique in well-nourished patients showed that, although in some groups there was a minimum bias that was directly proportional to an increase in age, there is an important dispersion in the values, and the limit of the differences was unacceptable. In the patients under 12 months of age, for example, although the bias was small, the limit of the difference between means is of approximately two kilograms, both in the group whose weight was estimated using the APLS Update and in that in which Best Guess was used. In our opinion, an error of two kilograms in patients who are only months old would correspond to a significant percentage of their body weight and could cause serious errors in their clinical management.

Lin's concordance correlation coefficient was also used, and its results reinforce those found with the Bland-Altman plots, in which agreement with measured weight, considered the gold standard, is poor with all the age-based methods. The imprecision of these methods was recently described by Wells et al. (19), who evaluated 28 estimation methods in a sample of around 1,000 children aged from one month to sixteen years admitted to four different paediatric emergency departments in Johannesburg, South Africa. None of the mathematical methods analysed presented satisfactory statistical results that would justify their use in that study population. To draw their conclusion, those authors also took

studies conducted to validate these methods into consideration, reporting on the performance of each one in the published paper (20).

The results obtained with the Broselow tape, nevertheless, were best when compared to the mathematical methods, confirming that the effectiveness of the length-based method is better than the effectiveness of those approaches that take only age into consideration (4,9,16,21-23). This finding agrees with the results reported by Geduld et al., who compared weight estimation methods in approximately 2,000 children. In that sample, agreement with measured weight was better when the Broselow tape was used (0.9% of difference between mean measurements) when compared to Best Guess and the methods proposed by Luscombe and Owens, which tended to overestimate weight by 15.4% and 12.4%, respectively (24).

Illnesses and hospitalization can result in weight loss, and this could be considered a limitation of the present study. Nevertheless, it was decided that the weight estimation methods should be applied under the actual circumstances in which they would need to be used, i.e., that the study should be performed in a sample of hospitalized patients. Another limitation of the study refers to the fact that the methods analysed here were instituted based on data from populations with different ethnic and anthropometric characteristics when compared to those of this sample. On the other hand, with the objective of identifying possible variations in the performance of the methods assessed, the sample was stratified according to patient nutritional status, evaluated in accordance with the WHO parameters.

Age-based weight estimation methods are imprecise, since the weight of children and adolescents is affected by various other factors in addition to age (1,9,25). In the present sample only the Broselow tape yielded satisfactory results when compared to the other methods evaluated. The age-based methods used were not accurate enough, and their use in this population cannot be endorsed. In this respect, whenever it is impossible to assess weight by the conventional method, and parents are absent or unable to supply this information, the use of weight estimation methods based on patient length is recommended.

REFERENCES

1. Loo PY, Chong SL, Lek N, Bautista D, Ng KC. Evaluation of three paediatric weight estimation methods in Singapore. *J Paediatr Child Health* 2013;49:E311-6. DOI: 10.1111/jpc.12141
2. Omisanjo AO, Orimadegun AE, Akinbami FO. Accuracy of Nelson and Best Guess method in estimation of weights in Nigerian children population. *Ann Ib Postgrad Med* 2014;12:80-8.
3. Abdel-Rahman SM, Ahlers N, Holmes A, Wright K, Harris A, Weigel J, et al. Validation of an improved pediatric weight estimation strategy. *J Pediatr Pharmacol Ther* 2013;18:112-21. DOI: 10.5863/1551-6776-18.2.112
4. O'Leary F, John-Denny B, McGarvey K, Hann A, Pegiazoglou I, Peat J. Estimating the weight of ethnically diverse children attending an Australian emergency department: a prospective, blinded, comparison of age-based and length-based tools including Mercy, PAWPER and Broselow. *Arch Dis Child* 2017;102:46-52. DOI: 10.1136/archdischild-2016-310917
5. Advanced Life Support Group. *Advanced paediatric life support: a practical approach to emergencies*. 5th ed. Oxford: Wiley-Blackwell; 2011.
6. Luscombe M, Owens B. Weight estimation in resuscitation: is the current methods still valid? *Arch Dis Child* 2007;92:412-5. DOI: 10.1136/adc.2006.107284
7. Tinning K, Acworth J. Make your Best Guess: an updated method for paediatric weight estimation in emergencies. *Emerg Med Australas* 2007;19:528-34. DOI: 10.1111/j.1742-6723.2007.01026.x
8. Lubitz DS, Seidel JS, Chameides L, Lutten RC, Zaritsky AL, Campbell FW. A rapid method for estimating weight and resuscitation drug dosages from length in the pediatric age group. *Ann Emerg Med* 1988;17:576-81. DOI: 10.1016/S0196-0644(88)80396-2
9. Young KD, Korotzer NC. Weight estimation methods in children: a systematic review. *Ann Emerg Med* 2016;68:441-51. DOI: 10.1016/j.annemergmed.2016.02.043

10. Graves L, Chayen G, Peat J, O'Leary F. A comparison of actual to estimated weights in Australian children attending a tertiary children's hospital, using the original and updated APLS, Luscombe and Owens, Best Guess method and the Broselow tape. *Resuscitation* 2014;85:392-6. DOI: 10.1016/j.resuscitation.2013.11.024
11. Cattermole GN, Leung PY, Mak PS, Graham CA, Rainer TH. Mid-arm circumference can be used to estimate children's weights. *Resuscitation* 2010;81:1105-10. DOI: 10.1016/j.resuscitation.2010.05.015
12. World Health Organization Multicentre Growth Reference Study Group. WHO child growth standards based on length/height, weight and age. *Acta Paediatr Suppl* 2006;450:76-85. DOI: 10.1111/j.1651-2227.2006.tb02378.x
13. De Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* 2007;85:660-7. DOI: 10.2471/BLT.07.043497
14. Bland JM, Altman D. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986;1:307-10.
15. McBride GB. A proposal for strength-of-agreement criteria for Lin's Concordance Correlation Coefficient. NIWA Client Report: HAM 2005-062; 2005.
16. Flannigan C, Bourke TW, Sproule A, Steveson M, Terris M. Are APLS method for estimating weight appropriate for use in children admitted to PICU? *Resuscitation* 2014;85:927-31. DOI: 10.1016/j.resuscitation.2014.03.313
17. Samerchua A, Suraseranivongse S, Komoltri C. A comparison of pediatric weight estimation methods for emergency resuscitation. *Pediatr Emerg Care* 2017.
18. Chavez H, Peterson RE, Lo K, Arel M. Weight estimation in an inner-city pediatric ED: the effect of obesity. *Am J Emerg Med* 2015;33:1364-7. DOI: 10.1016/j.ajem.2015.07.040

19. Wells M, Goldstein LN, Bentley A. It is time to abandon age-based emergency weight estimation in children! A failed validation of 20 different age-based methods. *Resuscitation* 2017;116:73-83. DOI: 10.1016/j.resuscitation.2017.05.018
20. Mishra DG, Kole T, Nagpal R, Smith JP. A correlation analysis of Broselow™ Pediatric Emergency Tape-determined pediatric weight with actual pediatric weight in India. *World J Emerg Med* 2016;7:40-3. DOI: 10.5847/wjem.j.1920-8642.2016.01.007
21. Tanner D, Negaard A, Huang R, Evans N, Hennes H. A prospective evaluation of the accuracy of weight estimation using the Broselow tape in overweight and obese pediatric patients in the emergency department. *Pediatr Emerg Care* 2017;33:675-8. DOI: 10.1097/PEC.0000000000000894
22. AlHarbi TM, AlGarni A, AlGamdi F, Jawish M, Wani TA, Abu-Shaheen AK. The accuracy of Broselow tape weight estimate among pediatric population. *BioMed Res Int* 2016;2016:7042947. DOI: 10.1155/2016/7042947
23. Meguerdichian MJ, Clapper TC. The Broselow tape as an effective medication dosing instrument: a review of the literature. *J Pediatr Nurs* 2012;27:416-20. DOI: 10.1016/j.pedn.2012.04.009
24. Geduld H, Hodgkinson PW, Wallis LA. Validation of weight estimation by age and length based methods in the Western Cape, South Africa population. *Emerg Med J* 2011;28:856-60. DOI: 10.1136/emj.2010.098640
25. Wells M, Coovadia A, Kramer E, Goldstein L. The PAWPER tape: a new concept tape-based device that increases the accuracy of weight estimation in children through the inclusion of a modifier based on body habitus. *Resuscitation* 2013;84:227-32. DOI: 10.1016/j.resuscitation.2012.05.028

Table I. Frequency distribution of paediatric patients admitted to a clinical ward according to their socioeconomic, demographic, and clinical parameters. IMIP, August 2016 - January 2017

	n	%
Sex (n = 446)		
Male	259	58.1
Female	187	41.9
Age (n = 446)		
< 1 year	55	12.3
1-5 years	192	43.0
6-10 years	125	28.0
11-14 years	74	16.7
Per capita income (n = 397)		
< 0.5 minimum salary*	343	86.4
> 0.5 minimum salary*	54	13.6
Chronic disease (n = 446)		
Yes	114	25.6
No	332	74.4
Nutritional status/BMI-for-age z-score (n = 446)		
< -2	24	5.4
≥ -2 and ≤ +2	379	85

> +2

43

9.6

*Per capita income based on the minimum salary for 2016, which was equivalent to US\$ 266.00.

Nutrición
Hospitalaria

Table II. Mean measured versus mean estimated weight as calculated by the different methods evaluated in paediatric patients admitted to a clinical ward, according to age group, sex, and ethnicity. IMIP, August 2016 - January 2017

		Methods used		
		APLS Update	Luscombe and Owens	Best Guess
Age group (years)				
< 1	Measured	7.2 ± 2.3	-	7.2 ± 2.3
	Estimated	6.8 ± 1.5	-	7.4 ± 1.5
	p-value*	0.183†	-	0.140†
1-5	Measured	13.9 ± 3.2	13.9 ± 3.2	13.9 ± 3.2
	Estimated	13.7 ± 2.6	15.5 ± 3.9	15.7 ± 2.8
	p-value*	0.200†	< 0.001	< 0.001
6-10	Measured	28.8 ± 8.8	28.8 ± 8.8	42.1 ± 13.5
	Estimated	31.8 ± 4.5	31.8 ± 4.5	44.9 ± 3.0
	p-value*	< 0.001	< 0.001	< 0.001
	Measured	35.8 ± 10.6	42.1 ± 13.5	42.1 ±

				13.5
11-14	Estimated	42.9 ± 1.7	44.9 ± 3.0	50.5 ± 4.0
	p-value*	< 0.001	< 0.001	< 0.001
Sex				



	Measured	19.1 ± 10.4	22.9 ± 11.4	20.5 ± 12
Male	Estimated	20.9 ± 12	25.8 ± 11.6	24.2 ± 14.2
	p-value*	< 0.001	< 0.001	< 0.001
	Measured	20.7 ± 12.5	25.1 ± 15.7	23.7 ± 15.8
Female	Estimated	22.3 ± 12.3	26.9 ± 12.8	27.0 ± 15.3
	p-value*	< 0.001	< 0.001	< 0.001
Ethnicity				
	Measured	19.9 ± 11.9	24.1 ± 14.1	22 ± 14.4
White	Estimated	21.5 ± 12.1	26.5 ± 12.2	25.5 ± 14.9
	p-value*	< 0.001	< 0.001	< 0.001
	Measured	19.7 ± 11.2	23.9 ± 13.4	21.7 ± 13.7
Non-white	Estimated	21.4 ± 12.3	26.3 ± 12.1	25.2 ± 14.7
	p-value*	< 0.001	< 0.001	< 0.001

Weight is expressed in kg, as mean ± SD. *p-value obtained using the t-test for paired samples. †A p-value greater than the significance level indicates that measurements tend to be the same with both methods.

Table III. Agreement between measured and estimated weight according to method and respective criteria in paediatric well-nourished patients admitted to a clinical ward. IMIP, August 2106 - January 2017

		0-12 months	1-5 years	6-12 years
APLS Update	N	4 2	176	135
	Bias	0.104	0.093	-5.162
	SD	1.04	1.8 5	5.11
	95% CI*	-1.9 to 2.1	-3.5 3.7	to -15.2 to 4.5
	CCC†	0.80	0.7 9	0.57
	Lower limit‡	0.69	0.7 5	0.48
	<hr/>			
1-14 years				
Luscombe and Owens	N		337	
	Bias		-2.823	
	SD		4.94	
	95% CI*	-	12.5 6.8	to
	CCC†		0.89	
	Lower limit‡		0.87	
<hr/>				
	N		328	
	Bias		-0.736	

Broselow tape	SD	2.35
	95% CI*	- 5.3 to 3.87
	CCC†	0.96
	Lower limit‡	0.94

Best Guess	1-11 months	1-4 years	5-14 years
-------------------	------------------------	----------------------	-----------------------



N	40	161	176
Bias	-0.424	-1.892	-6.515
SD	0.99	1.69	6.406
95% CI*	-2.3 to 1.5	-5.2 to 1.4	-19.1 to 6.0
CCC†	0.78	0.58	0.74
Lower limit***	0.65	0.51	0.65

*95% confidence interval. †CCC: Lin's concordance correlation coefficient (14). ‡Lower limit of the 95% confidence interval of the reliability of Lin's concordance correlation coefficient: < 0.90 = poor; 0.90-0.95 = moderate; 0.95-0.99 = good; > 0.99 = excellent (15).

Nutrición
Hospitalaria




Fig. 1. Bland-Altman plot showing the difference between measured and estimated weight as calculated by the APLS Update method in well-nourished patients. a) $\text{Weight} = (0.5 \times \text{age in months}) + 4$; b) $\text{weight} = (2 \times \text{age in years}) + 8$; c) $\text{weight} = (3 \times \text{age in years}) + 7$. The continuous line indicates the bias and the dotted lines indicate the limits of concordance. IMIP, August 2016 to January 2017.




Fig. 2. Bland-Altman plot showing the difference between measured and estimated weight as calculated by the Luscombe and Owens method [a) $\text{weight} = (3 \times \text{age in years}) + 7$] and by the Broselow tape in well-nourished patients. The continuous line indicates the bias and the dotted lines indicate the limits of concordance. IMIP, August 2016 to January 2017.




Fig. 3. Bland-Altman plot showing the difference between measured and estimated weight as calculated by the Best Guess method in well-nourished patients. a) $\text{Weight} = (\text{age in months} + 9)/2$; b) $\text{weight} = 2 (\text{age in years} + 5)$; c) $\text{weight} = 4 \times \text{age in years}$. The continuous line indicates the bias and the dotted lines indicate the limits of concordance. IMIP, August 2016 to January 2017.