



Trabajo Original

Pediatría

Stunting, wasting, and mid upper arm circumference status among children admitted to Nemazee Teaching Hospital

Desnutrición aguda y crónica, y circunferencia media del brazo en niños ingresados en el hospital universitario de Nemazee (Irán)

Seyed Mohsen Dehghani¹, Hazhir Javaherizadeh², Masoumeh Heydari¹, Naser Honar¹, Maryam Ataollahi¹, Homa Ilkhanipour³ and Hossein Moravej³

¹Gastroenterhepatology Research Center. Nemazee Teaching Hospital. Shiraz University of Medical Sciences. Shiraz, Iran. ²Department of Pediatric Gastroenterology. Abuzar Children's Hospital. Ahvaz Jundishapur University of Medical Sciences. Ahvaz, Iran. ³Department of Pediatric Endocrinology. Nemazee Teaching Hospital. Shiraz University of Medical Sciences. Shiraz, Iran

Abstract

Introduction and aim: The aim of this study was to evaluate nutritional status in children without prior hospital admission or evidence of chronic disease.

Subjects and methods: The current study is a cross-sectional and observational study which was conducted for assessing the nutritional status of children. In this study, consecutive sampling was used, with a sample size about 400 children aged 6 months to 18 years at first hospital admission. All subjects were hospitalized consecutively in the Pediatric Emergency Department of the Nemazee Teaching Hospital of Shiraz University of Medical Sciences (Shiraz, Islamic Republic of Iran). All children with evidence or history of chronic diseases such as liver cirrhosis, microcephaly, macrocephaly due to congenital infection or abnormal skull shape, patients with muscular or neurological problem, with prolonged steroid usage, nephrotic syndrome, cardiac problem, and protein losing enteropathy were excluded from the study. The duration of the study was six months, starting from January 2016.

Results: In the current study, 430 (225 boys and 175 girls) cases were included. Mean age was 5.39 (SEM 0.23) for boys and 5.78 (SEM 0.38) for girls. According to H/A criteria (current height/ideal height), 194 (48.5%) patients were stunted, 75 (18.8%) were mild, 49 (12.3%) were moderate and 70 (17.5%) were severe. According to W/A data (current weight/ideal weight), 188 (47%) of the patients studied were malnourished. Of these cases, 118 (29.5%) had mild malnutrition; 54 (13.5%), moderate; and 16 (4%), severe. According to body mass index (BMI), 40 (10%) patients were overweight, four (16%) were obese, and four (1%) were severe obese. In relation to middle upper arm circumference (MUAC), 56 (14%) patients were malnourished, 33 (8.3%) were at risk, 14 (3.5%) presented moderate malnutrition, and 9 (2.3%), severe malnutrition.

Conclusion: Prevalence of malnutrition among children without prior admission was 48.5% (according to H/A). According to W/A, 47% of children had different degrees of malnutrition.

Key words:

Malnutrition. Stunting. Wasting disease.

Resumen

Introducción y objetivo: el propósito de este estudio fue evaluar el estado nutricional de los niños ingresados por primera vez en el hospital y sin historia de enfermedad crónica.

Pacientes y métodos: se trata de un estudio transversal, observacional para valorar el estado nutricional al ingreso. La muestra estuvo constituida por algo más de 400 niños, de entre 6 meses y 18 años, ingresados por primera vez en el hospital. Se trataba de pacientes ingresados de forma consecutiva desde el servicio de Urgencias del hospital Nemazee de la Universidad Shiraz, en la República Islámica de Irán. Se excluyeron los pacientes con historia de enfermedad crónica como cirrosis hepática, microcefalia, macrocefalia por infección congénita o forma craneal anómala, pacientes con enfermedades musculares o neurológicas, con uso prolongado de corticoides, síndrome nefrótico, cardiopatía o enteropatía pierde-proteínas. La duración del estudio fue de seis meses, a partir de enero de 2016.

Resultados: se incluyeron 430 niños (225 varones y 175 niñas). La edad media fue de 5,39 años (SEM 0,23) para los varones y de 5,78 (SEM 0,38) en las niñas. De acuerdo al peso para la talla ideal, 194 (48,5%) tenían desnutrición crónica, 75 (18,8%) leve, 49 (12,3%) moderada y en 70 (17,5%) era grave. Según el porcentaje del peso ideal, 188 pacientes (47%) presentaban desnutrición aguda, de los cuales en 118 (29,5%) era leve, en 54 (13,5%) moderada y en 16 (4%), grave. Según el índice de masa corporal (IMC), 40 (10%) tenían sobrepeso, 16 (4%) eran obesos y cuatro (1%) presentaban obesidad mórbida. Según el perímetro del brazo (MUAC), 56 (14%) tenían desnutrición, 33 (8,3%) estaban en riesgo de desnutrición, 14 (3,5%) presentaban desnutrición moderada y 9 (2,3%) intensa.

Conclusión: la prevalencia de desnutrición en los niños ingresado por vez primera fue del 48,5%. De acuerdo al peso para la edad en 47% de los niños había algún grado de desnutrición.

Palabras clave:

Desnutrición. Enlentecimiento. Enfermedad debilitante.

Received: 22/03/2017 • Accepted: 25/06/2017

Dehghani SM, Javaherizadeh H, Heydari M, Honar N, Ataollahi M, Ilkhanipour H, Moravej H. Stunting, wasting, and mid upper arm circumference status among children admitted to Nemazee Teaching Hospital. *Nutr Hosp* 2018;35:33-37

DOI: <http://dx.doi.org/10.20960/nh.1143>

Correspondence:

Hazhir Javaherizadeh. Department of Pediatric Gastroenterology. Abuzar Children's Hospital. Ahvaz Jundishapur University of Medical Sciences. Ahvaz, Iran
e-mail: Hazhirja@yahoo.com; Hazhirja@ajums.ac.ir

INTRODUCTION

Childhood malnutrition is a major public health problem among children in developing countries. It can affect physical and intellectual growth and is also considered as the main cause of child morbidity and mortality.

High degrees of malnutrition have been reported in our country (1). The rate of stunting, underweight, and wasting were 9.53, 9.66 and 8.19%, respectively (2). In another study from Spain, the rate of malnutrition at the moment of hospitalization was 8.2 (3). In a Brazilian study, 16.3% of children under five years of age presented malnutrition at admission (4). Most of this studies were carried out in apparently healthy school children. There are little published studies about nutritional assessment at hospital admission. The aim of this study is to evaluate the frequency of obesity and malnutrition among hospitalized children with no past medical history at the moment of hospital admission.

METHODS

This is a prospective, cross-sectional and observational study which was conducted for the assessment of children nutritional status. The study sample included 430 patients: 225 boys (56%) and 175 girls (44%), with a mean age of 5.39 (SEM 0.32) years for boys and 5.78 (SEM 0.38) years for girls. All subjects were hospitalized consecutively in the Pediatric Emergency Department of the Nemazee Teaching Hospital of the Shiraz University of Medical Sciences (Shiraz, Islamic Republic of Iran).

All children with evidence or history of chronic diseases such as liver cirrhosis, microcephaly, macrocephaly due to congenital infection or abnormal skull shape, and patients with muscular or neurological problems, with prolonged steroid usage, nephrotic syndrome, cardiac problems, and protein losing enteropathy were excluded from the study. The duration of the study was six months starting from January 2016.

Parents or caregivers were informed about the aims of the study and verbal consent was obtained to take their children anthropometric measurements.

The instruments needed were a structured questionnaire and anthropometric measuring tools such as digital scale for weight with 0.1 kg accuracy and measuring tape with a 0.1 cm accuracy. The questionnaire included child age, sex, birth weight, height and head circumference (recorded from the child birth report card), which was completed by interviewing mothers or child caregivers.

Other required data were recorded through measuring height, weight, head circumference, and MUAC.

MEASURING WEIGHT

The child was weighed on a digital scale according to the standard way for the weight measurement. Evaluation of wasting was

done according to the following formula: $W/A: W/A: [\text{observed weight}/\text{median weight (same age and sex)}] \times 100$.

Classification of wasting was done as shown in table I.

MEASURING LENGTH

The height of children aged < 2 years was measured in supine position (top of the head to the bottom of the heel), and the rest of children were measured in standing position, with an accuracy of 0.10 cm.

Height measurement was taken on the flat floor and against a wall (flat surface). The measuring tape was placed 50 cm above the floor (and recorded height was gathered with that). The children stood with their feet flat, together, and against the wall. Legs were straight, arms were at sides, and shoulders were level. The child looked straight ahead, with the line of the sight parallel with the floor. Height was evaluated according to the following formula: $H/A: [\text{observed height}/\text{median height (same age and sex)}] \times 100$.

Stunting was classified as shown in table II and the classification of body mass index is shown in table III.

Table I. Classification of wasting according to weight-to-age ratio (Gómez classification)

Nutritional status	W/A
No malnutrition	More than 90
Mild malnutrition	75-90
Moderate malnutrition	60-74
Severe malnutrition	Less than 60

Table II. Classification of stunting according to H/A ratio (Waterlow classification)

Nutritional status	H/A
No malnutrition	More than 95
Mild malnutrition	90-95
Moderate malnutrition	85-89
Severe malnutrition	Less than 85

Table III. Classification of nutritional status according to BMI

BMI	
< 5	Underweight
5-85 th	Normal
85-95 th	Overweight
95-99 th	Obese
> 99 th	Severe obese

BMI: body mass index.

MEASURING MUAC

First, the child’s arms were relaxed and hanged down the sides of the body. The tip of the left shoulder (acromion) and the tip of the left elbow (olecranon) were located, and midpoint was marked. Then, the position of the tape was corrected for measuring arm circumference while ensuring that the tape neither pinched the arm nor was loose. Severity of malnutrition was calculated according to table IV.

MUAC has a high specificity and appears to be better predictor of childhood mortality than weight for height (5,6).

Gomez and Waterlow classifications were used for evaluation of weight for age and height for age ratio. World Health Organization (WHO) growth charts were used for weight, height, and BMI evaluation. Overweight, obesity, and severe obesity were defined according to BMI 85-95%, 95-99% and > 99%, respectively (Table III).

STATISTICAL ANALYSIS

Statistical analysis was performed with SPSS, version 16 (Chicago, IL, USA). Continuous variables were performed as mean and

standard error of mean (SEM), while categorical variables were presented as number and percentage. The Chi-squared test was used to compare differences in categorical variables and the independent t test for continuous variables between boys and girls. p value less than 0.05 was considered to be statistically significant.

RESULTS

The study sample included 430 patients: 225 boys (56%) and 175 girls (44%), with a mean age of 5.39 (SEM 0.32) for boys and 5.78 (SEM 0.38) for girls. Our patients were placed in four categories according to age: < 2 years (n = 124), 2-5 years (n = 120), 6-10 years (n = 78), and > 10 years (n = 78).

According to W/A data (current weight/ideal weight), 188 (47%) of the patients studied were malnourished: 118 (29.5%) presented mild malnutrition; 54 (13.5%), moderate; and 16 (4%), severe (Table V).

According to H/A data (current height/ideal height), 194 (48.5%) of the patients studied were stunted, 75 (18.8%) were mild, 49 (12.3%) were moderate, and 70 (17.5%) were severe (Table VI).

According to BMI, 40 (10%) patients from the sample were overweight, four (16%) were obese, and four (1%) were severe obese (Table VII). Regarding MUAC, 56 (14%) of the patients studied were malnourished, 33 (8.3%) were at risk, 14 (3.5%) presented moderate malnutrition, and nine (2.3%) were severely malnourished (Table VIII).

DISCUSSION AND CONCLUSION

In our study, we found that 47% of patients were underweight (low weight for age), 48.5% were stunted (low height for age),

Table IV. Classification of nutritional status according to MUAC

MUAC (mm)	Nutritional status
Less than 110	Severe acute malnutrition
110-125	Moderate acute malnutrition
125-135	At risk of malnutrition
More than 135	Well nourished

MUAC: mid upper arm circumference.

Table V. Malnutrition among children admitted to the hospital according to W/A ratio (Gómez classification)

Sex	No malnutrition	Mild malnutrition	Moderate malnutrition	Severe malnutrition
Male (n = 255)	126 (56.0%)	64 (28.4%)	27 (12.0%)	8 (3.6%)
Female (n = 175)	86 (49.1%)	54 (30.9%)	27 (15.4%)	8 (4.6%)

Table VI. Nutritional status according to height-for-age criteria (Waterlow classification)

Sex	Normal	Mild	Moderate	Severe
Male (n = 255)	110 (48.9%)	49 (21.8%)	24 (10.7%)	42 (18.6%)
Female (n = 175)	96 (54.9%)	26 (14.9%)	25 (14.3%)	28 (16.0%)

Table VII. BMI among children admitted to the hospital

Sex	Underweight	Normal	Overweight	Obese	Severe obesity
Male	8 (3.6%)	186 (82.7%)	19 (8.4%)	10 (4.4%)	2 (0.9%)
Female	12 (6.9%)	134 (76.6%)	21 (12.0%)	6 (3.4%)	2 (1.1%)

Table VIII. MUAC among children admitted to the hospital

At risk	33 (8.3%)
Normal	344 (86%)
Moderate	14 (3.5%)
Severe	9 (2.3%)

and 14% were malnourished according to MUAC. In the study by Moreno Vilares et al. (7), prevalence of malnutrition was 7.1% for moderate and 0.7% for severe malnutrition, which is significantly lower than that of the current study. In the study by Durá Travé et al., prevalence of malnutrition at the moment of hospitalization was 8.2%, which was lower than in our study (8). In their study, children with congenital malformations were included. In our study, children with chronic disease or abnormal skull shape were excluded. In the study by Lee and Ahmad (9), the prevalence rates of acute and chronic undernutrition were 11% and 14% respectively, that is, lower than in our study. In another study from Brazil, the prevalence of malnutrition was 10%, 18.5%, 21% and 14.7% according to weight for height, weight for age, height for age and BMI, respectively (10). In this study, children with Down syndrome and children with bone deformity were excluded (10). Consistent with our study, Mahdavi et al., in a study carried out in Tabriz (Iran) covering 140 patients aged 2-12 years, found chronic malnutrition in 30.7% of cases according to H/A, and acute malnutrition in 48.6% of cases according to W/A results (11). In a study conducted in Turkey including 528 patients with a mean age of 5.8 years, Dogan et al. found chronic malnutrition in 27% of cases according to H/A and acute malnutrition in 52.4% of cases according to W/A (12). In another study which was done by Veghari and Vakili, the rate of underweight and stunting was estimated to be 3.20% and 4.93%, respectively (13). A study done by Nouri Saelidou et al. on children under five years in Azerbaijan, Iran, indicated that the rate of underweight and stunting was 2.3% and 7.3%, respectively (14). In another study, prevalence of underweight among Iranian children according to CDC percentiles was 13.9% (15). In the study by Moeeny et al., malnutrition was present in 25.2% of inpatient children (16).

In another study from Turkey, according to the Gomez classification, malnutrition was present in 46.8% of the cases (17). In this study, patients with recurrent hospitalizations, neurological sequelae, and neonates were excluded; these exclusion criteria are similar to those of the current study. In the study by Ogunlesi et al., carried out in Nigeria, 18.3% of children had severe acute malnutrition according the Wellcome classification (18).

In the current study, 10% of the cases were overweight, 4% obese, and 1% were severe obese. In the study by Veghari and Vakili, prevalence of overweight was 3.3-5.2% between 1998 and 2013 (13). In a study from Iran, prevalence of overweight was 10.4% (19). In line with our study, a study was done in Azerbaijan by Nouri et al. where the prevalence of obesity and overweight was 1.3% and 5.1%, respectively (14). In a multicenter study from

Spain (20), overweight and obesity were seen in 37.9% of children, which is significantly higher than the percentage observed in the current study.

According to MUAC, 2.3% of the patients studied had severe malnutrition, 3.5% had moderate malnutrition and 8.3% were at risk of malnutrition. There are some differences between the results of these studies. These differences may be due to the different criteria which were used for the definition of malnutrition. In the current study, all children with a history of chronic disease or malformations were excluded. But in the study by Durá Travé et al., patients with congenital malformations were included (8).

CONCLUSION

In conclusion, our study demonstrates that the frequency of malnutrition among hospitalized children is relatively high. This higher prevalence of malnutrition needs more consideration because malnutrition has adverse effect on the outcome of the patients admitted in hospital.

The main limitation of the study was that it is a single center study.

FUNDING

This study was supported by research affairs of the Shiraz University of Medical Sciences.

ACKNOWLEDGMENTS

Data used in this paper was obtained from general physician thesis of Masoomeh Heidary (n= 93-01-01-8445), which was approved by research affairs of the Shiraz University of Medical Sciences.

REFERENCES

1. Karimi B, Ghorbani R, Niaki MA. Malnutrition and some related factors in primary school children, Semnan, Iran. *J Egypt Public Health Assoc* 2016;91:174-8.
2. Kavosi E, Hassanzadeh Rostami Z, Kavosi Z, et al. Prevalence and determinants of under-nutrition among children under six: A cross-sectional survey in Fars province, Iran. *Int J Health Policy Manag* 2014;3:71-6.
3. Dura-Trave T, San Martín-García I, Gallinas-Victoriano F, et al. Prevalence of malnutrition in hospitalised children: Retrospective study in a Spanish tertiary-level hospital. *JRSM Open* 2016;7:2054270416643889.
4. Saccardo Sarni RO, Suano de Souza F, Catherino P, et al. Treatment of severe malnourished children with WHO protocol: Experience of a referral center in Sao Paulo, Brazil. *Arch Latinoam Nutr* 2005;55:336-44.
5. Joseph B, Rebello A, Kullu P, et al. Prevalence of malnutrition in rural Karnataka, South India: A comparison of anthropometric indicators. *J Health Popul Nutr* 2002;20:239-44.
6. Briend A, Zimicki S. Validation of arm circumference as an indicator of risk of death in one to four year old children. *Nutr Res* 1986;6:249-61.
7. Moreno Villares JM, Varea Calderón V, Bousoño García C. Malnutrition in children admitted to hospital. Results of a national survey. *An Pediatr (Barc)* 2017;86:270-6.

8. Dura Trave T, San Martín García I, González Benavides A, et al. Nutritional status at the time of admission among patients admitted to a tertiary-care paediatric hospital. *Nutr Hosp* 2015;31:2465-71.
9. Lee WS, Ahmad Z. The prevalence of undernutrition upon hospitalization in children in a developing country: A single hospital study from Malaysia. *Pediatr Neonatol* 2017.
10. De Moraes Silveira CR, De Mello ED, Antonacci Carvalho PR. Evolution of nutritional status of pediatric patients of a tertiary care general hospital in Brazil. *Nutr Hosp* 2008;23:599-606.
11. Mahdavi AM, Ostadrahimi A, Safaiyan A. Nutritional status of children hospitalized in Tabriz Paediatric Hospital, Islamic Republic of Iran, 2008. *East Mediterr Health J* 2011;17:36-40.
12. Dogan Y, Erkan T, Yalvac S, et al. Nutritional status of patients hospitalized in pediatric clinic. *Turk J Gastroenterol* 2005;16:212-6.
13. Veghari G, Vakili M. Trend of stunting, overweight and obesity among children under five years in a rural area in Northern Iran, 1998-2013: Results of three cross-sectional studies. *Arch Iran Med* 2016;19:397-402.
14. Nouri Saeidlou S, Babaei F, Ayremlou P. Malnutrition, overweight, and obesity among urban and rural children in North of West Azerbaijan, Iran. *J Obes* 2014;2014:541213.
15. Kelishadi R, Ardalan G, Gheiratmand R, et al. Thinness, overweight and obesity in a national sample of Iranian children and adolescents: CASPIAN Study. *Child Care Health Dev* 2008;34:44-54.
16. Moeeni V, Walls T, Day AS. Assessment of nutritional status and nutritional risk in hospitalized Iranian children. *Acta Paediatr* 2012;101:e446-51.
17. Kapci N, Akcam M, Koca T, et al. The nutritional status of hospitalized children: Has this subject been overlooked? *Turk J Gastroenterol* 2015;26:351-5.
18. Ogunlesi TA, Ayeni VA, Fetuga BM, et al. Severe acute malnutrition in a population of hospitalized under-five Nigerian children. *Niger Postgrad Med J* 2015;22:15-20.
19. Jafari S, Fouladgar M, Naeeni MM, et al. Body mass index, weight-for-age, and stature-for-age indices in Iranian school children in relation to weight and growth disorders: A population-based survey. *Int J Prev Med* 2014;5:S133-8.
20. Moreno Villares JM, Varea Calderón V, Bousono García C, et al. Nutrition status on pediatric admissions in Spanish hospitals; DHOSPE study. *Nutr Hosp* 2013;28:709-18.