



Trabajo Original

Nutrición en el anciano

The association between obesity and vitamin D status among older adults in Ecuador: analysis of the SABE survey

La asociación entre obesidad y el estatus de vitamina D entre los adultos mayores en Ecuador: análisis de la encuesta SABE

Carlos H. Orces

Department of Medicine. Laredo Medical Center. Laredo, Texas

Abstract

Background: although it is well established that body fat mass is inversely associated with vitamin D (25(OH)D) concentrations, little is known whether obesity increases the risk of 25(OH)D insufficiency among older adults in Ecuador.

Methods: the present study used data from the National Survey of Health, Wellbeing, and Aging to describe the prevalence of obesity and 25(OH)D insufficiency (< 20 ng/ml) among Ecuadorians aged 60 years and older. Logistic regression models were used to examine the independent association between obesity and 25(OH)D insufficiency.

Results: a total of 2,270 participants with a mean age of 71.5 (SD 8.1) years comprised the sample size, representing an estimated 1.1 million older adults in Ecuador. Overall, the crude prevalence of obesity was 19.2% and 25(OH)D insufficiency was present in 14.0% of men and 34.6% of women. Moreover, the proportion of women with 25(OH)D insufficiency remained steady across BMI categories. In contrast, 25(OH)D insufficiency prevalence rates in men increased progressively as body mass index (BMI) categories also increase. Moreover, obese older men (OR 2.04; 95% CI: 1.99-2.09) were two times more likely to have 25(OH)D insufficiency compared with those defined as having an ideal weight, even after adjustment for potential confounders. In women, this association was attenuated. However, 25(OH)D insufficiency prevalence rates remained 12% higher in obese women (OR 1.12; 95% CI: 1.11-1.14) than their normal weight counterparts.

Conclusion: obesity is associated with increased risk of 25(OH)D insufficiency in Ecuador. Thus, obese older adults should be offered vitamin D supplementation and counseled regarding lifestyle modifications to improved their 25(OH)D status.

Key words:

Obesity. Older adults. Vitamin D. Ecuador.

Resumen

Antecedentes: aunque está bien establecido que la masa adiposa corporal esta inversamente asociada con las concentraciones de vitamina D (25(OH)D), no se ha descrito si la obesidad aumenta el riesgo de insuficiencia de 25(OH)D entre los adultos mayores en Ecuador.

Métodos: el presente estudio utiliza datos de la Encuesta Nacional de Salud, Bienestar y Envejecimiento (SABE) para describir la prevalencia de la obesidad y la insuficiencia de 25(OH)D (< 20 ng/ml) entre los ecuatorianos de 60 años de edad o mayores. Modelos de regresión logística se usaron para examinar la asociación independiente entre la obesidad y la insuficiencia de 25(OH)D.

Resultados: un total de 2.270 individuos con un promedio de edad de 71,5 (DE 8,1) años participaron en el estudio, el cual representa un estimado de 1,1 millones adultos mayores en Ecuador. La prevalencia de obesidad fue del 19,2%, y el 14,0% de los hombres y el 34,6% de las mujeres fueron clasificados con insuficiencia de 25(OH)D. En general, la prevalencia de insuficiencia de 25(OH)D entre las mujeres se mantuvo aumentada a través de las categorías de índice de masa corporal (IMC). En contraste, las tasas de prevalencia de insuficiencia de 25(OH)D en los hombres aumentaron progresivamente a medida que las categorías de IMC también aumentaron. Por otra parte, los hombres obesos mayores (OR 2,04; IC 95%: 1,99-2,09) fueron dos veces más propensos a tener insuficiencia de 25(OH)D en comparación con los que tenían un peso ideal, incluso después del ajuste por factores de confusión. En las mujeres, esta asociación fue atenuada. Sin embargo, las tasas de prevalencia de insuficiencia de 25(OH)D fueron un 12,0% mayores en mujeres obesas (OR 1,12, IC 95%: 1,11-1,14) que en sus contrapartes de peso normal.

Conclusión: la obesidad en los adultos mayores se asocia con un aumento en el riesgo de insuficiencia de 25 (OH)D en Ecuador. Por lo tanto, los adultos mayores con obesidad deberían recibir suplementos de vitamina D y consejos acerca de modificaciones del estilo de vida para mejorar el estatus de 25(OH)D.

Palabras clave:

Obesidad. Adultos mayores. Vitamina D. Ecuador.

Received: 06/01/2018 • Accepted: 04/03/2018

Orces CH. The association between obesity and vitamin D status among older adults in Ecuador: analysis of the SABE survey. Nutr Hosp 2018;35(5):1066-1071

DOI: <http://dx.doi.org/10.20960/nh.1752>

Correspondence:

Carlos H. Orces. Department of Medicine. Laredo Medical Center. 1700 East Saunders. 78041 Laredo, Texas
e-mail: corces07@yahoo.com

INTRODUCTION

Vitamin D deficiency leads to alterations in calcium and phosphorus homeostasis, resulting in secondary hyperparathyroidism with increased bone turnover, progressive bone loss, and increased risk of fractures (1,2). Numerous studies have demonstrated lower serum total 25-hydroxyvitamin D (25[OH]D) concentrations in obese subjects compared with their non-obese counterparts (3-10). Although the precise mechanism of decreased 25(OH)D levels in obesity has not been fully elucidated, research findings suggest that increased metabolic clearance and enhanced uptake of vitamin D by adipose tissue or decreased bioavailability of vitamin D once it is deposited in fat tissue may account for this finding (3,4). Moreover, Drincic et al. recently described that simple volumetric dilution may be the most parsimonious explanation for the low vitamin D status in obesity (5).

Aging is characterized by a reduction of muscle, bone mass, and strength levels with a concomitant increase of body fat mass, especially visceral fat mass (11). Moreover, older adults are particularly at higher risk of 25(OH)D deficiency because sunlight exposure is usually limited as a result of lifestyle changes. Likewise, the total production of previtamin D₃ after exposure to solar ultraviolet B radiation also decreases considerably with aging (12,13). In Ecuador, despite abundant sunlight throughout the year, a recent study among older adults reported a high prevalence of 25(OH)D insufficiency predominantly in women, indigenous, and residents in the Andes Mountains region of the country. In addition, subjects defined as having obesity had lower 25(OH)D concentrations compared with their non-obese counterparts (14). Because obesity is potentially a modifiable risk factor of 25(OH)D insufficiency, the present study aimed to extend previous research findings by examining the association between obesity and 25(OH)D status in a nationally representative sample of older adults in Ecuador.

METHODS

The present study was based on data from the National Survey of Health, Wellbeing, and Aging (Encuesta Nacional de Salud, Bienestar, y Envejecimiento; SABE II). Briefly, this survey is a probability sample of households with a least one person aged 60 years or older residing in the Andes Mountains and coastal regions of Ecuador. The SABE survey complex sampling design has been described elsewhere (15). Between April and August 2010, participants had biochemical evaluation to determine their nutritional status, including 25(OH)D concentrations. The survey data, including operation manuals, are publicly available and can be downloaded from the SABE survey website (16).

CHARACTERISTICS OF SUBJECTS

Age, sex, and race (white, black, mestizo, mulatto, or indigenous) were self-reported. Subjects were asked about their region (coastal vs mountains) and area of residence (urban vs rural). Lit-

eracy was defined by answering affirmatively to the question "Can you write and read a message". Smoking status was classified as current, former and never, and physical activity was evaluated by the question "Do you exercise regularly such as jogging and dancing, or have you performed rigorous work at least three times weekly for the past year?" Participants who responded affirmatively were considered to engage in vigorous physical activity. Self-reported general health was grouped as excellent to good or fair to poor.

Body height in centimeters and weight in kilograms were measured and the body mass index (BMI) was calculated (kg/cm^2). BMI was classified into three categories: under/normal ($< 25 \text{ kg}/\text{m}^2$), overweight ($25\text{-}29.9 \text{ kg}/\text{m}^2$), and obese ($\geq 30 \text{ kg}/\text{m}^2$) (17).

Subjects were defined as having diabetes according to the American Diabetes Association 2017 criteria if they reported a physician diagnosis of diabetes or had a fasting plasma glucose $\geq 126 \text{ mg}/\text{dl}$ (18). Serum 25(OH)D was measured by liquid chromatography at NetLab laboratory (Quito, Ecuador). The lowest limit of detection for the serum 25(OH)D assay was 4 ng/ml. A serum 25(OH)D $< 20 \text{ ng}/\text{ml}$ was the cut-off level to define subjects with 25(OH)D insufficiency as recommended by the Institute of Medicine (19).

STATISTICAL ANALYSIS

The descriptive characteristics of participants were stratified by BMI categories and examined using the ANOVA and Chi-square test for continuous and categorical variables, respectively. Data for men and women were analyzed separately since the prevalence of 25(OH)D insufficiency and obesity differed by gender among older adults in Ecuador (14,20). Subsequently, sex-specific multivariate logistic regression models were assembled in stages to evaluate the independent associations between BMI categories and the prevalence of 25(OH)D insufficiency. The first model was adjusted for age, race, and area of residency. A second model was further adjusted for literacy, smoking status, physical activity, self-reported health, and diagnosis of diabetes. Of 2,375 participants who completed the biochemical evaluation, 105 subjects were excluded from this analysis because of missing data on 25(OH)D concentrations ($n = 5$) and BMI ($n = 100$) measurements. All analyses used sample weights to account for nonresponse and the unequal probability of selection of the SABE survey and thus provide estimates representative of the older adult population in Ecuador. Statistical analyses were performed using SPSS, version 17 software (SPSS Inc., Chicago, IL).

RESULTS

A total of 2,270 participants with a mean age of 71.5 (SD 8.1) years comprised the sample size, representing an estimated 1.1 million older adults in Ecuador. Table 1 shows the demographic, behavioral, and health characteristics of participants. In general, women and self-reported race as mestizo accounted for 55.1%

Table I. Characteristics of participants in the SABE survey

Characteristics	Participants	Weighted % (SE)
<i>Age (years)</i>		
60-69	1,069	47.5 (1.3)
70-79	754	33.7 (1.3)
≥ 80	447	18.8 (1.0)
<i>Gender</i>		
Women	1,239	55.1 (1.3)
Men	1,031	44.9 (1.3)
<i>Race</i>		
Indigenous	198	10.2 (0.9)
Black	75	3.4 (0.5)
Mestizo	1,524	68.9 (1.3)
Mulatto	80	3.5 (0.5)
White	253	12.1 (0.9)
Other	44	1.9 (0.3)
<i>Area of residency</i>		
Urban Andes Mountains	663	30.2 (1.2)
Urban coast	827	36.5 (1.3)
Rural Andes Mountains	482	20.1 (1.0)
Rural coast	298	13.2 (0.9)
<i>BMI (kg/m²)</i>		
Normal	947	42.7 (1.3)
Overweight	890	38.1 (1.3)
Obesity	433	19.2 (1.1)
<i>Literacy</i>		
Yes	1,606	70.1 (1.2)
No	662	29.9 (1.2)
<i>Vigorous physical activity</i>		
Yes	739	34.1 (1.3)
No	1,530	65.9 (1.3)
<i>Smoking status</i>		
Current	239	11.1 (0.9)
Former	632	28.8 (1.2)
Never	1,349	60.0 (1.3)
<i>Self-reported health</i>		
Good to excellent	544	25.2 (1.1)
Fair to poor	1,721	74.8 (1.1)
<i>Diabetes</i>		
Yes	394	16.9 (1.0)
No	1,842	83.1 (1.0)
<i>Vitamin D insufficiency</i>		
Yes	569	25.4 (1.1)
No	1,701	74.6 (1.1)


and 68.9% of the participants, respectively. In addition, the crude prevalence of obesity was 19.2% and 25.4% of subjects were defined as having 25(OH)D insufficiency.

Table II shows the characteristics of participants stratified according to BMI categories. Overall, women, residents in the urban Andes Mountains, and subjects defined as having diabetes had higher obesity prevalence rates than those without. Moreover, obese subjects had significantly lower 25(OH)D concentrations compared with their non-obese counterparts. Likewise, obesity prevalence rates were higher among older adults with 25(OH)D insufficiency than in those without. As shown in figure 1, the prevalence of 25(OH)D insufficiency was considerably higher in women and remained steady across BMI categories. In contrast, 25(OH)D insufficiency prevalence rates in men increased progressively as BMI categories also increase.

As shown in table III, obese older men were two times more likely to have 25(OH)D insufficiency compared with those defined as having an ideal weight. Moreover, this strong association persisted even after adjusting for sociodemographic, behavioral, and health characteristics of the participants. In women, obesity was weakly associated with an inadequate 25(OH)D status. However, the prevalence of 25(OH)D insufficiency remained 12% higher in obese women compared with their normal weight counterparts.

DISCUSSION

The present findings indicate that obese older Ecuadorians had higher 25(OH)D insufficiency prevalence rates than their normal weight counterparts. This relationship was particularly evident in men. Indeed, older men with obesity had two-fold higher odds of having 25(OH)D insufficiency compared with those classified having a normal weight even after adjusting for sociodemographic characteristics previously associated with a high prevalence of 25(OH)D insufficiency nationwide (14). In contrast, obese women were not significantly associated with inadequate 25(OH)D status,

**Figure 1.**

Prevalence of 25(OH)D insufficiency according to BMI categories.

Table II. Characteristics of participants according to BMI categories

	Normal	Overweight	Obese	p-value
Age, mean (SD)	72.5 (8.4)	71.1 (7.8)	70.0 (7.3)	< 0.0001
<i>Gender, %</i>				< 0.0001
Men	52.2 (2.0)	37.7 (1.9)	10.1 (1.1)	
Women	35.0 (1.7)	38.4 (1.7)	26.5 (1.7)	
<i>Race, %</i>				< 0.0001
Indigenous	64.8 (4.2)	26.5 (3.9)	8.8 (2.0)	
Black	51.3 (7.0)	35.1 (6.5)	13.4 (4.1)	
Mestizo	39.2 (1.6)	39.5 (1.6)	21.2 (1.4)	
Mulatto	43.3 (7.1)	37.1 (6.6)	19.5 (4.8)	
White	41.3 (3.9)	43.2 (3.8)	15.6 (2.6)	
<i>Area of residency, %</i>				< 0.0001
Urban Andes Mountains	27.6 (2.2)	47.0 (2.5)	25.3 (2.2)	
Urban coast	42.3 (2.2)	35.2 (2.0)	22.5 (1.8)	
Rural Andes Mountains	61.2 (2.7)	20.5 (2.5)	22.5 (1.8)	
Rural coast	50.1 (3.7)	37.4 (3.5)	12.5 (2.8)	
<i>Literacy, %</i>				< 0.0001
Yes	37.8 (1.5)	42.7 (1.5)	19.4 (1.2)	
No	53.7 (2.5)	27.7 (2.1)	18.6 (2.2)	
<i>Vigorous physical activity, %</i>				0.767
Yes	44.0 (2.2)	37.4 (2.2)	18.5 (1.8)	
No	42.0 (1.6)	38.4 (1.5)	19.5 (1.3)	
<i>Smoking status, %</i>				< 0.0001
Current	59.4 (4.0)	29.0 (3.7)	11.6 (2.3)	
Former	44.5 (2.5)	40.4 (2.4)	15.2 (1.7)	
Never	38.7 (1.7)	38.7 (1.6)	22.5 (1.5)	
<i>Self-reported health, %</i>				0.132
Good to excellent	39.6 (2.5)	42.4 (2.5)	18.0 (1.9)	
Fair to poor	43.8 (1.6)	36.6 (1.5)	19.6 (1.3)	
<i>Diabetes, %</i>				< 0.0001
Yes	27.8 (2.9)	39.5 (3.0)	32.7 (3.1)	
No	45.2 (1.5)	38.1 (1.4)	16.7 (1.1)	
25(OH)D (ng/ml), mean (SD)	28.3 (18.4)	26.2 (10.1)	25.1 (10.9)	< 0.0001
<i>Vitamin D insufficiency, %</i>				0.013
Yes	40.0 (2.6)	35.5 (2.4)	24.5 (2.2)	
No	43.6 (1.6)	39.0 (1.5)	17.4 (1.2)	

which may have been partly explained by a high prevalence of 25(OH)D insufficiency across BMI categories, but particularly among those defined as having a normal weight. Indeed, up to 36.3% of women with normal weight had evidence of 25(OH)D insufficiency in Ecuador. Similarly, overweight men and women had 11% and 13% lower risk of having 25(OH)D insufficiency than those with an ideal weight, respectively. Of relevance, a recent

study described significantly higher obesity prevalence rates nationwide in women than in men (20). Thus, it is possible that the increased prevalence of 25(OH)D insufficiency in older women may be related to gender differences in body fatness, which has been documented by other investigators (21,22).

The present study results are consistent with those from the Mexican Health and Aging Study in which obese older adults were

Table III. Association between BMI categories and 25(OH)D insufficiency among older adults in Ecuador, SABE survey

	Normal	Overweight	Obese
Men			
	Reference	OR (95% CI)	OR (95% CI)
Model 1	1.00	0.84 (0.82-0.85)	1.83 (1.78-1.87)
Model 2	1.00	0.89 (0.87-0.91)	2.04 (1.99-2.09)
Women			
	Reference	OR (95% CI)	OR (95% CI)
Model 1	1.00	0.95 (0.93-0.96)	0.95 (0.93-0.96)
Model 2	1.00	0.87 (0.86-0.89)	1.12 (1.11-1.14)
Total			
	Reference	OR (95% CI)	OR (95% CI)
Model 1	1.00	0.93 (0.92-0.94)	1.23 (1.21-1.24)
Model 2	1.00	0.89 (0.88-0.90)	1.26 (1.24-1.27)

Model 1: adjusted for age, race, and place of residency. Model 2: Adjusted for model 1 and literacy, smoking status, physical activity, self-reported health, and diagnosis of diabetes.

1.7 times more likely to be associated with 25(OH)D concentrations in the lowest tertile (≤ 20.4 ng/ml) compared with their normal weight counterparts (22). Similarly, results of the Health, Aging and Body Composition Study demonstrated that older obese black and white were 1.5 and 1.3 times more likely to be defined as having 25(OH)D insufficiency, respectively (23). A previous study among participants of the Longitudinal Aging Study Amsterdam also reported that higher BMI, waist circumference, and skin folds were statistically significantly associated with lower 25(OH)D concentrations. However, the associations of total body fat with serum 25(OH)D levels were stronger than the associations of anthropometric measures (23). Furthermore, Young et al. demonstrated that among Hispanic and African-American, 25(OH)D levels were inversely associated with baseline BMI, and computed tomography derived measures of subcutaneous and visceral adipose tissue (25).

Notably, a recent study conducted to examine the effect of BMI categories on 25(OH)D concentrations and bone health demonstrated low free 25(OH)D levels in obesity, which were not due to differences in protein binding. Indeed, serum albumin and vitamin D binding proteins and genotype variation did not differ by BMI categories. Moreover, that particular study provided evidence that lower 25(OH)D concentrations in obesity were not due to more rapid metabolic clearance (9). Although 25(OH)D is fat soluble, and distributed into fat, muscle, liver, and serum, all of these compartments are increased in volume in obesity. Consequently, lower 25(OH)D concentrations likely reflect a volumetric dilution effect, and whole body stores of 25(OH)D may be adequate (26). In a recent study, Carelli et al. reported

that 25(OH)D concentrations measured by mass spectroscopy in omental and subcutaneous adipose tissue did not significantly differ in obese and normal weigh women. However, total body vitamin D stores were significantly greater in obese women than in their normal weight counterparts. Thus, these findings also support the hypotheses that the enlarged adipose mass in obese individuals serves as a reservoir for vitamin D and that the increased amount of vitamin D required to saturate this depot may predispose obese individuals to inadequate serum 25(OH)D (27). Of relevance, a large study conducted among healthy volunteers in the province of Alberta, Canada, demonstrated that the differences in serum 25(OH)D between normal, overweight, and obese subjects significantly differed by supplementation dose. For instance, supplementation with 600 IU per day would achieve a mean serum 25(OH)D levels of 33.2, 30.4, and 26.4 ng/ml in normal weight, overweight, and obese participants, respectively. Moreover, a mean serum 25(OH)D concentration of 40 ng/ml in normal, overweight, and obese subjects, was estimated to require 2,080 IU, 3,065 IU, and 5,473 IU per day, respectively (28). However, as previously described, overweight and obese subjects were less likely to use any dietary or vitamin supplements compared with their normal weight counterparts (29). Therefore, it is possible that the increased prevalence of 25(OH)D insufficiency among obese Ecuadorians may be partly explained by the low use of vitamin D supplements. However, this hypothesis may not be confirmed because no previous study has reported the prevalence of vitamin D supplement use among older adults in Ecuador. Of interest, results of a recent systematic review of randomized and nonrandomized control weight-loss trials reported that the effect of weight loss on circulating 25(OH)D levels, random assignment to weight loss compared with weight maintenance resulted in a small but significantly greater increase in serum 25(OH)D of 0.44 ng/ml. Similar results were found for nonrandomized trials (30). Thus, based on these findings, older obese subjects should be offered vitamin D supplementation and lifestyle modifications to improve their 25(OH)D status.

Several limitations should be mentioned while interpreting the present study results. First, the temporal relationship between obesity and 25(OH)D status may not be established due to the cross-sectional study design. Second, participants self-reported their demographic and certain behavioral characteristics, which may be a source of recall bias. Third, the dietary intake of and use of vitamin D supplements was not assessed in the survey. Fourth, participant's sunlight exposure was not examined. Fifth, because traditional anthropometric measures of adiposity such as BMI are weaker than direct measures of adiposity, it is possible that the association between obesity and 25(OH)D may be underestimated in the present study (25). Finally, the present study findings may be generalized to older adults residing in the coastal and Andes Mountains regions of the country.

In conclusion, obesity is associated with increased risk of 25(OH)D insufficiency among older adults in Ecuador. Thus, obese older subjects should be offered vitamin D supplementation and counseled regarding lifestyle modifications to improve their 25(OH)D status.

REFERENCES

- Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357:266-81.
- Lips P. Vitamin D deficiency and secondary hyperparathyroidism in the elderly: consequences for bone loss and fractures and therapeutic implications. *Endocr Rev* 2001;22:477-501.
- Liel Y, Ulmer E, Shary J, Hollis BW, Bell NH. Low circulating vitamin D in obesity. *Calcif Tissue Int* 1988;43:199-201.
- Wortsman J, Matsuoka LY, Chen TC, Lu Z, Holick MF. Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr* 2000;72:690-3.
- Drincic AT, Armas LA, Van Diest EE, Heaney RP. Volumetric dilution, rather than sequestration best explains the low vitamin D status of obesity. *Obesity (Silver Spring)* 2012;20:1444-8.
- Blum M, Dolnikowski G, Seyoum E, Harris SS, Booth SL, Peterson J, et al. Vitamin D(3) in fat tissue. *Endocrine* 2008;33:90-4.
- Samuel L, Borrell LN. The effect of body mass index on optimal vitamin D status in U.S. adults: the National Health and Nutrition Examination Survey 2001-2006. *Ann Epidemiol* 2013;23:409-14.
- Arunabh S, Pollack S, Yeh J, Aloia JF. Body fat content and 25-hydroxyvitamin D levels in healthy women. *J Clin Endocrinol Metab* 2003;88:157-61.
- Walsh JS, Evans AL, Bowles S, Naylor KE, Jones KS, Schoenmakers I, et al. Free 25-hydroxyvitamin D is low in obesity, but there are no adverse associations with bone health. *Am J Clin Nutr* 2016;103:1465-71.
- Shea MK, Houston DK, Tooze JA, Davis CC, Johnson MA, Hausman DB, et al. Health, Aging and Body Composition Study. Correlates and prevalence of insufficient 25-hydroxyvitamin D status in black and white older adults: the health, aging and body composition study. *J Am Geriatr Soc* 2011;59:1165-74.
- Jura M, Kozak LP. Obesity and related consequences to ageing. *Age (Dordr)* 2016;38:23. DOI: 10.1007/s11357-016-9884-3.
- MacLaughlin J, Holick MF. Aging decreases the capacity of human skin to produce vitamin D3. *J Clin Invest* 1985;76:1536-8.
- Van der Wielen RP, Löwik MR, Van den Berg H, De Groot LC, Haller J, Moreiras O, et al. Serum vitamin D concentrations among elderly people in Europe. *Lancet* 1995;346:207-10.
- Orces CH. Vitamin D status among older adults residing in the littoral and Andes Mountains in Ecuador. *Sci World J* 2015;2015:545297. DOI: 10.1155/2015/545297
- Guevara PE, Andrade FC. Socioeconomic and lifestyle factors associated with chronic conditions among older adults in Ecuador. *Rev Panam Salud Publica* 2015;38:226-32.
- Instituto Nacional de Estadística y Censos. Encuesta de Salud, Bienestar del Adulto Mayor. Quito, Ecuador. Accessed on Nov 23rd 2017. Available from: <http://www.ecuadorencifras.gob.ec/encuesta-de-salud-bienestar-del-adulto-mayor>
- Centers for Disease Control and Prevention. About adult BMI. Accessed on Nov 23rd 2017. Available from: https://www.cdc.gov/healthyweight/assessing/bmi/adult_bmi/index.html
- American Diabetes Association. 2. Classification and Diagnosis of Diabetes. *Diabetes Care* 2017;40:S11-S24.
- Institute of Medicine. Dietary Reference Intakes for Calcium and Vitamin D. Washington DC: National Academies Press; 2010.
- Orces CH, Montalvan M, Tettamanti D. Prevalence of abdominal obesity and its association with cardio metabolic risk factors among older adults in Ecuador. *Diabetes Metab Syndr* 2017;11:S727-33.
- Van Dam RM, Snijder MB, Dekker JM, Stehouwer CD, Bouter LM, Heine RJ, et al. Potentially modifiable determinants of vitamin D status in an older population in the Netherlands: the Hoorn Study. *Am J Clin Nutr* 2007;85:755-61.
- Rontoyanni VG, Ávila JC, Kaul S, Wong R, Veeranki SP. Association between obesity and serum 25(OH)D concentrations in older Mexican adults. *Nutrients* 2017;31:9. pii: E97. DOI: 10.3390/nu9020097.
- Shea MK, Houston DK, Tooze JA, Davis CC, Johnson MA, Hausman DB, et al. Health, Aging and Body Composition Study. Correlates and prevalence of insufficient 25-hydroxyvitamin D status in black and white older adults: the health, aging and body composition study. *J Am Geriatr Soc* 2011;59:1165-74.
- Snijder MB, Van Dam RM, Visser M, Deeg DJ, Dekker JM, Bouter LM, et al. Adiposity in relation to vitamin D status and parathyroid hormone levels: a population-based study in older men and women. *J Clin Endocrinol Metab* 2005;90:4119-23.
- Young KA, Engelman CD, Langefeld CD, Hairston KG, Haffner SM, Bryer-Ash M, et al. Association of plasma vitamin D levels with adiposity in Hispanic and African Americans. *J Clin Endocrinol Metab* 2009;94:3306-13.
- Walsh JS, Bowles S, Evans AL. Vitamin D in obesity. *Curr Opin Endocrinol Diabetes Obes* 2017;24:389-394.
- Carrelli A, Bucovsky M, Horst R, Cremers S, Zhang C, Bessler M, et al. Vitamin D storage in adipose tissue of obese and normal weight women. *J Bone Miner Res* 2017;32:237-42.
- Ekwaru JP, Zwicker JD, Holick MF, Giovannucci E, Veugelers PJ. The importance of body weight for the dose response relationship of oral vitamin D supplementation and serum 25-hydroxyvitamin D in healthy volunteers. *PLoS One* 2014;9:e111265.
- Radimer K, Bindewald B, Hughes J, Ervin B, Swanson C, Picciano MF. Dietary supplement use by US adults: data from the National Health and Nutrition Examination Survey, 1999-2000. *Am J Epidemiol* 2004;160:339-49.
- Mallard SR, Howe AS, Houghton LA. Vitamin D status and weight loss: a systematic review and meta-analysis of randomized and nonrandomized controlled weight-loss trials. *Am J Clin Nutr* 2016;104:1151-9.