



Trabajo Original

Epidemiología y dietética

Meat and meat products intake in pregnancy and risk of small for gestational age infants. A case-control study

Consumo de carne y productos cárnicos durante el embarazo y riesgo de tener un recién nacido pequeño para su edad gestacional. Un estudio de casos y controles

Naomi Cano-Ibáñez^{1,2,3}, Juan Miguel Martínez-Galiano^{2,4}, Carmen Amezcua-Prieto^{1,2,3}, Rocio Olmedo-Requena^{1,2,3}, Aurora Bueno-Cavanillas^{1,2,3} and Miguel Delgado-Rodríguez^{2,4}

¹Department of Preventive Medicine and Public Health. Universidad de Granada. Granada, Spain. ²CIBER of Epidemiology and Public Health (CIBERESP). Carlos III Institute of Health. Madrid, Spain. ³Granada Biosanitary Research Institute (IBS-Granada). Granada, Spain. ⁴Department of Health Sciences. Universidad de Jaén. Jaén, Spain

Abstract

Introduction: different diets during pregnancy might have an impact on the health as reflected by the birth weight of the newborn. The consumption of meat and meat products during pregnancy and its relationship with the newborn health status have been studied by several authors. The studies carried out show inconsistent results

Objective: to analyse the association between maternal dietary intake of meat and meat products and the risk of small for gestational age (SGA) newborn.

Methods: a matched case-control study of 518 cases and controls of pregnant women was performed in Spain. Cases were women with a SGA newborn. Data about demographic characteristics and diet were collected. Meat consumption was gathered through a validated food frequency questionnaire. Meat and meat products intakes were categorized in quintiles (Q1-Q5). The association between maternal meat and meat product intakes and SGA was assessed by logistic regression models with adjustment for confounding factors.

Results: an intake of meat products above 6.8 g/day was associated with a lower risk of SGA delivery (OR = 0.7; 95% CI, 0.53-0.93) after adjusting for smoking, body mass index, previous preterm-low birth weight, newborn gender and adherence to Mediterranean diet.

Conclusions: meat consumption was not associated with SGA, whereas meat products showed a moderate protective relationship.

Key words:

Small for gestational age. Maternal nutrition. Meat. Meat products. Pregnancy.

Resumen

Introducción: la ingesta dietética durante el embarazo es un factor clave en la salud maternal y fetal, pudiendo tener un impacto en el peso al nacer del recién nacido. El consumo de carne y productos cárnicos durante el embarazo y su asociación con el estado de salud del neonato ha sido estudiado, sin embargo los hallazgos encontrados muestran resultados contradictorios.

Objetivo: analizar la asociación entre la ingesta dietética materna de carne y productos cárnicos durante el embarazo y el riesgo de tener un bebé pequeño para la edad gestacional (PEG).

Metodología: se realizó un estudio de casos y controles emparejados en mujeres españolas embarazadas (518 casos y 518 controles). Los casos fueron mujeres con un recién nacido PEG. El consumo de carne se obtuvo a través de un cuestionario de frecuencia de alimentos validado. La carne y la ingesta de productos cárnicos fueron categorizados en quintiles (Q1-Q5). La asociación entre la ingesta materna cárnica durante el embarazo y el riesgo de tener un recién nacido PEG se evaluó mediante modelos de regresión logística ajustados por factores confusores.

Resultados: una ingesta de productos cárnicos $\geq 6,8$ g/día se asoció significativamente con un menor riesgo de tener un recién nacido PEG (OR = 0,7; IC 95%, 0,53-0,93) después de ajustar el modelo por factores confusores: fumar, índice de masa corporal, embarazos previos con bajo peso al nacer, sexo del recién nacido y adherencia materna a la dieta mediterránea.

Conclusiones: el consumo materno de carne no se asoció con recién nacidos PEG, sin embargo el consumo de productos cárnicos mostró una moderada asociación protectora. Estos hallazgos respaldan el consejo de una dieta variada para mujeres embarazadas, que proporciona ingesta de proteínas y otros micronutrientes de diferentes fuentes alimentarias.

Palabras clave:

Pequeño para su edad gestacional. Nutrición materna. Carne. Productos cárnicos. Embarazo.

Received: 25/10/2018 • Accepted: 14/11/2018

Funding: This work was supported by a grant from the National Institute of Health Carlos III (PI11/02199). N.C.-I. would like to acknowledge the support by the Ministry of Education of Spain (FPU14/03630).

Cano-Ibáñez N, Martínez-Galiano JM, Amezcua-Prieto C, Olmedo-Requena R, Bueno-Cavanillas A, Delgado-Rodríguez M. Meat and meat products intake in pregnancy and risk of small for gestational age infants. A case-control study. *Nutr Hosp* 2019;36(2):405-411

DOI: <http://dx.doi.org/10.20960/nh.2366>

Correspondence:

Juan Miguel Martínez-Galiano. Universidad de Jaén. Campus de Las Lagunillas, s/n. Edificio B3. 23071 Jaén, Spain
e-mail: juanmimartinezg@hotmail.com

INTRODUCTION

Small for gestational age (SGA) is defined as a birth weight below the tenth percentile birth weight for an infant of a specific gestational age and sex (1). SGA has associated with increased infant and child morbidity and mortality (2,3). A growing body of evidence has linked SGA to the risk of developing chronic diseases such as metabolic disease in adulthood (4). SGA prevalence changes depending on birth weight standards and population (5). Besides, although the prevalence of SGA infants is high among the low and middle-income countries (6), the SGA is also increasing in developed countries such as Europe, reaching figures around 10.5% (7). Risk factors for SGA include maternal age, body mass index (BMI), chronic diseases, socioeconomic status and unhealthy lifestyles (8). Only a number of these factors can be modified. Among the modifiable risk factors, maternal nutrition plays a crucial role, influencing fetal growth and birth outcomes (9).

Nutritional requirements for many nutrients increase during pregnancy in order to support fetal growth (10). This may produce several maternal deficiencies of essential nutrients such as protein, vitamin B12, iron, iodine, folate and thiamine (11,12). An inadequate intake of all of these nutrients has been linked to maternal morbidities, neurodevelopmental disease in babies and a higher risk of SGA in the newborn (13,16). In fact, inadequate intake of protein-energy is related to fetal growth failure (17). A balanced energy protein supplementation has been associated with a risk reduction of SGA around 32% and increasing birth weight by 73 g (18). Furthermore, anemia due to low nutritional iron intake is associated with the birth of small babies, affecting an average of 56% of pregnant women in developing countries and above 18% in developed countries (19). As a result, recommendations on supplementation or fortification food during pregnancy for these nutrients have been formulated. For this reason, pregnancy dietary counselling recommend the supplementation of these nutrients (20).

Among the dietary sources, meat and meat products provide the recommended daily amounts of these nutrients, especially of protein and heme-iron (21). Thus, meat intake should be related to a lower frequency of some nutritional deficiencies, such as iron, protein, iodine vitamin B12 and other vitamins, and by this pathway could be possible to reduce the risk of low birth weight (LBW) and SGA newborn (22).

A healthy diet during pregnancy should include moderate amounts of meat and meat products (23). Nowadays, maternal dietary advice is based on the reduction of the consumption of this food group in order to reduce the risk of exposure to potentially harmful food pathogens, such as toxoplasmosis, found in raw or undercooked meat (24). However, the occurrence of this infection among pregnant women in developed countries is unusual (25).

Few studies have analyzed the role of meat and meat products on the risk of SGA newborn, showing inconsistent results. Probably by the differences in nutritional status of the study population and the methods used to evaluate the dietary intake. Consequently, some of them have shown risk associations on SGA

delivery (26,27) meanwhile other authors have showed protective associations (28,29).

The aim of this study is to analyze the association between maternal dietary intake of meat and meat products during pregnancy and the risk of having a SGA newborn in Spanish population.

MATERIALS AND METHODS

The study population includes women attended to at five hospitals of Eastern Andalusia (Spain): Hospital Universitario de Jaén (HUJ), Hospital de Úbeda (UB), hospitales universitarios de Granada (two centers) (HUG) and Hospital de Poniente (HP), serving 1.8 million people. Case and control groups were collected from May 15th, 2012, through July 15th, 2015. The Ethics Committees of the hospitals authorized this study. Informed consent was sought from every eligible woman.

The appropriate sample size was estimated based on the results of a similar study (27). To detect a significant ($p < 0.05$) OR of 0.6 between extreme quintiles with a statistical power of 80%, it was estimated that 447 pairs of cases and controls were required.

CASES

SGA was defined as having a birth weight smaller than the tenth percentile for the infant's gestational age when compared with that expected for the same gestational age and sex, according to tables previously developed for the Spanish population (30), without congenital malformations during the study period and residence in the referral area of the hospital. Nineteen women rejected participation. A total of 533 cases were selected: 79 (HUJ), 369 (HUG), 46 (UB) and 39 (HP).

CONTROLS

A match pair by age at delivery (± 2 years) was selected within the next week after including a case at the same hospital. Eligible women were those having a non SGA newborn with the same inclusion criteria for cases (residence in the referral area of the hospital and no malformations). Sixty-five women declined participation.

DATA COLLECTION

Information was obtained in both groups on general *socio-demographic habits and personal characteristics*, including age at pregnancy, ethnicity, education level, marital status, monthly income, socioeconomic class, occupation, adherence to Mediterranean diet (MedDiet adherence) and body mass index (BMI). BMI was calculated as weight in kilograms/height in m². Weight

and height were obtained from medical records of women if possible, or self-reported if not. Social class was coded in five levels ranging from I (the highest) to V (the lowest level) according to the classification of the Spanish Society of Epidemiology (31), which is near to that of the other authors reported in the Black report (32). To measure the MedDiet adherence, the index proposed by Trichopoulou (33) was used, which includes the following components: intake of vegetables, legumes, fruits, cereals, fish, meat and meat products, milk and milk products, with scores ranging from 0 to 9, with higher scores indicating greater adherence. *Obstetric history* comprised parity, antecedents of abortions, previous adverse perinatal outcomes and morbidities during pregnancy (preeclampsia, diabetes, infections and other obstetric conditions). *Birth weight* was measured as weight in grams in the delivery room. *Toxic habits* (smoking during pregnancy and alcohol consumption during and before pregnancy) were assessed with a structured questionnaire, which included the number and type of drinks during a week, specifying the day at the week and holidays. *Drugs and medication* including prescribed medications and over-the-counter drugs. Finally, the prenatal care (number of visits and date of first visit) was also considered. Prenatal care was measured by using the Kessner index. This index takes into account timing of entry in prenatal care, number of prenatal visits and gestational age at delivery (34).

DIETARY ASSESSMENT

Information on dietary habits in both groups was collected. Trained midwives gave a food frequency questionnaire (FFQ) to women after birth, previously validated in Spain, with open questions about the use of dietary supplements (35,26). All participants were asked to complete and return the questionnaire, being a criterion for the participation of the study. FFQ included a 137-item and allowed the classification into eleven food groups (dairy products, meats and meat products, fish and seafood, vegetables, fruits, cereals, potatoes, fats, nuts, precooked food, sauces and beverages). The questionnaire included nine response options (never or almost never, 1-3 times a month, once a week, 2-4 times a week, 5-6 times a week, once a day, 2-3 times a week day, 4-6 times a day and more than six times a day). For each food item, we estimated the average amount of food consumed (grams) multiplied by the intake frequency, the average total energy intake and the average intake of macro and micronutrients. A dietitian updated the nutrient data using the information recorded in the Spanish tables of food composition (37,38).

Meat and meat products included in the FFQ were: *meat group*, including chicken with/without skin, beef, pork, lamb and rabbit; and *meat products group*, including liver (beef, pork and chicken), other entrails (brain, heart and sweetbread), cured ham, cooked ham, processed meats (sausages, black pudding, etc.), pâté, hamburger and bacon.

Meat and meat products sizes were pointed out in the FFQ and the midwives gave examples of portion sizes according to groups predefined as follows: meat group (100-150 g), liver and

other entrails (100-150 g), ham cured/cooked (one slice or 30 g), processed meats, hamburger and bacon (50 g) and pâté (25 g).

After computing total energy intake, a total of 15 matched pairs were excluded due to an unreliable dietary assessment (total energy intake above 4,000 kcal/day), leaving 518 pairs for analysis.

STATISTICAL ANALYSIS

All analyses were performed using Stata (14.0, StataCorp LP, Tx. USA). Qualitative variables were analyzed through frequency distribution, whereas quantitative variables were expressed as means and standard deviation (SD). Pearson χ^2 test and Student's t test were used to assess differences in the characteristics of the participants. All p values are two tailed. Statistical significance was set at $p < 0.05$. The intake frequencies were derived from the FFQ, thus, each respondent indicated intakes for the different meat groups as: never or more than once serving per week.

Meat and meat products intakes were adjusted for total energy intake using the residuals method as recommended by Willet et al. (39). Food intake was stratified into quintiles, according to intakes observed in the control group (used as criterion of general population intake). This categorization was applied to intakes reported in the SGA group.

Conditional logistic regression models were performed to calculate crude odds ratios (cORs) and adjusted odds ratios (aORs) at 95% confidence interval (CI). The lowest quintile (Q1) was taken as the reference. Intermediate variables were discarded and to control for confounding variables previously analyzed in a priori approach related to maternal diet. The models were adjusted for energy intake, smoking, previous preterm-low birth weight, BMI, newborn's gender and adherence to MedDiet.

RESULTS

One thousand and thirty-six women participated in this study. Table I shows characteristics among SGA and control participants. Women in the SGA group were more likely to be smokers and have previous preterm or low-birth weight newborn, preeclampsia and intrauterine growth retardation ($p < 0.001$). In contrast, more women in the control group were married ($p < 0.036$) and had a higher mean gestational weight gain and BMI prior to gestation ($p < 0.001$). No significant differences were observed with regard to education level, Kessner index and alcohol intake.

The frequency of intake of different types of meat and meat products and its effect on SGA are shown in table II. No significant association was found between the risk of SGA and the intake of most of them. The intake of chicken without skin more than twice a week and a frequent intake of cooked ham (once a day) yielded a lower risk of SGA in crude models (cOR = 0.65; 95% CI, 0.44-0.98; cOR = 0.48; 95% CI, 0.26-0.87, respectively), but both associations disappear in adjusted models.

The relationships between total mean intake of meat and meat products, measured in quintiles (g/day), are presented in table III.

Table I. Description of the study population characteristics in the study (n = 1,036)

	Cases (SGA)		Controls (AGA)		p value
	518		518		
<i>Marital status, n (%)</i>					
Single	37	(7.1)	42	(8.1)	
Stable couple	161	(31.1)	124	(23.9)	
Married	320	(61.8)	352	(68.0)	
<i>Education level, n (%)</i>					
Primary	112	(21.6)	93	(17.9)	
High school, not ended	42	(8.1)	28	(5.4)	
High school	185	(35.7)	190	(36.7)	
University	179	(34.6)	207	(40.0)	
Previous preterm/low-birth weight newborn, n (%)	64	(12.4)	26	(5.0)	< 0.001
<i>Kessner index (prenatal care), n (%)</i>					
Adequate	259	(50.0)	253	(48.8)	
Intermediate	185	(35.7)	182	(35.2)	
Inadequate	74	(14.3)	83	(16.0)	
Smoking during pregnancy, n (%)	149	(28.8)	80	(15.4)	< 0.001
Preeclampsia, n (%)	46	(8.9)	11	(2.1)	< 0.001
Intrauterine growth retardation, n (%)	141	(27.2)	8	(1.5)	< 0.001
Weight gain during pregnancy (g/week), mean (SD)	278	(121)	310	(114)	< 0.001
Prepregnancy body mass index, mean (SD)	23.1	(4.5)	23.9	(4.1)	< 0.001
Alcohol intake (g/week), mean (SD)	4.2	(18.5)	3.1	(15.2)	0.312

SGA: small for gestational age; AGA: adequate for gestational age; SD: standard deviation; BMI: body mass index. Pearson Chi-square test and Student's t test were performed for categorical and continuous variables respectively.

No significant association was detected with total mean intake and no trend was observed, either in crude results or in adjusted models. Regarding the intake of meat products, no association was observed in raw results. In adjusted results, a significant OR was observed for an isolated quintile (Q4 vs Q1). Given that the OR figures for quintiles Q3, Q4, and Q5 were quite similar, Q3-Q5 vs Q1-Q2 were compared, that is an intake above 6.8 g/d vs a lower one, and the resulting OR achieved statistical significance (aOR = 0.70; 95% CI, 0.53-0.93, p = 0.015).

DISCUSSION

In the present study, the association between the maternal intake of meat and meat products during pregnancy and the risk of having a SGA baby in a Spanish population was investigated. No significant association for total meat intake was found, although an intake above 6.8 g/day of meat products reduced the risk of having a SGA newborn.

Current pregnancy dietary guidelines are based on dietary reference intakes (DRI) recommendations; protein intake across pregnancy should be upper than in non-pregnant women (around 71.0 g/day) (40). Protein intake in late pregnancy has been tra-

ditionally associated with a reduced risk of SGA birth and other maternal and fetal complications (18). For this reason, the general advice is that pregnant women eat more protein, specifically from high biologic value protein sources (41). In our study, the reference group, first quintile, has a mean intake of meat recorded below 99.6 g/day, nearly 30 g above the recommendation. This finding is consistent with a previous research which reported that Spanish pregnant women have a high intake of protein from meat sources (42). The association between meat intake and SGA has been investigated, showing controversial results. Some recent works carried out in European countries (France and Italy) have revealed that a diet in pregnancy based on meat was significantly associated with increased risk for SGA (26,27), whereas other authors have pointed out that mothers of AGA infants ate more servings of meat compared to SGA mothers during pregnancy (28). No detectable effect on meat intake over SGA risk has been seen in this study. Our result may be explained because the lowest quintile intake was above the recommendations, so the reference group is linked to low risk. If this low risk group is located under a saturation line, further increases in meat intake could not show additional risk reductions.

After adjusting by potential confounding factors, no significant association was found when each item was analyzed.

Table II. Frequency of maternal intake of different types of meat and meat products and risk of SGA newborn (n = 1,036)

	Cases (SGA)		Controls					
	n	(%)	n	(%)	cOR	95% CI	aOR*	95% CI
Meat								
<i>Egg (1 unit)</i>								
≤ 1-3 a month	38	(7.3)	36	(7.0)	1 (ref.)		1 (ref.)	
> 5 a week	42	(8.1)	32	(6.2)	1.22	(0.65-2.30)	0.98	(0.49-1.97)
<i>Chicken with skin (100-150 g)</i>								
Never	205	(39.6)	206	(39.8)	1 (ref.)		1 (ref.)	
> 2 a week	64	(12.4)	75	(14.5)	0.86	(0.58-1.27)	0.89	(0.60-1.38)
<i>Chicken without skin (100-150 g)</i>								
Never	83	(16.0)	62	(12.0)	1 (ref.)		1 (ref.)	
> 2 a week	131	(25.9)	149	(28.8)	0.65 [†]	(0.44-0.98)	0.70	(0.44-1.11)
<i>Beef (100-150 g)</i>								
Never	168	(32.4)	157	(30.3)	1 (ref.)		1 (ref.)	
> 1 a week	215	(41.5)	226	(43.6)	0.88	(0.66-1.19)	0.88	(0.63-1.22)
<i>Pork (100-150 g)</i>								
Never	70	(13.5)	72	(13.9)	1 (ref.)		1 (ref.)	
> 2 a week	93	(18.0)	102	(19.7)	0.94	(0.61-1.44)	0.87	(0.54-1.41)
<i>Lamb (100-150 g)</i>								
Never	376	(72.6)	364	(70.3)	1 (ref.)		1 (ref.)	
> 1 a week	37	(7.1)	36	(7.0)	0.97	(0.60-1.58)	0.90	(0.52-1.54)
<i>Rabbit (100-150 g)</i>								
Never	373	(72.0)	397	(76.6)	1 (ref.)		1 (ref.)	
> 1 a week	41	(7.9)	33	(6.4)	1.37	(0.84-2.22)	1.20	(0.70-2.06)
Meat products								
<i>Liver</i>								
Never	426	(82.2)	406	(78.4)	1 (ref.)		1 (ref.)	
> 1 a week	28	(5.4)	37	(7.1)	0.70	(0.41-1.19)	0.57	(0.31-1.03)
<i>Other entrails</i>								
Never	456	(88.0)	463	(89.4)	1 (ref.)		1 (ref.)	
> 1 a week	22	(4.3)	28	(5.4)	0.81	(0.46-1.41)	0.93	(0.50-1.72)
<i>Cured ham</i>								
Never	66	(12.7)	57	(11.0)	1 (ref.)		1 (ref.)	
> 5 a week	51	(9.9)	47	(9.1)	0.92	(0.54-1.58)	0.91	(0.49-1.67)
<i>Cooked ham</i>								
Never	92	(17.8)	62	(12.0)	1 (ref.)		1 (ref.)	
> 1 a day	28	(5.4)	39	(7.5)	0.48 [†]	(0.26-0.87)	0.59	(0.30-1.14)
<i>Processed meats</i>								
Never	108	(20.9)	97	(18.7)	1 (ref.)			
> 5 a week	35	(6.8)	28	(5.4)	1.12	(0.62-2.01)	1.09	(0.56-2.12)
<i>Pâté</i>								
Never	132	(25.5)	141	(27.2)	1 (ref.)		1 (ref.)	
> 1 a week	61	(11.8)	54	(10.4)	1.22	(0.78-1.90)	1.02	(0.61-1.69)
<i>Hamburger</i>								
Never	94	(18.2)	92	(17.8)	1 (ref.)		1 (ref.)	
> 1 a week	218	(42.1)	192	(37.1)	1.12	(0.79-1.58)	1.04	(0.70-1.53)
<i>Bacon</i>								
Never	248	(47.9)	249	(48.1)	1 (ref.)		1 (ref.)	
> 1 a week	114	(22.0)	86	(16.6)	1.35	(0.95-1.91)	1.12	(0.76-1.65)

cOR: crude odds ratio; aOR: adjusted odds ratio. *Adjusted for education level, pre-pregnancy body mass index, smoking, previous preterm/low birth weight newborn, newborn's gender, and MedDiet adherence. [†]Statistically significant (p < 0.05) association.

Table III. Daily maternal intake (g/day) grouped by quintiles (Q) of meat and meat products and risk of SGA newborn (n = 1,036)

	Cases (SGA)		Controls (AGA)		SGA			
	n	(%)	n	(%)	cOR	95% CI	aOR*	(95% CI)
Meat (g/day)								
Q1 (\leq 99.6)	108	(20.9)	104	(20.1)	1 (ref.)		1 (ref.)	
Q2 (99.7-126.8)	139	(26.8)	104	(20.1)	1.33	(0.91-1.96)	1.21	(0.79-1.86)
Q3 (126.9-149.2)	86	(16.6)	103	(19.9)	0.80	(0.54-1.19)	0.71	(0.46-1.10)
Q4 (149.3-184.3)	92	(17.8)	104	(20.1)	0.84	(0.56-1.25)	0.79	(0.51-1.24)
Q5 ($>$ 184.3)	93	(18.0)	103	(19.9)	0.86	(0.58-1.28)	0.79	(0.50-1.23)
p for trend					0.471		0.371	
Meat products (g/day)								
Q1 (\leq 2.7)	125	(24.1)	104	(20.1)	1 (ref.)		1 (ref.)	
Q2 (2.8-6.7)	111	(21.4)	104	(20.1)	0.89	(0.61-1.30)	0.90	(0.59-1.37)
Q3 (6.8-12.6)	88	(17.0)	103	(19.9)	0.72	(0.49-1.05)	0.67	(0.44-1.03)
Q4 (12.7-20.7)	93	(18.0)	104	(20.1)	0.74	(0.51-1.10)	0.62 [†]	(0.40-0.96)
Q5 ($>$ 20.7)	101	(19.5)	103	(19.9)	0.81	(0.54-1.21)	0.72	(0.46-1.12)
Q3-5 vs Q1-2					0.80	(0.62-1.02)	0.70 [†]	(0.53-0.93)
p for trend					0.868		0.649	

cOR: crude odds ratio; aOR: adjusted odds ratio. *Adjusted for education level, pre-pregnancy body mass index, smoking, previous preterm/low birth weight newborn, newborn's gender, and MedDiet adherence. [†]Statistically significant ($p < 0.05$) association.

However, when stratifying meat products by quintiles a protective effect comparing quintiles 3 to 5 versus 1-2 quintiles was found. This kind of food is typically associated to a western dietary pattern, and has been related to a higher risk of SGA (43); however, this is not supported by our data, possibly because in our population the reference group shows very low intakes, under 7 g/d, and the intakes for upper quintiles are also low. The most frequent advice in pregnancy diet is to avoid raw or undercooked meat products in order to prevent infectious foodborne diseases. Our results do not show any pernicious effect of meat products on the risk of SGA.

The current study has some limitations. Firstly, it is difficult to separate out the specific effects of single food (meat and meat products) on SGA risk because of the highly interrelated nature of dietary exposures; however, our results are adjusted by total energy and diet quality (Mediterranean diet adherence). Secondly, we assessed food intake after birth, so the information registered might not be representative of dietary habits throughout pregnancy time. However, the last gestational dietary patterns could be stable during pregnancy stage (44). Thirdly, although food intake was adjusted by energy consumption and other confounding factors, some residual confounding cannot be excluded. And fourth, a misclassification bias may have occurred. In this sense the information was gathered by midwives, and given that there is no definite knowledge on dietary risk factors and SGA, this bias would be similar in both groups (i.e., non-differential bias), shifting the strength of association toward the null value.

Our study includes some strengths. It includes a vast representative sample of a reference population (around 12,000 healthy

pregnant women attending Andalusian public hospitals). In addition, established Spanish fetal growth curves to define SGA have been used (30) and diet was collected throughout a FFQ validated in the Spanish population (35,45). Finally, the control group was sampled by density in the same hospitals (to avoid influence of season in diet reporting).

The present case-control study in Spanish women showed no effect of meat consumption over SGA risk, and a moderate protective effect of meat products intake. These findings support the advice of a varied diet for pregnant women, providing the intake of protein and other micronutrients from different food sources.

REFERENCES

1. De Onis M, Habicht JP. Anthropometric reference data for international use: recommendations from a WHO expert committee. *Food Nutr Bull* 1997;18(2):179-89.
2. Jensen EA, Foglia EE, Dysart KC, Simmons RA, Aghai ZH, Cook A, et al. Adverse effects of small for gestational age differ by gestational week among very preterm infants. *Arch Dis Child Fetal Neonatal Ed* 2018.
3. Premru-Srsen T, Verdenik I, Ponikvar BM, Steblonik L, Geršak K, Cerar LK. Infant mortality and causes of death by birth weight for gestational age in non-malformed singleton infants: a 2002-2012 population-based study. *J Perinat Med* 2018;46(5):547-53.
4. Huang YT, Lin HY, Wang CH, Su BH, Lin CC. Association of preterm birth and small for gestational age with metabolic outcomes in children and adolescents: a population-based cohort study from Taiwan. *Pediatr Neonatol* 2018;59(2):147-53.
5. Katz J, Wu LA, Mullany LC, Coles CL, Lee ACC, Kozuki N, et al. Prevalence of small-for-gestational-age and its mortality risk varies by choice of birth-weight-for-gestation reference population. *PLoS One* 2014;9(3).

6. Lee ACC, Katz J, Blencowe H, Cousens S, Kozuki N, Vogel JP, et al. National and regional estimates of term and preterm babies born small for gestational age in 138 low-income and middle-income countries in 2010. *Lancet Glob Health* 2013;1(1):e26-e36.
7. Govarts E, Izsatt N, Trnovec T, De Cock M, Eggesbø M, Palkovicova Murinova L, et al. Prenatal exposure to endocrine disrupting chemicals and risk of being born small for gestational age: pooled analysis of seven European birth cohorts. *Environ Int* 2018;115:267-78.
8. Woods JR, Heazell AEP. Stillbirth: is it preventable? *Obstet Gynaecol Reprod Med* 2018;28(5):148-54.
9. Crume TL, Brinton JT, Shapiro A, Kaar J, Glueck DH, Siega-Riz AM, et al. Maternal dietary intake during pregnancy and offspring body composition: the Healthy Start Study. *Am J Obstet Gynecol* 2016;215(5):609.e1-e8.
10. O'Connor DL, Blake J, Bell R, Bowen A, Callum J, Fenton S, et al. Canadian consensus on female nutrition: adolescence, reproduction, menopause, and beyond. *J Obstet Gynaecol Can* 2016;38(6):508-54.e18.
11. Gernand AD, Schulze KJ, Stewart CP, West KP, Christian P. Micronutrient deficiencies in pregnancy worldwide: health effects and prevention. *Nat Rev Endocrinol* 2016;12(5):274-89.
12. Dubois L, Diasparra M, Bédard B, Colapinto CK, Fontaine-Bisson B, Tremblay RE, et al. Adequacy of nutritional intake during pregnancy in relation to prepregnancy BMI: results from the 3D Cohort Study. *Br J Nutr* 2018;1-10.
13. Chattopadhyay N, Saumitra M. Developmental outcome in children with malnutrition. *J Nepal Paediatr Soc* 2016;36(2):170-7.
14. Yang J, Cheng Y, Pei L, Jiang Y, Lei F, Zeng L, et al. Maternal iron intake during pregnancy and birth outcomes: a cross-sectional study in Northwest China. *Br J Nutr* 2017;117(6):862-71.
15. Takimoto H, Mito N, Umegaki K, Ishiwaki A, Kusama K, Abe S, et al. Relationship between dietary folate intakes, maternal plasma total homocysteine and B-vitamins during pregnancy and fetal growth in Japan. *Eur J Nutr* 2007;46(5):300-6.
16. Hovdenak N, Haram K. Influence of mineral and vitamin supplements on pregnancy outcome. *Eur J Obstet Gynecol Reprod Biol* 2012;164(2):127-32.
17. Liberato SC, Singh G, Mulholland K. Effects of protein energy supplementation during pregnancy on fetal growth: a review of the literature focusing on contextual factors. *Food Nutr Res* 2013;57.
18. Bhutta ZA, Das JK, Rizvi A, Gaffey MF, Walker N, Horton S, et al. Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost? *Lancet* 2013;382(9890):452-77.
19. Haider BA, Olofin I, Wang M, Spiegelman D, Ezzati M, Fawzi WW. Anaemia, prenatal iron use, and risk of adverse pregnancy outcomes: systematic review and meta-analysis. *BMJ* 2013;346:f3443.
20. Simpson JL, Bailey LB, Pietrzik K, Shane B, Holzgreve W. Micronutrients and women of reproductive potential: required dietary intake and consequences of dietary deficiency or excess. Part I: folate, vitamin B12, vitamin B6. *J Matern Fetal Neonatal Med* 2010;23(12):1323-43.
21. De Gavelle E, Huneau JF, Mariotti F. Patterns of protein food intake are associated with nutrient adequacy in the general french adult population. *Nutrients* 2018;10(2).
22. Abu-Saad K, Fraser D. Maternal nutrition and birth outcomes. *Epidemiol Rev* 2010;32(1):5-25.
23. Ortega RM. Food, pregnancy and lactation: dietary guidelines for pregnant women. *Public Health Nutr* 2001;4(6A):1343-6.
24. Williamson CS. Nutrition in pregnancy. *Nutr Bull* 2006;31(1):28-59.
25. Naumenko NS, Nikonov AP, Aleksandrova LS, Stsaturova R, Belova V. Screening for infections in pregnancy: its significance for perinatal outcomes. *Voprosy Ginekologii, Akusherstva i Perinatologii* 2017;16(6):44-9.
26. Matta J, Carette C, Marchal CL, Bertrand J, Pétéra M, Zins M, et al. Dietary intake in young adults born small or appropriate for gestational age: data from the Haguenau cohort. *BMJ Open* 2016;6(7).
27. Ricci E, Chiaffarino F, Cipriani S, Malvezzi M, Parazzini F. Diet in pregnancy and risk of small for gestational age birth: results from a retrospective case-control study in Italy. *Matern Child Nutr* 2010;6(4):297-305.
28. Mitchell EA, Robinson E, Clark PM, Becroft DMO, Glovish N, Pattison NS, et al. Maternal nutritional risk factors for small for gestational age babies in a developed country: a case-control study. *Arch Dis Child Fetal Neonatal Ed* 2004;89(5):F431-F5.
29. Lu MS, Chen QZ, He JR, Wei XL, Lu JH, Li SH, et al. Maternal dietary patterns and fetal growth: a large prospective cohort study in China. *Nutrients* 2016;8(5).
30. Delgado Beltrán P, Melchor Marcos JC, Rodríguez-Alarcón Gómez J, Linares Uribe A, Fernández-Llebreg del Rey L, Barbazan Cortes MJ, et al. The fetal development curves of newborn infants in the Hospital de Cruces (Vizcaya). II. Length, head circumference and ponderal index. *An Esp Pediatr* 1996;44(1):55-9.
31. Álvarez-Dardet CAJ, Domingo A, Regidor E. La medicina de la clase social en Ciencias de la Salud. Informe de un Grupo de Trabajo de la Sociedad Española de Epidemiología. SG Editores Ed.; 1995.
32. Black D TP. Inequalities in health. Harmondsworth: Penguin; 1983.
33. Trichopoulou A, Costacou T, Bamia C, Trichopoulos D. Adherence to a Mediterranean diet and survival in a Greek population. *N Engl J Med* 2003;348(26):2599-608.
34. Kessner DM SJ, Kalk CE, Schlesinger ER. Washington, D.C.: Institute of Medicine and National Academy of Sciences; 1973.
35. Martín-Moreno JM, Boyle P, Gorgojo L, Maisonneuve P, Fernández-Rodríguez JC, Salvini S, et al. Development and validation of a food frequency questionnaire in Spain. *Int J Epidemiol* 1993;22(3):512-9.
36. Fernández-Ballart JD, Piñol JL, Zazpe I, Corella D, Carrasco P, Toledo E, et al. Relative validity of a semi-quantitative food-frequency questionnaire in an elderly Mediterranean population of Spain. *Br J Nutr* 2010;103(12):1808-16.
37. Mataix Verdú J. Tabla de composición de alimentos españoles (Spanish food composition tables). Granada: Universidad de Granada; 2003.
38. Moreiras O CA, Cabrera L, Cuadrado C. Tablas de composición de alimentos. (Spanish food composition tables). Madrid, Spain: Pirámide; 2003.
39. Willet W. Nutritional epidemiology. 3rd ed. New York: Oxford University Press; 2013.
40. Trumbo P, Schlicker S, Yates AA, Poos M. Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein and amino acids. *J Am Diet Assoc* 2002;102(11):1621-30.
41. Lee A, Belski R, Radcliffe J, Newton M. What do pregnant women know about the healthy eating guidelines for pregnancy? A web-based questionnaire. *Matern Child Health J* 2016;20(10):2179-88.
42. Rodríguez-Bernal CL, Ramón R, Quiles J, Murcia M, Navarrete-Muñoz EM, Vioque J, et al. Dietary intake in pregnant women in a Spanish Mediterranean area: as good as it is supposed to be? *Public Health Nutr* 2013;16(8):1379-89.
43. Lowensohn RI, Stadler DD, Naze C. Current concepts of maternal nutrition. *Obstet Gynecol Surv* 2016;71(7):413-26.
44. Crozier SR, Robinson SM, Godfrey KM, Cooper C, Inskip HM. Women's dietary patterns change little from before to during pregnancy. *J Nutr* 2009;139(10):1956-63.
45. De la Fuente-Arrillaga C, Vázquez Ruiz Z, Bes-Rastrollo M, Sampson L, Martínez-González MA. Reproducibility of an FFQ validated in Spain. *Public Health Nutr* 2010;13(9):1364-72.