



Trabajo Original

Valoración nutricional

A pilot intervention to reduce postpartum weight retention at primary health care in Brazil *Una intervención piloto para reducir la retención de peso después del parto en la atención primaria de salud en Brasil*

Karina dos Santos^{1,2}, Paulla Tavares Patrício¹, Thaïssa Santana Vieira Lima¹, Denise Cavalcante de Barros^{1,3} and Cláudia Saunders^{1,2}

¹Maternal and Child Health Research Group (GPSM). Universidade Federal do Rio de Janeiro (UFRJ). Ilha do Fundão, Rio de Janeiro. Brazil. ²Department of Nutrition and Dietetics. Josué de Castro Nutrition Institute. Universidade Federal do Rio de Janeiro (UFRJ). Ilha do Fundão, Rio de Janeiro. Brazil. ³National School of Public Health. Fundação Oswaldo Cruz - ENSP/FIOCRUZ. Manginhos, Rio de Janeiro. Brazil

Abstract

Background: postpartum weight retention (PWR) strongly predicts obesity, the major nutritional concern of this century; however, there is a gap in nutritional care for postpartum women in Brazil.

Objective: to evaluate the effect of nutritional counselling provided at primary health care on the reduction of PWR.

Method: pilot study carried out in a low-income community in southeastern Brazil, involving postpartum women with PWR, who were provided with individual monthly appointments with a nutritionist over three months of follow-up. Nutritional counselling was based on the Dietary Approach to Stop Hypertension (DASH diet) aiming at healthy and gradual weight loss. Anthropometric evaluation included measurement of weight, height, waist circumference and percentage of body fat. Dietary intake was assessed using a food frequency questionnaire and adherence to diet was evaluated using a DASH score.

Results: women who participated in the study (n = 26) showed a reduction in PWR (median -1.80 kg, p = 0.004), body mass index (-0.57 kg/m², p = 0.004), and waist circumference (-2.50 cm, p = 0.024), as well as 91.67% of them presented good adherence to diet.

Conclusion: nutritional counselling provided to low-income postpartum women at primary health care contributed to the reduction of PWR, body mass index and waist circumference, as the study participants presented good adherence to a healthy dietary pattern.

Key words:

Postpartum period.
Maternal nutrition.
Women's health.
Primary health care.
Public health.

Resumen

Introducción: la retención de peso después del parto (RPDP) predice firmemente la obesidad, que es la principal preocupación nutricional de este siglo; sin embargo, existe una brecha en la atención nutricional para las mujeres después del parto en Brasil.

Objetivo: evaluar el efecto del asesoramiento nutricional brindado en Atención Primaria sanitaria sobre la reducción de RPDP.

Método: estudio piloto realizado en una comunidad de bajos ingresos en el sureste de Brasil, en el que participaron mujeres postparto con RPDP, que recibieron citas mensuales individuales con un nutricionista durante tres meses de seguimiento. El asesoramiento nutricional se basó en la dieta Dietary Approach to Stop Hypertension (DASH) y apuntó a una pérdida de peso saludable y gradual. La evaluación antropométrica incluyó la medición del peso, la altura, la circunferencia de la cintura y el porcentaje de grasa corporal. La ingesta dietética se evaluó mediante un cuestionario de frecuencia alimentaria y la adherencia a la dieta se evaluó mediante un *score* DASH.

Resultados: las mujeres que participaron en el estudio (n = 26) mostraron una reducción en la RPDP (mediana de -1,80 kg, p = 0,004), índice de masa corporal (-0,57 kg/m², p = 0,004) y circunferencia de la cintura (-2,50 cm, p = 0,024); el 91,67% de ellas presentaron buena adherencia a la dieta.

Conclusión: el asesoramiento nutricional brindado a las mujeres de bajos ingresos después del parto en Atención Primaria sanitaria contribuyó a la reducción de la RPDP, el índice de masa corporal y la circunferencia de la cintura, ya que los participantes del estudio presentaron una buena adherencia a un patrón de dieta saludable.

Palabras clave:

Periodo posparto.
Nutrición materna.
Salud de la mujer.
Atención Primaria sanitaria.
Salud pública.

Received: 18/01/2019 • Accepted: 26/04/2019

Author's contributions: KS, DCB, and CS designed the study; KS, PTP, and TSVL participated in the data collection and intervention; KS, DCB, and CS analysed the data and wrote the manuscript. All authors read, reviewed, and approved the final manuscript.

Santos K, Patrício PT, Lima TSV, de Barros DC, Saunders C. A pilot intervention to reduce postpartum weight retention at primary health care in Brazil. *Nutr Hosp* 2019;36(4):854-861

DOI: <http://dx.doi.org/10.20960/nh.02508>

Correspondence:

Karina dos Santos. Centro de Ciências da Saúde, bloco J, 2º andar, sala 26. Av. Carlos Chagas Filho, 373. 21941-590 Ilha do Fundão, Rio de Janeiro. Brazil
e-mail: karsantos@gmail.com

INTRODUCTION

Overweight among women of reproductive age has been growing at an alarming rate in recent decades, becoming a serious public health issue in Brazil. Currently, 50.5% of adult women are overweight and 19.6% are obese, with a higher risk of developing chronic diseases and a tendency towards worse obstetric outcomes (1,2).

The pregnancy-puerperal cycle involves physiological adaptations that induce significant changes in body weight and composition (3). Excessive gestational weight gain and postpartum weight retention (PWR) are strong predictors of obesity development (3-5). The greater the gestational weight gain, the higher the PWR verified in one year, and the higher the body mass index (BMI) in subsequent years (5). Therefore, it is recommended that women return to pre-gestational weight within 12 months after delivery (5,6).

Nutritional counselling is effective in promoting PWR reduction (7-10). However, there is a lack of a specific protocol at Brazilian public primary health care to monitor health and nutrition of women in the postpartum period (11). Moreover, commitment to nutritional care during the postpartum period is challenging due to the demands of motherhood, which are notably worse for low-income women (12). Thus, there is a current call to action to development of new strategies that could better achieve this specific public (11,12).

The present study aimed to evaluate the effect of nutritional counselling at primary health care on the reduction of PWR in a low-income community in southeastern Brazil.

MATERIAL AND METHODS

STUDY DESIGN AND POPULATION

This pilot study was conducted from November 2016 to September 2017 in two primary health care centers located in a low-income community in southeastern Brazil.

All women registered in the health center's information system were identified with the potential to participate in the study. Individuals that met the following inclusion criteria were selected: women 6-9 months postpartum, with weight retention of any magnitude, BMI above 18.5 kg/m², older than 18 years old, single-fetus pregnancy, a living infant, absence of diagnosis of chronic diseases (except for obesity), no history or diagnosis of eating disorders, and no bariatric surgery.

Individuals were invited to participate through phone calls or personally in the health centers waiting rooms and childcare groups. The project was widely disseminated among health professionals and community using posters and leaflets. Women who agreed to participate had their first appointment scheduled, during which the data regarding sociodemographic characteristics, obstetric history, pre- and postnatal care, and conditions of the child at birth were verified through a face-to-face interview and medical records.

INTERVENTION

The study participants attended four individual appointments with a nutritionist over three months of follow-up. Nutritional counselling was performed based on the Dietary Approach to Stop Hypertension (DASH) diet (13). The DASH diet is recognized as a healthy and beneficial dietary pattern for women of reproductive age (14). Most daily energy consumption should derive from plant foods, such as fruits, vegetables, seeds, oilseeds, beans, and whole grains, restricting the consumption of meat and other foods with high amounts of saturated fat, salt, and sugar (13). However, it is an American dietary pattern, so it had to be translated and adapted for this study, especially considering typical foods that were accessible to the Brazilian population (in press).

The meal plan delivered to women was organized into six meals, setting up a daily menu model with the recommended number of portions of the adapted DASH diet groups: 7-8 servings of cereals, preferably whole-grain cereals; 4-5 servings of vegetables; 4-5 servings of fruits; 2-3 servings of dairy products, preferably low-fat; 2-3 servings of oils; 1-2 servings of beans (typical Brazilian food); and up to two servings of meats. The consumption of seeds and/or nuts was indicated 4-5 times per week. Sugars and sugary beverages were advised against, without definition of a tolerable portion [in press].

The nutritional guidance material was developed in line with the proposal of the new Food Guide for the Brazilian Population (15), similar to others already used in the routine practice where the study was developed. A daily energy value of approximately 2,000 kcal was considered to meet the recommendation of minimum intakes for nursing mothers (1,900 kcal/day) (16).

The meal plan was delivered and explained in the first appointment. General recommendations were also made for a healthy diet, such as the amount of water intake, replacement of industrial seasonings with natural herbs and spices, reduction of consumption of ultra-processed food and moderation in the addition of salt, sugar, and fats. In subsequent consultations, with 30-day intervals, all orientation was reinforced or adjusted, according to the obtained results.

Despite the use of a common meal plan for all participants, nutritional counselling was conducted in an individualized and contextualized way based on the routine of each woman. Suggestions for foods and preparations were discussed with each participant, according to availability and affordability of certain items (17,18). Among available options, the best choices were suggested, so that women would be closest to the adapted DASH diet proposal. Physical activity was encouraged throughout the follow-up period and evaluated before and after the intervention (19).

To improve adherence to the diet, study participants received a kit containing three packages of skimmed milk powder (400 g each), two packages of linseed and/or sunflower seeds (250 g) and three seed and cereal bars (30 g each), at each visit.

In addition, as part of the intervention, the free instant messaging application WhatsApp® and the social media site Facebook® were used as tools to reinforce nutritional counselling (10,20,21). Messages regarding information on healthy eating were sent weekly. These digital tools were also used to communicate with

the study participants, in order to promote self-motivation and self-monitoring (10,20,21). Women were encouraged to send pictures of meals and questions about food and nutrition, always in a private way, and were receiving answers from expert nutritionists.

ASSESSMENT OF DIET ADHERENCE

Food intake was assessed in the last appointment using a food frequency questionnaire (FFQ) validated for the Brazilian population (22). Analyses of individual food consumption were performed in Microsoft Excel. The amount consumed for each food, reported in usual units, was converted to grams/milligrams or liters/milliliters.

The frequencies of consumption reported in the FFQ as weekly or monthly were transformed into daily frequency (23). Using the Table of Nutritional Composition of Foods Consumed in Brazil (24), it was possible to verify daily consumption in kilocalories of energy and grams or milligrams of nutrients.

To evaluate the diet adherence, a DASH score (25) was used, considering the number of recommended portions in each food group, for each 1,000 kcal daily consumed: fruits (> 2.10 servings of 80 g), vegetables (> 2.10 servings of 50 g), dairy products (> 0.96 portion of 120 ml of milk or 40 g of cheese), whole grains (> 3.60 servings of 20 g), seeds/legumes (\geq 0.33 portion of 30 g of seeds/oilseeds or 80 g of beans), meat (\leq 0.80 portion of 170 g), oils/fats (\leq 1.20 portion of 5 ml) and percentage of saturated fat (< 7% of the total daily energy value).

For each food group in which the consumption recommendation was met, one point was recorded in the DASH score (25). Therefore, the score ranging from 0 (lower adherence) to 8 (higher adherence) was individually calculated. A score equal or greater than 4 was considered as good adherence to diet, and scores less than 4, as bad adherence.

ANTHROPOMETRIC EVALUATION

Anthropometric evaluation was performed in all visits, with measurements of weight (kg), percentage of body fat (%) and waist circumference (cm). Weight and percentage of body fat were measured with an electronic field balance, with bipolar bioimpedance. Participants removed their shoes and accessories and wore light clothing (26). Waist circumference was measured with a flat steel anthropometric tape, at the mean height between the iliac crest and lower costal margin (26).

Height (m) was measured at the first visit, with a fixed vertical stadiometer, in the absence of shoes and hair accessories (26). Blood pressure was measured in all consultations using a high precision digital device. The analyses for determining nutritional status were performed according to the current protocols published by the Ministry of Health of Brazil (26).

PWR was defined as the difference between the actual weight and pre-gestational weight found on the medical records. In the absence of such information, the weight recorded at the first prenatal

visit was used, if it has occurred during the first trimester. The BMI was calculated using the formula $BMI = \text{weight (kg)}/\text{height}^2 \text{ (m)}$.

STATISTICAL ANALYSIS

PWR was considered as the primary outcome of the study. Secondary outcomes were weight, BMI, waist circumference and body fat percentage. Due to the reduced sample size and its non-normality condition, tested with the Kolmogorov-Smirnov test, non-parametric statistical tests were adopted.

For the descriptive analyses, the results were presented as absolute and relative frequency for the qualitative variables and as median and interquartile range for the quantitative variables. The Mann-Whitney test was used to compare the medians of independent samples, while the Wilcoxon test was used for the comparison between dependent samples. The Chi-square test was used to compare qualitative variables using the Fisher's test.

Spearman's correlation was used to verify possible associations between covariables and the main outcome. Analyses were performed in the IBM SPSS Statistics version 21.0 software, adopting $p < 0.05$ for statistical significance of the results.

ETHICAL STANDARDS DISCLOSURE

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures were approved by the Ethics Committee in Research of the National School of Public Health (CAAE 0251.0.031.031-10) and Brazilian Registry of Clinical Trials (RBR-4t46ry). Written informed consent was obtained from all study participants.

RESULTS

Following consultation of the records in the health center's information system, 478 women were identified, who would be 6-9 months postpartum during the time of data collection. Of these, 283 could not be reached. Among those who accepted to participate in the study and attended the first visit ($n = 60$), 19 did not meet the eligibility criteria. Forty-one women started the intervention and 26 completed it, accounting for a 36.6% loss of follow-up (Fig. 1). The mean duration of follow-up, encompassing the four visits, was 109 days (3.6 months).

The characteristics of the sample, as well as the comparison between women who completed or did not completed the study, are presented in table I. The median PWR was 5.50 kg. Regarding the nutritional status, 48.8% ($n = 20$) were obese and 39% ($n = 16$) were overweight. The median pre-gestational BMI value was 27.06 kg/m² (24.63-30.77). There was a greater loss of follow-up for younger women with higher BMI values ($p < 0.05$) (Table I). There were no significant differences in relation to the sociodemographic characteristics between women who completed or did not completed the study (Table I).




Figure 1.
Flowchart of the study.

The sample contained a majority of black women (68.3%), married (85.4%), with high school education (65.9%), adequate sanitation at home (87.8%), multiparous (56.1%) and with low

adherence to physical activities (78.1%). Most women maintained breastfeeding (70.7%), but exclusive breastfeeding up to six months was reported by only nine of them (22%).

Table II shows the anthropometric results before and after the intervention. Measurements of all parameters presented a reduction ($p < 0.05$) after intervention, except for body fat percentage ($p = 0.346$). The median systolic (SBP) and diastolic (DBP) blood pressure did not change after the intervention (initial SBP = 109 *versus* final SBP = 112, $p = 0.909$ and baseline DBP = 68 *versus* final DBP = 68.5, $p = 0.808$).

Regarding the adherence to diet (Table III), the frequency of women who consumed the portions indicated for each food group was 50.00% for fruits, 33.33% for vegetables, 87.50% for dairy products, 16.67% for whole grains, 91.67% for seeds and legumes, 70.83% for oils and fats, 91.67% for meat and 75.00% for saturated fat. Analysis of DASH score showed that 91.67% of women presented good adherence to diet. Two women did not respond to the FFQ, as they claimed little time for the consultation.

About the level of physical activity, only three of the 26 women who completed the consultations were classified as “active” and this was maintained before and after the intervention. All others were “irregularly active” ($n = 14$) or “sedentary” ($n = 9$). Therefore, it was not possible to evaluate the potential impact of physical activity on postpartum weight retention.

Time after delivery and the breastfeeding period (in days) presented a negative correlation with PWR after the intervention ($\rho -0.406$, $p = 0.040$ and $\rho -0.437$, $p = 0.025$, respectively). Energy intake presented a positive correlation with PWR ($\rho 0.536$, $p = 0.007$), which did not occur for the individually assessed food groups. No correlation was observed between PWR and the following covariates tested: maternal age ($p = 0.244$), gestational weight gain ($p = 0.203$), parity ($p = 0.171$) and per capita family income ($p = 0.677$).

DISCUSSION

Our results showed that nutritional care offered to women during the postpartum period, at primary health care, contributed to the reduction of PWR, BMI and waist circumference, as women presented good adherence to a healthy dietary pattern.

The location of the study was a neighborhood with elevated levels of violence and police confrontations, which contributed to a limitation of the study, which is the small sample size. The selection and follow-up of women were constantly hampered by this social context, especially exacerbated during the study period. Lack of security limits the routine activities of women residing on-site. Attending consultations, following a diet plan, and focusing on personal goals related to nutritional status has become especially challenging in this hostile environment.

To mitigate the damages caused by these conditions, it was strategic to use communication technologies (WhatsApp® and Facebook®). Thus, it was possible to maintain communication with the study participants and promote motivation for adherence to nutritional guidelines, even when the attendance schedule was compromised. Participants sent photos, questions and comments

Table I. Characteristics of participants at baseline and comparison between individuals who have or have not completed the intervention (n = 41)

	Total (n = 41)		Did not complete the intervention (n = 15)		Completed the intervention (n = 26)		p
	Med	IR	Med	IR	Med	IR	
Age, years	29.00	24.00-33.00	24.00	21.00-31.00	31.00	25.75-33.25	0.032*
Pre-gestational BMI, kg/m ²	27.06	24.63-30.77	29.59	26.99-32.66	25.81	23.30-29.38	0.058*
BMI, kg/m ²	29.92	26.60-35.13	30.90	29.20-37.20	28.59	25.81-31.71	0.021*
PWR, kg	5.50	2.45-10.65	7.60	2.60-12.90	4.05	2.22-8.52	0.120*
	n	%	n	%	n	%	p
<i>Color of the skin</i>							
White/yellow	13.00	31.70	4.00	26.70	9.00	34.60	0.734 [‡]
Brown/black	28.00	68.30	11.00	73.30	17.00	65.40	
<i>Marital status</i>							
With partner	35.00	85.40	13.00	86.70	22.00	84.60	1.000 [‡]
Without partner	6.00	14.60	2.00	13.30	4.00	15.40	
<i>Education</i>							
Incomplete HS	14.00	34.10	5.00	33.30	9.00	34.60	0.934 [‡]
Complete HS	27.00	65.90	10.00	66.70	17.00	65.40	
<i>Work</i>							
Yes	19.00	46.30	6.00	40.00	13.00	50.00	0.536 [‡]
No	22.00	53.70	9.00	60.00	13.00	50.00	
<i>Family income per capita[§]</i>							
Up to 0.5 MW	17.00	41.50	7.00	46.70	10.00	38.50	0.680 [‡]
0.5 MW or more	23.00	56.10	8.00	53.30	15.00	57.70	
<i>Sanitation</i>							
Suitable	36.00	87.80	12.00	80.00	24.00	92.30	0.336 [‡]
Unsuitable	5.00	12.20	3.00	20.00	2.00	7.70	
<i>Parity</i>							
Primiparous	18.00	43.90	8.00	53.40	10.00	38.50	0.355 [‡]
Multiparous	23.00	56.10	7.00	46.60	16.00	61.50	
<i>PA level</i>							
Active	9.00	21.90	2.00	13.30	7.00	26.90	0.444 [‡]
Irregularly active	20.00	48.80	7.00	46.70	13.00	50.00	
Sedentary	12.00	29.30	6.00	40.00	6.00	23.10	
<i>Breastfeeding</i>							
Yes	29.00	70.70	10.00	66.70	19.00	73.10	0.664 [‡]
No	12.00	29.30	5.00	33.30	7.00	26.90	
<i>6 m exclusive breastfeeding</i>							
Yes	9.00	22.00	3.00	20.00	6.00	23.10	1.000 [‡]
No	32.00	78.00	12.00	80.00	20.00	76.90	

Med: median; IR: interquartile range; BMI: body mass index; PWR: postpartum weight retention; HS: high school; MW: minimum wage; PA: physical activity; 6 m: six months. *Mann-Whitney test. [‡]Chi-square test. [‡]Fisher's test. Confidence interval 95%. [§]Missing (n = 1), 2.4%.

about food and nutrition. In this way, it was possible to provide constant support and enhance the link between the research team and study participants.

A second limitation refers to the lack of a comparison group (control). This is because the project is part of an initiative to

increase the supply of services to the population of the territory under study. Therefore, it was understood that all identified eligible women who expressed an interest in participation should have access to nutritional care. This option limited the elaboration of more robust conclusions regarding the cause and effect relationship of

Table II. Comparison of anthropometric measurements before and after the intervention (n = 26)

Outcome	Baseline/1 st visit		After intervention/4 th visit [‡]		Δ median 4 th visit-baseline [§]	p [†]
	Med	IR	Med	IR		
PWR (kg)	4.05	2.22-8.52	2.25	0.82-5.97	-1.80	0.004
Weight (kg)	70.55	66.15-77.40	69.00	64.25-75.25	-1.55	0.004
BMI (kg/m ²)	28.59	25.81-31.71	28.02	24.79-30.84	-0.57	0.004
WC (cm) [¶]	89.75	80.87-97.30	87.25	79.75-96.50	-2.50	0.024
BF (%)	36.80	34.07-40.37	35.75	33.35-40.85	-1.05	0.346

Med: median; IR: interquartile range; PWR: postpartum weight retention; BMI: body mass index; WC: waist circumference; BF: body fat. ^{*}Mean time between 1st and 2nd visits: 38 days. [†]Mean time between 2nd and 3rd visits: 37 days. [‡]Mean time between 3rd and 4th visits: 34 days. [§]Mean duration of the intervention: 109 days (3.6 months). [¶]Wilcoxon test, comparison with baseline, confidence interval 95%. ^{††}Missing (n = 2).

Table III. Assessment to diet adherence and DASH score (n = 24)

Food groups (servings per 1,000 kcal of EI)	Consumed (median number of servings)	IR	Recommended (number of servings)	Frequency of adequacy* n (%)
Fruits	2.09	1.53-2.63	> 2.10	12 (50.00)
Vegetables	1.80	0.92-2.79	> 2.10	8 (33.33)
Dairy food	2.03	1.14-3.17	> 0.96	21 (87.50)
Whole grains	0.58	0.00-1.30	> 3.60	4 (16.67)
Seeds, nuts/beans	1.07	0.74-1.38	≥ 0.33	22 (91.67)
Oil/fat	0.96	0.47-1.54	≤ 1.20	17 (70.83)
Meat	0.42	0.23-0.57	≤ 0.80	22 (91.67)
Saturated fat (% of EI)	6.05	5.28-7.29	< 7.00	18 (75.00)
DASH score (points)	5.00	4.00-6.00	8.00	Score < 4.00 = 2 (8.33%) Score ≥ 4.00 = 22 (91.67%)

EI: energy intake; IR: interquartile range; DASH: Dietary Approach to Stop Hypertension. Median (IR) of daily energy intake: 1,876.19 (1,639.80-2,488.42) kcal. ^{*}Number and percentage of women who individually reached the recommended intake.

the intervention; however, it increased the range of the benefits resulting from the study.

Nutritional care initiated during pregnancy and continued postpartum is more effective in reducing weight retention than those provided only after delivery (7,9). However, in the present study, the intervention was initiated in the late puerperium. It is likely that the results may be more robust if counselling and monitoring of gestational weight gain were provided during prenatal care.

To date, no studies have been identified using the DASH diet to reduce PWR. The Mediterranean diet, with similar composition, was tested in a US study (27), over four months, in overweight postpartum women. The difference after the intervention was -2.31 kg in weight, -0.85 kg/m² in BMI, -3.47 cm in waist circumference and -1.19% in the percentage of body fat. These results were similar to ours (-1.55 kg, -0.57 kg/m², -2.50 cm and -1.05%, respectively). However, the American study was one month longer than ours.

In a survey performed at primary health care in Sweden (9), overweight postpartum women who received individual nutritional coun-

selling for 12 weeks showed a mean reduction of 6.1 kg in weight at the end of the intervention. This weight loss was higher than the one presented here, however, an energy restriction of 500 kcal was applied in the individual food plan. In the present study, there was no energy restriction, because the meal plan had a qualitative focus.

There was a greater follow-up for older women with lower BMI values, which cannot be explained with other sociodemographic factors. However, the values of pre-gestational BMI were lower among these women, with a borderline result for statistical significance (25.81 kg/m² versus 29.59 kg/m², p = 0.058). The lower pre-gestational BMI is associated with greater weight change during and after gestation (3,4). It is possible that this more pronounced body change has motivated participants to better adhere to nutritional care than those who already have higher pre-gestational BMI.

We observed good adherence to the adapted DASH diet, especially limiting the intake of meat and saturated fat while improving consumption of dairy products and seeds, nuts and beans. These recommendations were emphasized during the nutritional orien-

tation because they are recognized DASH markers (13) and are part of the recommendations of the Food Guide for the Brazilian Population (15). Thus, these results were associated with reduction of RPPP in other studies (28-30).

It is important to consider that adopting a healthy dietary pattern is a challenge for low-income women living in areas of social vulnerability, either because of financial constraints or little availability of fresh and non-industrialized foods, locally (31,32). Popular beliefs and food-related myths that prioritize the consumption of foods with higher energy density during the lactation period were also verified (33). Whole grains, vegetables and fruits had the lowest frequency of consumption adequacy, which may be associated to some of these factors. On the other hand, the better adherence to consumption of dairy products and seeds/nuts was possibly favored by the donation of these products during the intervention.

Another relevant issue, reported by a large part of the study participants, was the lack of time due to the excess of responsibilities related to family and maternity, which was considered as the main barrier to attend appointments, to commit to physical activity and to adhere to the food plan. Most studies involving postpartum women reported large loss of follow-up, generally associated with such factors (12,35).

However, the improvement in the anthropometric indicators of nutritional status and overall diet quality after the intervention indicate that, even though there is limited access to some types of foods and barriers to postpartum care for women's health, adherence to most of the recommendations has been achieved. Therefore, it is important to implement personalized guidelines to meet individual's needs, with the aim to provide available and affordable food options and share practical suggestions for meal planning and preparation.

It is known that weight loss is natural during the first year after delivery, due to the return to physiological and anatomical pre-gestational conditions and to breastfeeding (34). In fact, postpartum and breastfeeding time showed negative correlations with PWR at the end of the study. Corroborating previous findings (36), the total daily energy intake, rather than the consumption of certain food groups, correlated positively with the outcome.

The DASH diet was originally developed for the reduction of systemic arterial pressure (13). In this study, it was not possible to observe a decrease in SBP and DBP; however, the median blood pressure levels were within the normal range before and after the intervention.

Despite the unfavorable context, it was possible to successfully implement this pilot intervention in a community of economic and social vulnerability in southeast Brazil. There was a significant decrease in PWR, BMI and waist circumference. Positive results can be attributed to the good adherence to the adapted DASH diet and to contextualized nutritional counselling, with a focus on women's health. It is relevant to highlight the importance of the digital communication tools used to maintain the follow-up, although this strategy was not measured alone as an outcome of the study, but rather as part of it.

Larger trials are needed to show whether counselling on diet can improve nutritional status of postpartum women and how it should

be effectively done. Currently, there is no protocol that prioritizes and guides nutritional care for women in the postpartum period at Brazilian public primary health care. Thus, this kind of study may support the improvement of pre- and postnatal care in public health services.

FINANCIAL SUPPORT

This research was supported by the "Teias-Escola Manguinhos" project, a partnership between Osvaldo Cruz Foundation (Fundação Oswaldo Cruz) and the Municipal Secretary of Health and Civil Defense of Rio de Janeiro (Secretaria Municipal de Saúde e Defesa Civil do Rio de Janeiro). This work is part of a master's degree project, which had a scholarship supported by the Coordination for the Improvement of Higher Education Personnel (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - CAPES) and Foundation for Research Support of the State of Rio de Janeiro (Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro - FAPERJ). The funders had no role in the design, analysis or writing of this article.

ACKNOWLEDGMENTS

The authors would like to acknowledge Ana Lúcia Fittipaldi, Roberta Araújo and Beatriz Della Libera for the precious support regarding the participant recruitment and data collection; and thank all the women that volunteered for the study.

REFERENCES

1. Ministry of Health of Brazil. *Vigitel Brasil 2016*. Brasília: Ministry of Health; 2017.
2. Howell KR, Powell TL. Effects of maternal obesity on placental function and fetal development. *Reproduction* 2017;153(3):R97-R108.
3. Bogaerts A, De Baetselier E, Ameys L, Dilles T, Van Rompaey B, Devlieger R. Postpartum weight trajectories in overweight and lean women. *Midwifery* 2017;49:134-41.
4. Widen EM, Whyatt RM, Hoepner LA, Ramirez-Carvey J, Oberfield SE, Hassoun A, et al. Excessive gestational weight gain is associated with long-term body fat and weight retention at 7 y postpartum in African American and Dominican mothers with underweight, normal, and overweight prepregnancy BMI. *Am J Clin Nutr* 2015;102(6):1460-7.
5. Linné Y, Dye L, Barkeling B, Rössner S. Long-term weight development in women: a 15-year follow-up of the effects of pregnancy. *Obes Res* 2004;12(7):1166-78.
6. Stang J, Huffman LG. Position of the Academy of Nutrition and Dietetics: obesity, reproduction, and pregnancy outcomes. *J Acad Nutr Diet* 2016;116(4):677-91.
7. Huang TT, Yeh CY, Tsai YC. A diet and physical activity intervention for preventing weight retention among Taiwanese childbearing women: a randomised controlled trial. *Midwifery* 2011;27(2):257-64.
8. Jaakkola J, Isolauri E, Poussa T, Laitinen K. Benefits of repeated individual dietary counselling in long-term weight control in women after delivery. *Matern Child Nutr* 2015;11(4):1041-8.
9. Huseinovic E, Bertz F, Leu Agelii M, Hellebø Johansson E, Winkvist A, Brekke HK. Effectiveness of a weight loss intervention in postpartum women: results from a randomized controlled trial in primary health care. *Am J Clin Nutr* 2016;104(2):362-70.
10. Herring SJ, Cruice JF, Bennett GG, Darden N, Wallen JJ, Rose MZ, et al. Intervening during and after pregnancy to prevent weight retention among African American women. *Prev Med Rep* 2017;7:119-23.

11. Cornell A, McCoy C, Stampfel C, Bonzon E, Verbiest S. Creating new strategies to enhance postpartum health and wellness. *Matern Child Health J* 2016;20(Suppl 1):39-42.
12. Graham M, Uesugi K, Olson C. Barriers to weight-related health behaviours: a qualitative comparison of the socioecological conditions between pregnant and post-partum low-income women. *Matern Child Nutr* 2016;12(2):349-61.
13. Windhauser MM, Ernst DB, Karanja NM, Crawford SW, Redican SE, Swain JF, et al. Translating the Dietary Approaches to Stop Hypertension diet from research to practice: dietary and behavior change techniques. *J Am Diet Assoc* 1999;99(Suppl 8):S90-5.
14. Tabibian S, Daneshzad E, Bellissimo N, Brett NR, Dorosty-Motlagh AR, Azadbakht L. Association between adherence to the Dietary Approaches to Stop Hypertension diet with food security and weight status in adult women. *Nutr Diet* 2018;75(5):481-7.
15. Ministry of Health of Brazil. *Guia Alimentar para a População Brasileira*. 2nd ed. Brasília: Ministry of Health; 2014.
16. Kolasa KM, Firnhaber G, Haven K. Diet for a healthy lactating woman. *Clin Obstet Gynecol* 2015;58(4):893-901.
17. Whitt-Glover MC, Hunter JC, Foy CG, Quandt SA, Vitolins MZ, Leng I, et al. Translating the Dietary Approaches to Stop Hypertension (DASH) diet for use in underresourced, urban African American communities, 2010. *Prev Chronic Dis* 2013;10:120088.
18. Mackenbach JD, Burgoine T, Lakerveld J, Forouhi NG, Griffin SJ, Wareham NJ, et al. Accessibility and affordability of supermarkets: associations with the DASH Diet. *Am J Prev Med* 2017;53(1):55-62.
19. Matsudo SM, Araújo TL, Matsudo VKR, Andrade DR, Andrade EL, Oliveira LC, et al. Questionário Internacional de Atividade Física (IPAQ): estudo de validade e reprodutibilidade no Brasil. *Rev Bras Ativ Saude* 2001;6(2):5-18.
20. Van der Pligt P, Ball K, Hesketh KD, Teychenne M, Crawford D, Morgan PJ, et al. A pilot intervention to reduce postpartum weight retention and central adiposity in first-time mothers: results from the mums OnLINE (Online, Lifestyle, Nutrition & Exercise) study. *J Hum Nutr Diet* 2018;31(3):314-28.
21. Phelan S, Hagobian T, Brannen A, Hatley KE, Schaffner A, Muñoz-Christian K, et al. Effect of an internet-based program on weight loss for low-income postpartum women. *JAMA* 2017;317(23):2381-91.
22. Molina MC, Benseñor IM, Cardoso LO, Velásquez-Meléndez G, Drehmer M, Pereira TS, et al. Reproducibility and relative validity of the Food Frequency Questionnaire used in the ELSA-Brasil. *Cad Saude Publica* 2013;29(2):379-89.
23. Costa Louzada ML, Martins AP, Canella DS, Baraldi LG, Levy RB, Claro RM, et al. Ultra-processed foods and the nutritional dietary profile in Brazil. *Rev Saude Publica* 2015;49:38.
24. Brazilian Institute of Geography and Statistics. *Pesquisa de orçamentos familiares 2008-2009: tabelas de composição nutricional dos alimentos consumidos no Brasil*. Rio de Janeiro: Brazilian Institute of Geography and Statistics; 2011.
25. De Paula TP, Steemburgo T, De Almeida JC, Dall'Alba V, Gross JL, De Azevedo MJ. The role of Dietary Approaches to Stop Hypertension (DASH) diet food groups in blood pressure in type 2 diabetes. *Br J Nutr* 2012;108(1):155-62.
26. Ministry of Health of Brazil. *Orientações para a coleta e análise de dados antropométricos em serviços de saúde: norma técnica do sistema de vigilância alimentar e nutricional – SISVAN*. Brasília: Ministry of Health; 2011.
27. Stendell-Hollis NR, Thompson PA, West JL, Wertheim BC, Thomson CA. A comparison of Mediterranean-style and MyPyramid diets on weight loss and inflammatory biomarkers in postpartum breastfeeding women. *J Womens Health (Larchmt)* 2013;22(1):48-57.
28. Althuisen E, Van Poppel MNM, De Vries JH, Seidell JC, Van Mechelen W. Postpartum behaviour as predictor of weight change from before pregnancy to one year postpartum. *BMC Public Health* 2011;11:165.
29. Martins APB, Benicio MHDA. Influence of dietary intake during gestation on postpartum weight retention. *Rev Saude Publica* 2011;45(5):870-7.
30. Drehmer M, Camey SA, Nunes MA, Duncan BB, Lacerda M, Pinheiro AP, et al. Fibre intake and evolution of BMI: from pre-pregnancy to postpartum. *Public Health Nutr* 2013;16(8):1403-13.
31. Oliveira TC, Czeresnia D, Vargas EP, Barros DC. Concepções sobre práticas alimentares em mulheres de camadas populares no Rio de Janeiro, RJ, Brasil: transformações e ressignificações. *Interface (Botucatu)* 2018;22(65):435-46.
32. Baião MR, Deslandes SF. Práticas alimentares na gravidez: um estudo com gestantes e puérperas de um complexo de favelas do Rio de Janeiro (RJ, Brasil). *Ciênc Saúde Colet* 2010;15(Supl. 2):3199-206.
33. Baião MR, Santos MMAS, Libera BD, Machado RCM. Puerperium and its sociocultural dimension from the perspective of women living in Manguinhos - Rio de Janeiro, RJ, Brazil. *Demetra* 2013;8(Suppl. 1):309-20.
34. Schmitt NM, Nicholson WK, Schmitt J. The association of pregnancy and the development of obesity - Results of a systematic review and meta-analysis on the natural history of postpartum weight retention. *Int J Obes* 2007;31:1642-51.
35. Christenson A, Johansson E, Reynisdóttir S, Torgerson J, Hemmingsson E. Women's perceived reasons for their excessive postpartum weight retention: a qualitative interview study. *PLoS One* 2016;11(12):e0167731.
36. Boghossian NS, Yeung EH, Lipsky LM, Poon AK, Albert PS. Dietary patterns in association with postpartum weight retention. *Am J Clin Nutr* 2013;97(6):1338-45.