



Trabajo Original

Pediatría

The association between children's food habits, anthropometric parameters and health-related quality of life in Chilean school-age children

Asociación entre los hábitos alimentarios de los niños, los parámetros antropométricos y la calidad de vida relacionada con la salud en escolares chilenos

Pedro Delgado-Floody¹, Felipe Caamaño-Navarrete², Daniel Jerez-Mayorga³, Alfonso Cofré-Lizama^{4,5} and Iris Paola Guzmán-Guzmán⁶

¹Department of Physical Education, Sport and Recreation. Universidad de La Frontera. Temuco, Chile. ²Faculty of Education. Universidad Católica de Temuco. Temuco, Chile. ³Faculty of Rehabilitation Sciences. Universidad Andres Bello. Santiago de Chile, Chile. ⁴School of Psychology. Faculty of Social Sciences. Universidad Santo Tomás. Temuco, Chile. ⁵Universidad Mayor. Santiago de Chile, Chile. ⁶Faculty of Chemical-Biological Sciences. Universidad Autónoma de Guerrero. Guerrero, México

Abstract

Introduction: children's food habits have an influence on physical and mental health and should be monitored.

Objective: the purpose of this study was to determinate the association between children's food habits (i.e., Mediterranean diet adherence), anthropometric parameters and health-related quality of life (HRQoL) in Chilean schoolchildren.

Material and methods: girls (n = 282, 11.86 ± 0.82 years) and boys (n = 352, 12.02 ± 0.87 years) enrolled in public schools were included in this study (n = 634). Body mass index (BMI), waist circumference (WC), waist-to-height ratio (WtHR), body fat (BF), nutritional level and HRQoL were evaluated.

Results: the schoolchildren with high nutritional levels reported higher HRQoL (p = 0.018) and presented lower BMI, WC and WtHR (p < 0.001). Likewise, according to the following questions: Have your parent(s) treated you fairly? and Have you felt full of energy? The schoolchildren with high nutritional levels reported the major proportion in very and extremely answers (p < 0.001). The BMI was negatively correlated with HRQoL (r = -0.33, p < 0.05). Beside, children's food habits had significant correlation with HRQoL (r = 0.48, p = 0.002). The BMI (B = -0.41, 95% CI = -0.55, 0.93, p = 0.001) and children's food habits (B = -0.78, 95% CI = -0.09, -0.02, p = 0.002) reported associations with HRQoL.

Conclusion: in conclusion, the nutritional level (i.e., Mediterranean diet adherence) and BMI presented associations with HRQoL in Chilean schoolchildren; therefore, it is important to consider these results and develop different strategies in schoolchildren to improve their nutritional levels, as HRQoL represents a measure of happiness or satisfaction with life.

Key words:

Quality of life.
Schoolchildren.
Nutritional level.

Resumen

Introducción: los hábitos alimentarios de los niños influyen en su salud física y mental, por lo tanto, deben controlarse.

Objetivo: el propósito de este estudio fue determinar la asociación entre los hábitos alimentarios de los niños (adherencia a la dieta mediterránea), los parámetros antropométricos y la calidad de vida relacionada con la salud (CVRS) en escolares chilenos.

Material y métodos: se incluyeron en este estudio (n = 634) niñas (n = 282, 11,86 ± 0,82 años) y niños (n = 352, 12,02 ± 0,87 años) inscritos en escuelas públicas. Se evaluaron el índice de masa corporal (IMC), la circunferencia de la cintura (CC), la relación cintura-altura (RCE), la grasa corporal (GC), el nivel nutricional y la CVRS.

Resultados: los escolares con niveles nutricionales altos tuvieron una CVRS más alta (p = 0,018) y presentaron un IMC, CC y RCE más bajos (p < 0,001). Del mismo modo, de acuerdo a las preguntas "¿te han tratado tus padres de manera justa?" y "¿te has sentido lleno de energía?", los escolares con altos niveles nutricionales obtuvieron la mayor proporción en respuestas muy y muy altas (p < 0,001). El IMC se correlacionó negativamente con la CVRS (r = -0,33, p < 0,05). Además, los hábitos alimenticios de los niños tuvieron una correlación significativa con la CVRS (r = 0,48, p = 0,002). El IMC (B = -0,41; IC 95% = -0,55, 0,93, p = 0,001) y los hábitos alimentarios de los niños (B = -0,78, IC 95% = -0,09, -0,02, p = 0,002) reportaron asociaciones con la CVRS.

Conclusión: en conclusión, el nivel nutricional (es decir, la adherencia a la dieta mediterránea) y el IMC presentaron asociaciones con la CVRS en escolares chilenos; por lo tanto, es importante tener en cuenta estos resultados y desarrollar diferentes estrategias en los escolares para mejorar sus niveles nutricionales, ya que la CVRS representa una medida de felicidad o satisfacción con la vida.

Palabras clave:

Calidad de vida.
Escolares. Nivel
nutricional.

Received: 26/04/2019 • Accepted: 27/05/2019

Delgado-Floody P, Caamaño-Navarrete F, Jerez-Mayorga D, Cofré-Lizama A, Guzmán-Guzmán IP. The association between children's food habits, anthropometric parameters and health-related quality of life in Chilean school-age children. Nutr Hosp 2019;36(4):1061-1066

DOI: <http://dx.doi.org/10.20960/nh.02643>

Correspondence:

Pedro Delgado-Floody. Department Of Physical Education, Sport and Recreation. Universidad de La Frontera. Francisco Salazar 1145. Temuco, Chile
e-mail: pedro.delgado@ufrontera.cl

INTRODUCTION

Health-related quality of life (HRQoL) is a broad term generally representing a measure of happiness or satisfaction with life (1) and includes social, mental and physical well-being (2) as well as other factors such as purpose, fulfilment, motivation, social engagement, role, and participation in daily activities (3). In recent years, HRQoL has aroused the interest of the public health community as a relevant measure related to childhood obesity and lifestyle (4,5).

In contemporary society, food habits are one of the most frequent human behaviours, and healthy food habits generally have a strong participative role in the overall state of well-being (6). Science has reported that adherence to the Mediterranean diet is associated with positive effects on specific components of HRQoL and subjective happiness (7). Besides, it is also associated with physical and mental health status as well as HRQoL. Likewise, children's food habits have an influence on health in later life, present associations with different factors of mental wellness (8), and can be protective factors for subjective well-being in children (9). Therefore, children's food habits should be monitored.

Childhood obesity is a multifactorial health condition and is associated with psychosocial alterations, including deficiencies in social co-existence, with consequences for HRQoL (4,5). It has been observed that obese children tend to have affective problems, which may negatively affect their academic performance (10). Childhood obesity has come up as a threat to the physical as well as the mental health of children and adolescents (11). Therefore, the relationship between mental illness (i.e., HRQoL), and obesity has been increasingly recognised in the paediatric population. However, research exploring the association between nutritional level and HRQoL in children has been inconsistent. Therefore, the purpose of this study was to determinate the association between children's food habits (i.e., Mediterranean diet adherence), weight status and HRQoL in Chilean schoolchildren.

MATERIAL AND METHOD

This cross-sectional study included 634 schoolchildren, girls ($n = 282$, 11.86 ± 0.82 years) and boys ($n = 352$, 12.02 ± 0.87 years) from four primary public schools in Chile selected by convenience. The sample size is similar to previous studies (12,13). Parents and guardians were informed about the study and provided signed written consent for participation. Additionally, all children gave their written assent on the day of the assessment.

The inclusion criteria were: a) presenting informed consent of the parents and the assent of the participant; b) belong to educational centres, and c) be between 10 and 13 years of age. The exclusion criteria were having a musculoskeletal disorder or any other known medical condition, which might alter the participant's health and physical activity levels. Moreover, schoolchildren with physical, sensorial or intellectual disabilities were excluded.

The investigation complied with the Declaration of Helsinki (2013) and was approved by the Ethics Committee of Universidad de La Frontera, Chile (DFP16-0013). The tests were explained to all the participants before the study began.

MATERIAL

ANTHROPOMETRIC ASSESSMENT

The body mass index (BMI), calculated as the body mass divided by the square of the height in metres (kg/m^2), was used to estimate the degree of obesity. The body mass (kg) was measured using a TANITA scale, model Scale Plus UM – 028 (Tokyo, Japan); the children were weighed in their underclothes, without shoes, and the height (m) was estimated with a Seca® stadiometer, model 214 (Hamburg, Germany), graduated in mm. The BMI is shown in the growth table of the Centers for Disease Control and Prevention (CDC), Overweight and Obesity, verifying the corresponding age and the sex-related percentile. Child obesity is defined as a BMI equal to or greater than the 95th percentile and overweight as a BMI equal to or greater than the 85th percentile among children of the same age and sex (14). The waist circumference (WC) was measured using a Seca® tape measure model 201 (Hamburg, Germany) at the height of the umbilical scar (15). The waist-to-height ratio (WtHR) was obtained by dividing the WC by the height and was used as a tool for estimating the accumulation of fat in the central zone of the body following international standards (16). To measure the % body fat (BF), we used the tricipital fold and the subscapular fold (Lange Skinfold Caliper, 102-602L, Minneapolis, USA) applying Slaughter's formula (17) ♀ ♂ : Girls: $\% \text{BF} = 1.33 (\text{tricipital} + \text{subscapular}) - 0.013 (\text{tricipital} + \text{subscapular})^2 - 2.5$; Boys: $\% \text{BF} = 1.21 (\text{tricipital} + \text{subscapular}) - 0.008 (\text{tricipital} + \text{subscapular})^2 - 1.7$.

The research assistant was submitted to the test-retest ($n = 64$, 10% of study sample) protocol to verify its technical measurement error with an intra-class correlation coefficient (ICC), in WC (ICC = 0.94), tricipital fold (ICC = 0.91) and subscapular fold (ICC = 0.91).

NUTRITIONAL LEVEL ACCORDING CHILDREN'S FOOD HABITS

The children's food habits were assessed by the Krece Plus test (18), which is a tool to assess the eating pattern and the relationship with the nutritional status based on the Mediterranean diet. The questionnaire has 15 items and the format assesses a set of items about the food consumed in the diet. Each item has a score of +1 or -1, depending on whether it approximates the ideal of the Mediterranean diet. The total points are added, and according to the score the nutritional status is classified as follows:

1. Low nutritional level: Less than or equal to 5.
2. Moderate nutritional level: From 6 to 8.
3. High nutritional level: Greater than or equal to 9.

QUALITY OF LIFE

The HRQoL for Children and Young People was measured with the KIDSCREEN-10 questionnaire (19). The HRQoL was assessed using the Spanish version. This instrument of self-report consists of 10 items derived from the version of the KIDSCREEN-27 questionnaire. It is answered using a Likert scale with 5 possibilities (not at all, slightly, moderately, very often, always). The scale was reset to have a score from 0 to 10.

STATISTICAL ANALYSIS

Statistical analysis was performed using SPSS v23.0 software (SPSS™ IBM Corporation, NY, USA). Normal distribution was tested using the Kolmogorov-Smirnov test. For continuous variables, values are presented as mean and standard deviation, whereas for categorical variables data are presented as proportions. Differences between weight status and nutritional level were determined using the one-way ANOVA. To determine the association between HRQoL with nutritional level and anthropometric parameters, a multivariable logistic regression was used. The chi-squared test was applied to compare proportions according to weight status and nutritional level about different questions of quality of life. Values of $p < 0.05$ were considered statistically significant.

RESULTS

Table I shows the results according to nutritional level, which considered children's food habits (i.e., Mediterranean diet adherence). The schoolchildren with high nutritional levels reported higher quality of life ($p = 0.018$) and presented lower BMI, WC and WtHR ($p < 0.001$).

The distribution of the response (significantly different) between groups according to nutritional level in the categories of the KID-

SCREEN-10 are presented in the table II. In relationship to the following questions: Have your parent(s) treated you fairly? and Have you felt full of energy? The schoolchildren with high nutritional levels reported the major proportion in very and extremely answers ($p < 0.001$). Instead, in relationship to the question: Have you felt sad? The schoolchildren with low nutritional levels presented the major proportion in very and extremely answers ($p < 0.001$).

Linear correlations (adjusted by sex) indicated that BMI was negatively correlated with food habits ($r = 0.29, p < 0.0001$) and HRQoL ($r = -0.33, p < 0.05$). Besides, children's food habits had a significant correlation with HRQoL ($r = 0.48, p = 0.002$) (Table III).

The multivariable logistic regression reported associations between HRQoL, BMI ($B = -0.41, 95\% \text{ CI} = -0.55, 0.93, p = 0.001$), and food habits ($B = -0.78, 95\% \text{ CI} = -0.09, -0.02, p = 0.002$) (Table IV).

DISCUSSION

The purpose of this study was to determinate the association between food habits (i.e., Mediterranean diet adherence), anthropometric parameters and HRQoL in Chilean schoolchildren. The main findings were: a) schoolchildren with high nutritional levels presented higher quality life than low and moderate nutritional level groups; and (b) BMI and food habits (i.e., Mediterranean diet adherence) reported associations with quality of life.

The schoolchildren with high nutritional levels reported higher HRQoL and they felt full of energy in comparison with moderate and low nutritional level groups. Instead, schoolchildren with low nutritional levels felt sad in greater proportion. A previous study has reported that food habits, such as adherence to the Mediterranean diet, presented positive correlations with different factors of mental wellness (8). Likewise, another investigation has reported that Japanese children with undesirable lifestyles, such as skipping breakfast, were more likely to have poor HRQoL, independent of sex, BMI, social background, and somatic symptoms (20).

Table I. Characteristics of sample study according nutritional level

	Low nutritional level (n = 222) A	Moderate nutritional level (n = 301) B	High Nutritional level (n = 111) C	p value	Post hoc
Age (years)	11.86 ± 0.83	11.97 ± 0.86	12.08 ± 0.88	$p = 0.082$	-
Body mass (kg)	54.05 ± 14.45	52.87 ± 14.54	47.14 ± 11.31	$p < 0.001$	A > B > C
Size (m)	1.54 ± 0.10	1.56 ± 0.11	1.54 ± 0.11	$p = 0.097$	-
BMI (kg/m ²)	22.58 ± 4.46	21.58 ± 4.67	19.71 ± 4.02	$p < 0.001$	A > B > C
WC (cm)	75.55 ± 12.53	73.32 ± 11.03	69.57 ± 9.51	$p < 0.001$	A > B > C
WtHR (WC / height)	0.49 ± 0.08	0.47 ± 0.07	0.45 ± 0.06	$p < 0.001$	A > B > C
BF (%)	25.20 ± 6.66	24.21 ± 7.34	23.37 ± 8.07	$p = 0.077$	-
Quality of life (score)	37.00 ± 4.59	37.23 ± 4.89	38.52 ± 4.72	$p = 0.018$	A, B < C

Data are presented as mean and SD. $p < 0.05$ considered statistically significant. BMI: body mass index; WC: waist circumference; WtHR: waist-to-height ratio; BF: body fat. (A) low nutritional level, (B) denotes moderate nutritional level and (C) high nutritional level in post hoc analysis.

Table II. Answer in relation with quality of life according nutritional level

	Low nutritional level (n = 222)		Moderate nutritional level (n = 301)		High nutritional level (n = 111)		p value
Have you felt fit and well?							p = 0.006
Not at all	0	0.0%	4	1.3%	0	0.0%	
Slightly	6	2.7%	9	3.0%	5	4.5%	
Moderately	63	28.4%	113	37.5%	37	33.3%	
Very	122	55.0%	131	43.5%	41	36.9%	
Extremely	31	14.0%	44	14.6%	28	25.2%	
Have you felt full of energy?							p = 0.010
Slightly	0	0.0%	10	3.3%	0	0.0%	
Moderately	47	21.2%	76	25.2%	18	16.2%	
Very	130	58.6%	154	51.2%	70	63.1%	
Extremely	45	20.3%	61	20.3%	23	20.7%	
Have you felt sad?							p < 0.001
Not at all	20	9.0%	30	10.0%	13	11.7%	
Slightly	93	41.9%	138	45.8%	61	55.0%	
Moderately	69	31.1%	112	37.2%	37	33.3%	
Very	37	16.7%	14	4.7%	0	0.0%	
Extremely	3	1.4%	7	2.3%	0	0.0%	
Have you felt lonely?							p < 0.001
Not at all	48	21.6%	89	29.6%	24	21.6%	
Slightly	83	37.4%	109	36.2%	46	41.4%	
Moderately	38	17.1%	86	28.6%	30	27.0%	
Very	44	19.8%	9	3.0%	2	1.8%	
Extremely	9	4.1%	8	2.7%	9	8.1%	
Have you had enough time for yourself?							p < 0.001
Slightly	18	8.1%	6	2.0%	9	8.1%	
Moderately	58	26.1%	56	18.6%	16	14.4%	
Very	95	42.8%	145	48.2%	38	34.2%	
Extremely	51	23.0%	94	31.2%	48	43.2%	
Have you been able to do the things that you want to do in your free time?							p < 0.001
Slightly	42	18.9%	11	3.7%	0	0.0%	
Moderately	43	19.4%	56	18.6%	26	23.4%	
Very	76	34.2%	151	50.2%	45	40.5%	
Extremely	61	27.5%	83	27.6%	40	36.0%	
Have your parent(s) treated you fairly?							p < 0.001
Not at all	1	.5%	1	.3%	0	0.0%	
Slightly	2	.9%	8	2.7%	3	2.7%	
Moderately	60	27.0%	47	15.6%	6	5.4%	
Very	59	26.6%	92	30.6%	63	56.8%	
Extremely	100	45.0%	153	50.8%	39	35.1%	

The data shown represent n (%). p values < 0.05 were considered statistically significant.

Table III. Linear correlation adjusts by sex between anthropometrics parameters and food habits and quality of life

	Children's foods habits		Quality of life	
BMI	-0.29	(p < 0.001)	-0.33	(p < 0.001)
WC	-0.16	(p < 0.001)	-0.20	(p < 0.001)
WtHR	-0.19	(p < 0.001)	-0.19	(p < 0.001)
BF	-0.08	(p = 0.044)	-0.03	(p = 0.398)
Children's foods habits		-	0.48	(p = 0.002)

The data shown represent r (p values). p < 0.05 were considered statistically significant. BMI: body max index; WC: waist circumference; WtHR: waist-to-height ratio; BF: body fat.

Likewise, a study that evaluated the adherence to the Mediterranean diet with HRQoL in Portuguese adolescents reported that the adherence was positively associated with HRQoL (21). Moreover, a recent study carried out in Spanish children and adolescents concluded that adherence to the Mediterranean diet was found to behave as a protective factor for positive well-being in a cross-sectional analysis (22). A recent systematic review and meta-analysis reported that unhealthy dietary behaviour or lower diet quality was associated with decreased HRQoL among children and adolescents (23).

On the other hand, schoolchildren with high nutritional levels presented better anthropometric parameters. A previous study has reported that regular breakfast consumption is significantly associated with lower body fatness and healthier dietary habits (24). In the same line, a study carried out in Chilean schoolchildren reported that adherence to the Mediterranean diet was positively associated with a healthier lifestyle, mental wellness and physical fitness (8). Another study that examined the association between breakfast consumption and cardiovascular risk factors in European adolescents showed that adolescents who regularly consumed breakfast had lower body fat content, higher cardiorespiratory fitness and a healthier cardiovascular profile (25). Similarly, a study that evaluated the associations between habitual school day breakfast consumption, BMI, physical activity and cardiorespiratory fitness in children reported that habitual breakfast consumption was associated with healthy BMI and higher PA levels in schoolchildren (26).

In our sample, the multivariable logistic regression reported a negative association between BMI and HRQoL. In this line, a study that evaluated the relationships between HRQoL and BMI in children reported that physical, social and school functioning was significantly lower for obese children when compared with normal weight children (27). Likewise, a study showed that HRQoL was inversely related to the extent of obesity measured as either BMI or BF in children (28). Moreover, another investigation reported

Table IV. Association of quality of life with sex, anthropometrics parameters and children's food habits

	B	(95% CI)		p value
(Constant)	42.047	38.90,	45.186	p < 0.000
Sex	0.187	-0.55,	.929	p = 0.621
BMI	-0.41	-0.34,	-.083	p < 0.001
WC	-0.02	-.097,	.056	p = 0.601
WtHR	-0.81	-12.497,	10.871	p = 0.891
BF	0.04	-.013,	.096	p = 0.134
Children's food habits	0.78	-0.08,	0.24	p < 0.001

The data shown represent B, 95% CI and p value. p < 0.05 were considered statistically significant. BMI: body max index; WC: waist circumference; WtHR: waist-to-height ratio; BF: body fat.

that the BMI was inversely correlated to adolescent self-reported HRQoL compared with normal weight adolescents (29). However, an investigation in elementary school girls reported no significant association between HRQoL and being overweight (30). In this line, BMI was not found to be a strong explanatory factor for variations in HRQoL in a Norwegian school sample; however, BMI was negatively associated with the HRQoL subscales of self-perception and physical well-being (31). A study examining HRQoL across weight categories in adolescents using both a general and a condition-specific measure reported that differences in HRQoL across weight categories differed based on the measure utilized. Clinicians should use caution when interpreting and comparing HRQoL findings (32). Childhood obesity can profoundly affect children's physical, social, and emotional well-being as well as self-esteem. It is also associated with a lower quality of life experienced by the child (33).

LIMITATIONS

The main limitation of this study is its cross-sectional design. As a result, causal relationships cannot be established. Despite this limitation, it is noteworthy that the information available regarding the association between HRQoL and children's food habits is important, especially for professionals in the Nutrition Sciences. So, the current study provides some insights into this field.

CONCLUSION

In conclusion, children's food habits (i.e., Mediterranean diet adherence) and BMI were associated with HRQoL in Chilean schoolchildren; therefore, it is important to consider these results and develop different strategies in schoolchildren to improve their nutritional levels, as HRQoL represents a measure of happiness or satisfaction with life.

REFERENCES

- Auld G, Baker S, Hernández-Garbanzo Y, et al. The Expanded Food and Nutrition Education Program's Impact on Graduates' Quality of Life. *J Nutr Educ Behav* 2018;51(2):217-23.
- Wake M, Hughes EK, Collins CM, et al. Parent-reported health-related quality of life in children with congenital hearing loss: A population study. *Ambul Pediatr* 2004;4(5):411-7.
- Watson P, Zhang J-P, Rizvi A, et al. A meta-analysis of factors associated with quality of life in first episode psychosis. *Schizophr Res* 2018;202:26-36.
- Walker L, Hill A. Obesity: The role of child mental health services. *Child and Adolescent Mental Health* 2009;14(3):114-20.
- Gálvez Casas A, Rosa Guillamón A, García-Cantó E, et al. Estado nutricional y calidad de vida relacionada con la salud en escolares del sureste español. *Nutr Hosp* 2015;31:737-43.
- Bundhun D, Rampadarath S, Puchooa D, et al. Dietary intake and lifestyle behaviors of children in Mauritius. *Heliyon* 2018;4(2):e00546.
- Ferrer-Cascales R, Albaladejo-Blázquez N, Ruiz-Robledillo N, et al. Higher Adherence to the Mediterranean Diet is Related to More Subjective Happiness in Adolescents: The Role of Health-Related Quality of Life. *Nutrients* 2019;11(3):698.
- Muros JJ, Cofre-Bolados C, Arriscado D, et al. Mediterranean diet adherence is associated with lifestyle, physical fitness, and mental wellness among 10-y-olds in Chile. *Nutrition* 2017;35:87-92.
- Lindberg L, Swanberg I. Well-being of 12-year-old children related to interpersonal relations, health habits and mental distress. *Scand J Caring Sci* 2006;20(3):274-81.
- Barmola K. Family Environment, Mental Health and Academic Performance of Adolescents. *Sociology* 2013;2(12).
- Sagar R, Gupta T. Psychological aspects of obesity in children and adolescents. *Indian J Pediatr* 2018;85(7):554-9.
- Delgado-Floody P, Cofré A, Guzmán I, et al. Perception of obese schoolchildren regarding their participation in the Physical Education class and their level of self-esteem: comparison according to corporal status. *Nutr Hosp* 2018;35(6):1270-4.
- Delgado-Floody P, Caamaño-Navarrete F, Palomino-Devia C, et al. Relationship in obese Chilean schoolchildren between physical fitness, physical activity levels and cardiovascular risk factors. *Nutr Hosp* 2019;36(1):13-9.
- Centers for Disease Control and Prevention, Overweight and obesity. Defining overweight and obesity; 2010.
- Schröder H, Ribas L, Koebnick C, et al. Prevalence of abdominal obesity in Spanish children and adolescents. Do we need waist circumference measurements in pediatric practice? *PLoS One* 2014;9(1):e87549.
- Chung IH, Park S, Park MJ, et al. Waist-to-Height Ratio as an Index for Cardiometabolic Risk in Adolescents: Results from the 1998-2008 KNHANES. *Yonsei Med J* 2016;57(3):658-63.
- Slaughter MH, Lohman TG, Boileau RA, et al. Skinfold equations for estimation of body fatness in children and youth. *Hum Biol* 1988;60(5):709-23.
- Majem LS, Barba LR, Bartrina JA, et al. Obesidad infantil y juvenil en España. Resultados del Estudio enKid (1998-2000). *Med Clin* 2003;121(19):725-32.
- Ravens-Sieberer U, Erhart M, Rajmil L, et al. Reliability, construct and criterion validity of the KIDSCREEN-10 score: a short measure for children and adolescents' well-being and health-related quality of life. *Qual Life Res* 2010;19(10):1487-500.
- Chen X, Sekine M, Hamanishi S, et al. Lifestyles and health-related quality of life in Japanese school children: a cross-sectional study. *Prev Med* 2005;40(6):668-78.
- Evaristo OS, Moreira C, Lopes L, et al. Associations between physical fitness and adherence to the Mediterranean diet with health-related quality of life in adolescents: results from the LabMed Physical Activity Study. *Eur J Public Health* 2018;28(4):631-5.
- Esteban-Gonzalo L, Turner AI, Torres SJ, et al. Diet quality and well-being in children and adolescents: the UP&DOWN longitudinal study. *Br J Nutr* 2019;121(2):221-31.
- Wu XY, Zhuang LH, Li W, et al. The influence of diet quality and dietary behavior on health-related quality of life in the general population of children and adolescents: a systematic review and meta-analysis. *Qual Life Res* 2019;28(8):1989-2015.
- Ahadi Z, Qorbani M, Kelishadi R, et al. Association between breakfast intake with anthropometric measurements, blood pressure and food consumption behaviors among Iranian children and adolescents: the CASPIAN-IV study. *Public Health* 2015;129(6):740-7.
- Hallström L, Labayen I, Ruiz JR, et al. Breakfast consumption and CVD risk factors in European adolescents: the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) Study. *Public Health Nutr* 2013;16(7):1296-305.
- Sandercock GR, Voss C, Dye L. Associations between habitual school-day breakfast consumption, body mass index, physical activity and cardiorespiratory fitness in English schoolchildren. *Eur J Clin Nutr* 2010;64(10):1086.
- Khodaverdi F, Alhani F, Kazemnejad A, et al. The relationship between obesity and quality of life in school children. *Iran J Public Health* 2011;40(2):96.
- Morrison KM, Shin S, Tarnopolsky M, et al. Association of depression & health related quality of life with body composition in children and youth with obesity. *J Affect Disord* 2015;172:18-23.
- Jalali-Farahani S, Chin Y, Amiri P, et al. Body mass index (BMI)-for-age and health-related quality of life (HRQoL) among high school students in Tehran. *Child Care Health Dev* 2014;40(5):731-9.
- Jalali-Farahani S, Alamdari S, Karimi M, et al. Is overweight associated with health-related quality of life (HRQoL) among Tehranian school children? *SpringerPlus* 2016;5(1):313.
- Helseth S, Haraldstad K, Christophersen K-A. A cross-sectional study of Health Related Quality of Life and body mass index in a Norwegian school sample (8-18 years): a comparison of child and parent perspectives. *Health Qual Life Outcomes* 2015;13(1):47.
- Dalton AG, Smith C, Dalton III WT, et al. Examining general versus condition-specific health-related quality of life across weight categories in an adolescent sample. *J Pediatr Health Care* 2015;29(5):453-62.
- Sahoo K, Sahoo B, Choudhury AK, et al. Childhood obesity: causes and consequences. *J Family Med Prim Care*. 2015;4(2):187-92.