



Trabajo Original

Epidemiología y dietética

Vitamin D deficiency in primary health care users at risk in Spain *Deficiencia de vitamina D en usuarios de atención primaria con riesgo en España*

Andrés Díaz-López^{1,2,3}, Indira Paz-Graniel^{1,2}, Ramón Alonso-Sanz⁴, Carolina Marqués Baldero⁵, César Mateos Gil², and Victoria Arijá⁷

¹Facultad de Medicina y Ciencias de la Salud. Universitat Rovira i Virgili (URV). Reus, Tarragona. Spain. ²Institut d'Investigació Sanitària Pere Virgili (IISPV). Reus, Tarragona. Spain. ³Centro de Investigación Biomédica en Red, Fisiopatología de la Obesidad y la Nutrición (CIBEROBN). Instituto de Salud Carlos III. Madrid, Spain. ⁴Centro de Salud de Alfaro. Alfaro, La Rioja. Spain. ⁵Hospital Reina Sofía. Tudela, Navarra. Spain. ⁶Research Group in Nutrition and Mental Health (NUTRISAM). Universitat Rovira i Virgili (URV). Reus, Tarragona. Spain. ⁷Tarragona-Reus Research Support Unit. Institut Universitari d'Investigació en Atenció Primària Jordi Gol (IDIAP Jordi Gol). Barcelona, Spain

Abstract

Background and objectives: despite long hours of sunlight in Spain, vitamin D deficiency has been rising of late. We aimed to describe vitamin D deficiency and insufficiency in at-risk patients in La Rioja, a region in northern Spain.

Methods: a retrospective, cross-sectional study involving 21,490 patients (74.3 % women) aged 14-105 years, who were primary health care (PHC) users in La Rioja, with an available 25-hydroxyvitamin D (25(OH)D) test result corresponding to the first request from January 2013 to October 2015. Vitamin D deficiency, insufficiency, and sufficiency were defined as 25(OH)D levels < 12 ng/mL, 12-20 ng/mL, and ≥ 20 ng/mL, respectively. Hypovitaminosis D was defined considering 2 cut-off points for 25(OH)D: < 20 ng/mL and < 30 ng/mL.

Results: mean 25(OH)D levels were 18.3 (SD, 11.6) ng/mL in the whole sample, being lower in men than in women (17.6 vs 18.5 ng/mL, $p < 0.001$). Significantly higher 25(OH)D levels were found in the groups with 55-59, 60-64, 65-69, and 70-74 years of age (range, 19.4-20.4 ng/mL), and lower levels were found in the groups with 80-85 years and older (16.7 and 14.5 ng/mL), as compared to other groups. Two thirds of the population were vitamin D-deficient (< 12 ng/mL, 30.9 %) or insufficient (12-20 ng/mL, 32.8 %), and only 32.8 % of patients were found to be vitamin D-sufficient (> 20 ng/ml). Hypovitaminosis D rate was higher (89.7 %) when the 25(OH)D cut-off point was set at < 30 ng/mL. A higher hypovitaminosis D rate (< 20 ng/mL) was found in men (67.1 %) vs women (62.6 %) ($p < 0.001$), and it was more prevalent in patients over 75 years (72.6 %). The prevalence of hypovitaminosis D followed a seasonal pattern ranging from 51.8 % to 76.5 %; the lowest 25(OH)D levels were found in winter, and the highest in summer, with levels above 20 ng/mL in all age groups except for patients over 75 years (17.7 ng/mL). Patients over 75 years exhibited high rates of deficiency or insufficiency across all seasons of the year.

Conclusions: in La Rioja, Spain, hypovitaminosis D was highly prevalent among PHC users, and especially in older age groups and during the winter months.

Keywords:

Vitamin D deficiency.
Hypovitaminosis D.
Primary care. La Rioja.

Received: 05/02/2021 • Accepted: 13/03/2021

Acknowledgments: this research is based on data and information compiled and provided by San Pedro Hospital, Logroño (Spain). The analyses, conclusions, opinions, and statements expressed herein are solely those of the authors and do not reflect those of the data sources; no endorsement is intended or should be inferred.

Funding: this work declares no funding sources.

Disclosure of interests: the authors declare that they have no conflict of interest.

Authors' contributions: study concept and design: AD-L and VA. Statistical analyses: AD-L and VA. Drafting of the manuscript: AD-L, IP-G, and VA. All authors reviewed the manuscript for important intellectual content, and approved the final version to be published. AD-L and VA are the guarantors of this work and, as such, had full access to all the data in the study, and take responsibility for the integrity of the data and the accuracy of the data analysis.

Díaz-López A, Paz-Graniel I, Alonso-Sanz R, Marqués Baldero C, Mateos Gil C, Arijá V. Vitamin D deficiency in primary health care users at risk in Spain. *Nutr Hosp* 2021;38(5):1058-1067

DOI: <http://dx.doi.org/10.20960/nh.03565>

Correspondence:

Victoria Arijá. Unit of Preventive Medicine and Public Health. Faculty of Medicine and Health Science. Universitat Rovira i Virgili. C/Sant Llorenç 21. 43201 Reus, Tarragona. Spain
e-mail address: victoria.arija@urv.cat

Resumen

Justificación y objetivos: a pesar de las largas horas de exposición solar en España, la deficiencia de vitamina D ha ido en aumento. Nuestro objetivo fue describir la deficiencia e insuficiencia de vitamina D en pacientes de riesgo en La Rioja, España.

Métodos: estudio transversal retrospectivo en el que participaron 21.490 pacientes (74,3 % mujeres), usuarios de Atención Primaria (APS) en La Rioja, con resultado de prueba de 25-hidroxivitamina D (25(OH)D) disponible, correspondiente a las solicitudes efectuadas entre enero de 2013 y octubre de 2015. La deficiencia, insuficiencia y suficiencia de vitamina D se definieron como niveles de 25(OH)D < 12 ng/mL, = 12-20 ng/mL y \geq 20 ng/mL, respectivamente. La hipovitaminosis D se definió considerando 2 puntos de corte de 25(OH)D: < 20 ng/mL y < 30 ng/mL.

Resultados: los valores medios de 25(OH)D fueron de 18,3 (DE: 11,6) ng/mL en toda la muestra, siendo menores en los hombres que en las mujeres (17,6 vs. 18,5 ng/mL, $p < 0,001$). Se encontraron niveles de 25(OH)D significativamente más altos en los grupos de 55-59, 60-64, 65-69 y 70-74 años (rango, 19,4-20,4 ng/mL), y más bajos en los grupos de 80-85 años y mayores (16,7 y 14,5 ng/mL) en comparación con otros grupos. Dos tercios de la población presentaban deficiencia de vitamina D (< 12 ng/mL, 30,9 %) o insuficiencia (12-20 ng/mL, 32,8 %), y solo el 32,8 % de los pacientes presentaban niveles suficientes de vitamina D (> 20 ng/mL). La hipovitaminosis D fue mayor (89,7 %) cuando el punto de corte de 25(OH)D se estableció en < 30 ng/mL. Se encontró una mayor tasa de hipovitaminosis D (< 20 ng/mL) en los hombres (67,1 %) que en las mujeres (62,6 %) ($p < 0,001$), y dicha hipovitaminosis fue más prevalente en los pacientes mayores de 75 años (72,6 %). La prevalencia de la hipovitaminosis D siguió un patrón estacional que varió del 51,8 % al 76,5 %. Los niveles más bajos de 25(OH)D se encontraron en invierno y los más altos en verano, con niveles por encima de 20 ng/mL en todos los grupos de edad excepto en los pacientes mayores de 75 años (17,7 ng/mL). Los pacientes mayores de 75 años presentaron altas tasas de deficiencia o insuficiencia durante todas las estaciones del año.

Conclusiones: en La Rioja, España, la hipovitaminosis D fue altamente prevalente entre los usuarios de APS y, especialmente, en los grupos de mayor edad y durante los meses de invierno.

Palabras clave:

Deficiencia de vitamina D.
Hipovitaminosis D.
Atención primaria.
La Rioja.

INTRODUCTION

Vitamin D deficiency is an epidemic worldwide, turning into a major public health concern even in developed countries with sufficient sunshine and enough food availability (1-3). About 1 billion people worldwide suffer from this condition, while 30-50 % of the general population has vitamin D insufficiency (4-6). Although vitamin D is naturally present in a few foods, it is added to others, and consumed as a supplement, most of the requirements of vitamin D are covered by its synthesis through skin exposure to ultraviolet-B (UVB) radiation in sunlight (80-100 %) (4,7). So far, the best marker of vitamin D status is circulating 25-hydroxyvitamin D (25(OH)D) concentration, though no consensus exists on optimal serum 25(OH)D levels and the threshold for defining vitamin D insufficiency is a controversial subject (3,8).

Vitamin D is a prohormone indispensable for appropriate functioning in the human body. It is classically implicated in the regulation of calcium homeostasis and healthy bone growth, so its deficiency could cause several diseases during life (4). While, in children, vitamin D deficiency increases the risk of developing rickets and muscle-skeletal disruption, in adults is associated with osteomalacia, osteopenia, and osteoporosis, increasing the risk of fractures and falls (4,9). However, these clinically overt manifestations are only "the top of the vitamin D deficiency iceberg" (10,11). Nowadays, it is clear that vitamin D is also involved in a multitude of complex metabolic pathways with potential extraskeletal functions, and because of this, vitamin D deficiency may play other roles in a number of diseases (12). In fact, emerging research supports the possible role of vitamin D against cancer, infection, autoimmune disease, depression, diabetes, and poor outcomes in cardiovascular health (4,13-16). Although convincing, this evidence does not prove causality but supports a hypothesis for further study.

Vitamin D inadequacy not only is frequently observed in vulnerable populations with low consumption of vitamin D food sources and short exposure to sunlight (6), but also children and

younger and middle-aged adults, postmenopausal women, and the elderly are at equally high risk for insufficiency worldwide (1,2,6). This state of vitamin D hypovitaminosis is replicated in Spain, a region of southern Europe. Previous studies conducted in Spanish populations have observed that vitamin D intakes are lower than recommended (17,18). In addition, available data from a recent review of vitamin D status in Spain suggest the presence of widespread vitamin D deficiency and/or insufficiency in several population groups of all ages (19) (children (24.3 %) (20), young individuals (61 %) (21), primary health care users (87 %) (22), pregnant women (50.2 %) (23), postmenopausal osteoporotic women (44 %) (24), the elderly whether in the community (79-80 %) or institutionalized (91 %) (25,26)), ranging from 24.3 % to 91 %. Probably this wide variation in prevalence rates across studies partly relates to individual differences in the study population such as age or pre-existing risk factors, together with geographical and seasonal variances, and the 25(OH)D cutoff used to define vitamin D inadequacy.

Nonetheless, in Spain, despite being a country with many hours of sunlight, it seems that vitamin D deficiency has been on the rise. To our knowledge, studies on this matter conducted in La Rioja population are still lacking. Thus, we aimed to describe the prevalence of vitamin D deficiency and insufficiency in at-risk patients attending primary healthcare facilities in La Rioja, a northern Spanish region, according to age, gender, and seasonal variations. In addition, we also examined the prevalence of hypovitaminosis D considering 2 cutoff 25(OH)D levels available in the literature.

MATERIAL AND METHODS

STUDY DESIGN AND STUDY POPULATION

This was a retrospective cross-sectional, descriptive, population-based study that analyzed data from primary health care (PHC)

users in La Rioja, a region located in northern Spain (located 42 ° latitude N, 2 ° LW). The study population included 21,490 patients aged 15 to 105 years who regularly or occasionally attended the PHC clinic, with an available serum 25(OH)D measurement corresponding to the first requests from January 2013 to October 2015 (inclusion criteria).

Eligible individuals were users who for any medical reason or on their own initiative attended a primary health care center and underwent blood tests for serum vitamin D levels as requested by a doctor. While no consensus exists on precise indications for 25(OH)D testing in at-risk patients, in our study population vitamin D testing was carried out when high suspicion or diagnostic criteria for rickets or osteomalacia existed, or for any other clinical indication based on the professional clinical judgment of a clinician according to the Endocrine Society Clinical Practice Guideline (27). Consequently, from a clinical perspective, candidates for screening were considered patients at high risk for vitamin D deficiency, which therefore justified a 25(OH)D measurement. No other inclusion or exclusion criteria were defined. It should be noted that no individuals were receiving vitamin D supplementation.

The study was conducted in agreement with the 1990 Declaration of Helsinki and subsequent amendments. The use of data for the purpose of this research was authorized by the steering committee of Hospital San Pedro in Logroño, La Rioja, Spain. Since all data handled by the researchers were anonymous and no additional contact with patients was needed, this study did not require a review by a Research Ethics Board.

DATA COLLECTION

From the electronic medical records of the individuals selected we retrospectively collected the following data: age, gender, first serum 25(OH)D measurement, and date of the blood sample request.

The study population was divided into age groups of five years each. Only the first (14-19 years) and last (≥ 85 years) age groups were different in class interval width. Furthermore, the following 4 age groups were also arbitrarily considered in our analyses; 14-29, 30-54, 55-74, and ≥ 75 years. The season during which vitamin D testing took place was categorized into months (January to December), and seasonality by quarters of a year: winter (January to March), spring (April to June), summer (July to September), and autumn (October to December).

ASSESSMENT OF CIRCULATING 25(OH)D CONCENTRATION

Serum 25(OH)D levels were quantified by using a direct competitive chemiluminescent immunoassay method based on the DiaSorin® Liaison System XL, according to the manufacturer's instructions. All tests for serum vitamin D levels were conducted in the central laboratory of San Pedro Hospital, Logroño, which belongs to the Rioja Health Service (SERIS).

Vitamin D status was evaluated according to the different cut-off points indicated in the literature. Because there is no consensus regarding the 25(OH)D cutoff value for defining vitamin D inadequacy, for the purpose of this study the reference ranges used were based upon the recommendations of the US Institute of Medicine (USIOM) (28) and the US Endocrine Society (USES) (27). The USIOM defines vitamin D deficiency, insufficiency, and sufficiency as 25(OH)D concentrations of < 12 ng/mL (30 nmol/L), 12-20 ng/mL (30-50 nmol/L), and ≥ 20 ng/mL (50 nmol/L), respectively. The USES-recommended 25(OH)D cutoffs to define vitamin D deficiency, insufficiency, and sufficiency/adequacy were < 20 ng/mL, 20-30 ng/mL, and ≥ 30 ng/mL, respectively. Hypovitaminosis D was also defined considering 2 additional cutoff points for 25(OH)D (< 20 ng/mL and < 30 ng/mL).

STATISTICS

All descriptive and comparative analyses were performed using the STATA software, version 15.0 194 (StataCorp LP, Tx. USA). The results are expressed as mean (SD) for quantitative variables and number (%) for qualitative variables, unless otherwise indicated. Differences between groups were examined by an independent Student's t-test or by χ^2 or one-way ANOVA tests with Bonferroni's post hoc analysis, as appropriate. All tests were two-tailed, and p values less than 0.05 were considered statistically significant.

RESULTS

The study results are based on the data analysis of 21490 PHC users with a serum vitamin D test measurement corresponding to the first requested blood test from January 2013 to October 2015. Of these, 74.3 % ($n = 16,000$) were women. The mean ages of the women and men were 62 (range, 14-105) and 63 (range, 14-101) years, respectively. As shown in table I, the average 25(OH)D level was 18.3 (SD, 11.6) ng/mL in the whole sample; 25(OH)D levels were significantly lower in men (17.6 ng/mL) than in women (18.5 ng/mL) ($p < 0.001$). Regarding age groups, the highest 25(OH)D levels were found in the groups of 55-59, 60-64, 65-69, and 70-74 years of age, and the lowest in the groups of 80-85 years of age and older. In women, the distribution of 25(OH)D levels among age groups showed a similar pattern. However, in men, the highest mean 25(OH)D concentrations were found in the youngest groups (14-19, 20-24, and 25-29 years) whereas the lowest mean levels were recorded in the oldest groups (80-85 years and older).

Overall, the prevalence rate of vitamin D inadequacy varied according to the different cutoff values, with significant differences between genders (Table II and Table III). Using the USIOM interpretation of serum 25(OH)D levels, hypovitaminosis D (25(OH)D < 20 ng/mL) was reported in 63.7 % of patients; about half of them (30.9 % of study sample) had vitamin D deficiency (25(OH)D < 12 ng/mL). Furthermore, 32.8 % of subjects were found to have vitamin D insufficiency (25(OH)D = 20-30 ng/mL),

Table I. Mean 25(OH)D values in the study population and according to age group and gender

	n	%	Total population		n	%	Women		n	%	Men		p value ¹
			Mean (SD), ng/mL	Mean (SD), ng/mL			Mean (SD), ng/mL	Mean (SD), ng/mL					
Total population	21,490	100	18.3 (11.6)	18.5 (12.0)	16,000	74.3	18.5 (12.0)	17.6 (10.5)	5490	25.7	17.6 (10.5)	< 0.001	
Age group (years)													
14-19 ⁽¹⁾	364	1.7	18.9 (10.4) ^{(15)*}	18.2 (10.9) ^{(15)*}	250	1.6	18.2 (10.9) ^{(15)*}	20.3 (8.9) ^{(14,15)*}	114	2.1	20.3 (8.9) ^{(14,15)*}	0.07	
20-24 ⁽²⁾	397	1.9	18.6 (11.1) ^{(15)*}	18.1 (11.4) ^{(10,11,15)*}	286	1.8	18.1 (11.4) ^{(10,11,15)*}	19.7 (10.2) ^{(14,15)*}	111	2.0	19.7 (10.2) ^{(14,15)*}	0.20	
25-29 ⁽³⁾	543	2.5	18.5 (11.4) ^{(11,15)*}	17.7 (11.1) ^{(8,10,12,15)*}	419	2.6	17.7 (11.1) ^{(8,10,12,15)*}	21.4 (11.9) ^{(8,13,15)*}	124	2.3	21.4 (11.9) ^{(8,13,15)*}	0.001	
30-34 ⁽⁴⁾	699	3.3	17.6 (9.19) ^{(9,12,15)*}	17.5 (9.5) ^{(9,12,15)*}	571	3.6	17.5 (9.5) ^{(9,12,15)*}	18.1 (7.9) ^{(15)*}	128	2.3	18.1 (7.9) ^{(15)*}	0.51	
35-39 ⁽⁵⁾	916	4.3	18.3 (10.4) ^{(10,11,14,15)*}	17.6 (8.9) ^{(9,12,15)*}	708	4.4	17.6 (8.9) ^{(9,12,15)*}	20.6 (14) ^{(13,15)*}	208	3.8	20.6 (14) ^{(13,15)*}	0.001	
40-44 ⁽⁶⁾	977	4.6	17.9 (10.1) ^{(9,11,15)*}	17.9 (10.3) ^{(9,12,15)*}	761	4.8	17.9 (10.3) ^{(9,12,15)*}	18.4 (9.2) ^{(14,15)*}	216	3.9	18.4 (9.2) ^{(14,15)*}	0.53	
45-49 ⁽⁷⁾	1347	6.3	17.8 (9.7) ^{(9,12,15)*}	17.5 (9.8) ^{(9,12,15)*}	1043	6.5	17.5 (9.8) ^{(9,12,15)*}	18.7 (9.7) ^{(14,15)*}	304	5.5	18.7 (9.7) ^{(14,15)*}	0.08	
50-54 ⁽⁸⁾	1756	8.2	18.3 (9.3) ^{(9,11,14,15)*}	18.6 (9.4) ^{(9,11,15)*}	1337	8.4	18.6 (9.4) ^{(9,11,15)*}	17.6 (8.8) ^{(3,15)*}	419	7.6	17.6 (8.8) ^{(3,15)*}	0.06	
55-59 ⁽⁹⁾	1910	8.9	19.6 (11.5) ^{(4,6,8,13,15)*}	20.2 (12.1) ^{(3,8,13,15)*}	1494	9.3	20.2 (12.1) ^{(3,8,13,15)*}	17.8 (8.7) ^{(15)*}	416	7.6	17.8 (8.7) ^{(15)*}	< 0.001	
60-64 ⁽¹⁰⁾	1914	8.9	20.2 (11.2) ^{(4,5,13,15)*}	20.8 (11.4) ^{(2,9,13,15)*}	1427	8.9	20.8 (11.4) ^{(2,9,13,15)*}	18.3 (10.2) ^{(14,15)*}	487	8.9	18.3 (10.2) ^{(14,15)*}	< 0.001	
65-69 ⁽¹¹⁾	2134	9.9	20.4 (11.5) ^{(3,5,13,15)*}	21 (11.4) ^{(2,8,13,15)*}	1569	9.8	21 (11.4) ^{(2,8,13,15)*}	18.9 (11.7) ^{(14,15)*}	565	10.3	18.9 (11.7) ^{(14,15)*}	< 0.001	
70-74 ⁽¹²⁾	2055	9.6	19.4 (12.4) ^{(4,7,13,15)*}	20 (13) ^{(3,7,13,15)*}	1463	9.1	20 (13) ^{(3,7,13,15)*}	17.9 (10.6) ^{(14,15)*}	592	10.8	17.9 (10.6) ^{(14,15)*}	< 0.001	
75-79 ⁽¹³⁾	2042	9.5	18.1 (12.8) ^{(9,12,14,15)*}	18.3 (13.6) ^{(9,12,15)*}	1462	9.1	18.3 (13.6) ^{(9,12,15)*}	17.7 (10.3) ^{(3,5,14,15)*}	580	10.6	17.7 (10.3) ^{(3,5,14,15)*}	0.28	
80-84 ⁽¹⁴⁾	2163	10.1	16.7 (12.7) ^{(5,8,9,13,15)*}	17.2 (13.9) ^{(9,12,15)*}	1526	9.5	17.2 (13.9) ^{(9,12,15)*}	15.5 (9.2) ^{(1,3,5,7,10,12)*}	637	11.6	15.5 (9.2) ^{(1,3,5,7,10,12)*}	0.005	
≥ 85 ⁽¹⁵⁾	2273	10.6	14.5 (13.0) ^{(1,4)*}	14.8 (13.6) ^{(1,4)*}	1684	10.5	14.8 (13.6) ^{(1,4)*}	13.6 (11.2) ^{(1,13)*}	589	10.7	13.6 (11.2) ^{(1,13)*}	0.05	

⁽¹⁻¹⁵⁾Indicates the age group number. SD: standard deviation. *p-value < 0.05 for the differences in mean 25(OH)D levels between age groups, by ANOVA tests with Bonferroni's post hoc analysis. *p value¹ for the differences in mean 25(OH)D levels between women and men within each age group, by independent Student's t-tests.

and only 10.3 % of the whole sample had sufficient/optimal levels of vitamin D (25(OH)D > 30 ng/mL as USES cutoff) (Table II). As shown in table III, the prevalence rate of hypovitaminosis D was significantly higher in men than in women regardless of 25(OH)D cutoffs (p < 0.001). Hypovitaminosis D (25(OH)D < 20 ng/mL) was highly prevalent at 72.6 % among patients over 75 years of age, whereas in the groups of 14-29 and 55-74 years prevalence was 59.8 % and 57.6 %, respectively (Table II). The highest rate of hypovitaminosis D was found in patients over 85 years of age (79.4 %), and the lowest rate in the group of 65-69 years (54.9 %) (Fig. 1). When the cutoff value for defining hypovitaminosis D was set at < 30 ng/mL, prevalence was as high as 89.7 % in the whole sample, and it was slightly but significantly higher in men than in women (91.7 % vs 88.9 %, respectively, p > 0.001), and also in all age groups (Table III and Fig. 2).

As expected, season had a significant effect on 25(OH)D levels (Table IV and Fig. 3). March (15.1 ng/mL) was the month of the year in which the lowest 25(OH)D levels were found, whereas the highest levels were observed in September (22.2 ng/mL) (Fig. 3). The mean level of 25(OH)D in winter was significantly lower than in the other seasons (p < 0.001), with significant differences in levels between spring, summer, and autumn regardless of age group. The highest levels of 25(OH)D were found in the summer (20.9 ng/mL), with mean values above 20 ng/ml in all age groups except in patients over 75 years of age (17.7 ng/mL) (Table IV).

Hypovitaminosis D rates during winter, spring, summer, and autumn were 76.5 %, 68.7 %, 51.8 % and 58.3 %, respectively (p < 0.001) (Table V). Winter and spring were the seasons in which hypovitaminosis D was higher in all age groups as compared to summer or autumn (p < 0.001). Patients whose age was over 75 years exhibited rates of hypovitaminosis D in all seasons when compared to other age groups except for winter, where hypovitaminosis D rate was higher in the youngest (14-29 years) and/or middle-aged (30-54 years) groups (Table V).

DISCUSSION

This study explored for the first time vitamin D status in at-risk patients attending primary healthcare facilities in La Rioja, Spain. Our results have shown that hypovitaminosis D was highly prevalent in our region of study. More than two-thirds of the population were vitamin D deficient (30.9 %) or insufficient (32.8 %) according to the USIOM criteria. Vitamin D hypovitaminosis was more prevalent in men (67.1 %) than in women (62.6 %), and up to 72.6 % of patients older than 75 years had 25(OH)D levels below 20 ng/mL. Using a stricter criterion, as the USES cutoff point (30 ng/mL) for defining vitamin D insufficiency, the prevalence rate rose to as high as 92.7 %. As expected, winter was shown to be an important risk factor for decreased 25(OH)D levels.

Our findings indicate a breakthrough in identifying the burden vitamin D represents in order to control and improve the prognosis of this condition. In addition, they support the relevance

Table II. Prevalence rate (% (95 % CI)) of vitamin D deficiency, insufficiency, and sufficiency/adequacy in the study population and according to gender

25(OH)D concentrations	n	Total population		Women		Men		p-value ¹
		% (95 % CI)	n	% (95 % CI)	n	% (95 % CI)		
<i>US Institute of Medicine</i>								
< 12 ng/mL	6640	30.9 (30.3-31.5)	4920	30.7 (30.0-31.4)	1720	31.3 (30.1-32.6)	0.42	
12 to 20 ng/mL	7052	32.8 (32.2-33.4)	5089	31.8 (31.1-32.5)	1963	35.7 (34.4-37.0)	< 0.001	
> 20 ng/mL	7798	36.3 (35.6-36.9)	5991	37.4 (34.8-36.3)	1807	32.9 (30.5-33.0)	< 0.001	
p-value ²					< 0.001			
<i>US Endocrinology Society</i>								
< 20 ng/mL	13,692	63.7 (63.1-64.3)	10,009	62.6 (61.8-63.3)	3683	67.1 (65.8-68.3)	< 0.001	
20 to 30 ng/mL	5580	26.0 (25.4-26.5)	4229	26.4 (25.7-27.1)	1351	24.6 (23.5-25.7)	0.007	
> 30 ng/mL	2218	10.3 (9.9-10.8)	1762	11.0 (10.5-11.5)	456	8.3 (7.6-9.0)	< 0.001	
p-value ²					< 0.001			

p-value¹ for the differences in vitamin D status between women and men within each 25(OH)D concentration category, by the χ^2 test; p-value² for the differences in vitamin D status between women and men, by the χ^2 test.




Figure 1.

Distribution of the study population according to the USA Institute of Medicine criteria to define vitamin D status by age groups and gender. (1-4) Indicates age group number. P value < 0.05 (groups 5, 8, 9, 10, 11, 12, 14, and 15) of the differences in vitamin D (25 (OH)D <20 ng/mL) deficiency prevalence between gender for ages groups.

of vitamin D status screening as part of the routine laboratory bloodwork usually ordered by primary care physicians, especially for groups potentially at risk of inadequacy, and providing appropriate population-based interventions to combat this global public issue associated with serious consequences for health (16). Along these lines, some countries have adopted public health policies that include food fortification, and vitamin D supplementation for high-risk population groups (4).

Circulating 25(OH)D levels in the blood are subject to cyclical variations over the course of the year based on genetic (ethnicity, gender, polymorphisms), environmental, and lifestyle factors (UVB exposure, skin pigmentation, clothing style, diet, dietary supplements use), advanced age, and the presence of certain physiological or chronic diseases (4). This suggests that the number of factors contributing to vitamin D deficiency, and their interrelations, should be taken into account when it comes to making an adequate clinical decision (29). Furthermore, it is essential and imperative to guide future public health policies.

Globally, vitamin D insufficiency is prevalent in all regions of the world (2,3). The high prevalence of vitamin D deficiency observed in the current study is comparable to what was previously found in various subpopulations in Spain (17,18,30); the prevalence of 25(OH)D levels lower than 20 ng/mL ranged from 61 % among the youth (21) to 87 % among middle-aged adults (22,23); and in the elderly, whether institutionalized or living in the community, this prevalence ranged from 79 % to 91 % (24-26). Clearly, vitamin D insufficiency is the result of a complex interplay of the aforementioned factors. However, in Spain, despite being a country with many sunlight hours (mean, 2,900 hours/year) (31), a diminished or absent vitamin D synthesis in the skin might partly explain the high prevalence of insufficiency in the Spanish population, including that of La Rioja. It has been proposed that climate change has affected the availability of ultraviolet rays in the region; to be sure, in Spain's location above parallel 35 ° N, the oblique incidence of solar rays conditions that most UVB radiation be absorbed by the ozone layer.

Table III. Prevalence rate (% (95 % CI)) of hypovitaminosis D in the study population and according to age groups and gender

25(OH)D levels < 20 ng/mL										
	n	Mean (SD)	Total population % (95 % CI)	n	Mean (SD)	Women		Men	p-value ¹	p-value ²
						% (95 % CI)	n			
Total population (n = 21,490)	13,692	12.1 (4.6)	63.7 (63.1-64.3)	10,009	12.0 (4.6)	62.6 (61.8-63.3)	3683	12.3 (4.4)	67.1 (65.8-68.3)	< 0.001
Age group (years)										
14-29 (n = 1304) ⁽¹⁾	780	11.8 (4.9)	59.8 (57.1-62.4)	590	11.4 (5.1)	61.7 (58.7-64.8)	190	13.3 (4.3)	54.4 (49.2-59.7)	< 0.001
30-54 (n = 5695) ⁽²⁾	3592	12.5 (4.5)	63.2 (61.8-64.3)	2809	12.4 (4.6)	63.5 (62.1-64.9)	783	12.8 (4.3)	61.4 (58.7-64.1)	0.01
55-74 (n = 8013) ⁽³⁾	4617	13.1 (4.3)	57.6 (56.5-58.7)	3284	13.1 (4.3)	55.2 (53.9-56.4)	1333	12.9 (4.4)	64.7 (62.6-66.7)	0.10
> 75 (n = 6478) ⁽⁴⁾	4703	10.8 (4.5)	72.6 (71.5-73.7)	3326	10.6 (4.5)	71.2 (69.9-72.5)	1377	11.3 (4.4)	76.2 (74.3-78.2)	< 0.001
p value ³		1-2, 1-3, 1-4, 2-3, 2-4, 3-4	< 0.001		1-2, 1-3, 1-4, 2-3, 2-4, 3-4	< 0.001		1-4, 2-4, 3-4	< 0.001	
25(OH)D levels < 30 ng/mL										
Total population (n = 21,490)	19,272	12.1 (4.6)	89.7 (89.2-90.1)	14,238	15.6 (6.9)	88.9 (88.5-89.5)	5034	15.5 (6.6)	91.7 (90.6-92.4)	0.21
Age group (years)										
14-29 (n = 1304) ⁽¹⁾	1140	15.7 (7.2)	87.4 (85.6-89.2)	843	15.2 (7.3)	88.3 (86.2-90.3)	297	17.3 (6.5)	85.0 (81.3-88.8)	< 0.001
30-54 (n = 5695) ⁽²⁾	5181	16.1 (6.7)	90.9 (90.2-91.7)	4015	15.9 (6.7)	90.8 (89.9-91.6)	1166	16.6 (6.6)	91.3 (89.9-92.9)	0.003
55-74 (n = 8013) ⁽³⁾	7020	16.9 (6.5)	87.6 (86.9-88.3)	5135	17.2 (6.6)	86.3 (85.4-87.1)	1885	16.2 (6.4)	91.5 (90.3-92.7)	< 0.001
> 75 (n = 6478) ⁽⁴⁾	5931	13.6 (6.9)	91.6 (90.8-92.2)	4245	13.6 (7.1)	90.8 (90.0-91.6)	1686	13.7 (6.5)	93.2 (92.2-94.5)	0.001
p-value ³		1-3, 1-4, 2-3, 2-4, 3-4	< 0.001		1-2, 1-3, 1-4, 2-3, 2-4, 3-4	< 0.001		1-3, 1-4, 2-4, 3-4	< 0.001	

⁽¹⁻⁴⁾Indicates age group number. SD: standard deviation. p-value¹ of the differences in mean 25(OH)D levels between women and men within each age group, by independent Student's t-tests. p-value² for the differences in hypovitaminosis D between women and men within each age group, by the χ^2 test. p-value³ < 0.05 for the differences in mean 25(OH)D levels (by ANOVA tests with Bonferroni's post hoc analysis) or hypovitaminosis D (by χ^2 tests) between age groups.




Figure 2.

Distribution of the study population according to the USA Endocrinology Society criteria to define vitamin D status by age groups and gender. (1-4) Indicates age group number. P value < 0.05 (groups 8-12, and 14) of the differences in vitamin D (25 (OH)D <30 ng/mL) deficiency prevalence between gender for ages groups.

Table IV. Mean vitamin D (ng/mL) levels according to age group and season of the year

	n	Winter	n	Spring	n	Summer	n	Autumn	p-value ¹	p-value ²	p-value ³
		Mean (SD)		Mean (SD)		Mean (SD)		Mean (SD)			
Total population (n = 21,490)	4991	15.3 (11.1)	5652	17.2 (11.1)	5043	20.9 (11.9)	5804	19.6 (11.6)	< 0.001	< 0.001	< 0.001
<i>Age group (years)</i>											
14-29 (n = 1304) ⁽¹⁾	284	13.5 (8.9)	286	16.4 (10.29)	354	23.5 (11.2)	380	19.6 (10.7)	0.001	< 0.001	< 0.001
30-54 (n = 5695) ⁽²⁾	1348	14.2 (9.4)	1436	16.8 (8.9)	1301	22.3 (9.7)	1610	18.9 (9.2)	< 0.001	< 0.001	< 0.001
55-74 (n = 8013) ⁽³⁾	1906	16.8 (11.0)	2130	19.1 (11.9)	1760	22.4 (11.7)	2217	21.5 (11.3)	< 0.001	< 0.001	< 0.001
> 75 (n = 6478) ⁽⁴⁾	1453	14.7 (12.7)	1800	15.5 (11.6)	1628	17.7 (13.3)	1597	17.5 (13.9)	0.09	< 0.001	< 0.001
p-value ⁴		1-3, 2-3, 3-4		1-3, 2-3, 2-4, 3-4		1-4, 2-4, 3-4		1-3, 1-4, 2-3, 2-4, 3-4			

⁽¹⁻⁴⁾Indicate age group number. p-values for the differences in mean 25(OH)D levels between winter and spring (p-value¹), or summer (p-value²), or autumn (p-value³) within each age group, by independent Student's t-tests. p-value⁴ < 0.05 for significant differences in mean 25(OH)D levels between age groups within each season of the year, by ANOVA test with Bonferroni's post hoc analysis.




Figure 3.

Mean vitamin D (ng/ml) levels according to month of vitamin D test.

As a consequence, in the last few years, European and Spanish populations have been less exposed to ultraviolet rays, which has compromised cutaneous vitamin D synthesis, particularly during the winter (32). In this context, Serrano et al. (33) estimated the time needed to obtain the recommended daily dose of 1000 IU in Valencia, a Spanish province at 39 ° N. It was found that around midday (between 12:30 and 13:30 h) in January, with 10 % of the body exposed, more than two hours of solar exposure are necessary to obtain an optimal vitamin D dose, whereas the rest of the year exposure time ranges from 7 min in July to 31 min in October (33). Thus, the low 25(OH)D levels observed in our at-risk population in La Rioja (42 ° N) perhaps do not represent an unexpected problem, which may be speculatively accounted for by scarce contents of vitamin D in the diet that cannot be compensated for by vitamin D synthesis in the skin following exposure to UVB, all of this combined with darker skin pigmentation, sunscreen use, and clothing.

In the current study, serum 25(OH)D levels were significantly lower in men than in women. Also in agreement with previous

Table V. Prevalence rate (% (95 % CI)) of hypovitaminosis D in the study population, according to age group and season of the year

	25(OH)D levels < 20 ng/mL											
	Winter		Spring		Summer		Autumn		p-value ¹		p-value ²	
	n	% (95 % CI)	n	% (95 % CI)	n	% (95 % CI)	n	% (95 % CI)				
Total population (n = 21,490)	3817	76.5 (63.1-64.3)	3883	68.7 (61.8-63.3)	2611	51.8 (65.8-68.3)	3381	58.3 (65.8-68.3)	< 0.001	< 0.001	< 0.001	< 0.001
Age group (years)												
14-29 (n = 1304)	231	81.3 (76.8-85.8)	209	73.1 (67.9-78.2)	132	37.3 (32.2-42.3)	208	54.7 (49.7-59.7)	0.03	< 0.001	< 0.001	< 0.001
30-54 (n = 5695)	1102	81.7 (79.7-83.8)	1007	70.1 (67.7-72.5)	543	41.7 (39.1-44.4)	940	58.4 (55.9-60.8)	< 0.001	< 0.001	< 0.001	< 0.001
55-74 (n = 8013)	1356	71.1 (69.1-73.1)	1312	61.6 (59.5-63.6)	825	46.9 (44.5-49.2)	1124	50.7 (48.6-52.8)	< 0.001	< 0.001	< 0.001	< 0.001
> 75 (n = 6478)	1128	77.6 (75.4-79.8)	1355	75.8 (73.3-77.3)	1111	68.2 (65.9-70.5)	1109	69.4 (67.2-71.7)	0.13	< 0.001	< 0.001	< 0.001
p-value ⁴		< 0.001		< 0.001		< 0.001		< 0.001				

p-value for the differences in hypovitaminosis D between winter and spring (p-value¹), summer (p-value²), or autumn (p-value³) within age groups, by the χ^2 test; p-value⁴ is for the differences in hypovitaminosis D between age groups within each season of the year, by the χ^2 test.

studies (2,30), vitamin D deficiency (< 20 ng/mL) affected males and females differently, being more prevalent among men, especially among middle-aged adults (55-74 years) and the elderly over 75 years of age. The presence of intrinsic sexual dimorphisms at the molecular and cellular levels, and of differing sex hormones including sex hormone-binding globulin (SHBG), estradiol, and testosterone, may be responsible for these sex differences (34,35). Furthermore, sex differences might also be partly explained by other factors such as adiposity and fat distribution, smoking, physical activity, and clothing/eating habits. However, neither sex hormones nor these sex-specific factors were determined in the current study. Perhaps, sex disparities regarding the use of health services result in a sample where women represent almost 75 % of subjects. This might lead to an underestimation of the overall prevalence of vitamin D deficiency in men. The highest rate of hypovitaminosis D observed in patients with advancing age might be explained by changes in lifestyle and living conditions. The elderly population has been shown to increase indoor and sedentary activities, and to reduce social life; as a consequence, there is a decrease in sunlight exposure (36) that, along with a reduced ability to synthesize vitamin D (4) and conservative clothing habits, might result in a suboptimal vitamin D status. The decrease in food intake and gut absorption (37), medication use, and pre-existing risk factors usually associated with ageing could provide further explanations. We observed an opposite gender association for vitamin D deficiency in younger people (14-29 years), with women exhibiting here a higher prevalence. This may have been due to the fact that in populations at productive ages regular sun exposure tends to decrease because of changes in lifestyle, with an increased amount of time spent in indoor activities. The liberal use of sunscreen, a pattern especially common among younger women when compared to men, may also support these findings (38). Therefore, in order to ensure sunlight exposure, a promotion of suitable outdoor activities for the different age groups should be included in the setting of health care counseling.

We observed, as in the rest of the world and in other regions of Spain (2,30,33,39), a seasonal variation in serum 25(OH)D levels, with the highest levels being found at the end of summer (August and September), and the lowest levels at the end of winter (February and March). In the current study, the prevalence of hypovitaminosis D followed a seasonal pattern ranging from 51.8 % to 76.5 %, with higher rates in winter in all age groups as compared to summer or autumn. Few hours of sunlight exposure, limited cutaneous synthesis of vitamin D, and the excessive exposure time that is required may account for the low 25(OH)D levels recorded during the months of winter. Patients over 75 years exhibited high rates of deficiency or insufficiency throughout the year; however, in winter, the most severely affected age groups included the youngest sample and middle-aged adults. In winter, one individual may require more than two hours of sun exposure to obtain an optimal dose of vitamin D (33). Thus, it is possible that the oldest group of people had spent more minutes in the sun during the middle hours of the day during winter.

Finally, as of today, the optimum serum level of 25-hydroxyvitamin D remains controversial. It was remarkable for us to find that, when 30 ng/mL and 20 ng/mL were the USES-based cutoff values for vitamin D insufficiency and deficiency, respectively, as many as 89.7 % of patients had insufficient levels and 63.7 % of these were even classified as deficient in vitamin D. This finding suggests that the cutoff value proposed by different international (27,40) and Spanish (41) scientific societies for sufficient or desirable vitamin D levels, namely above 30 ng/mL, invariably shows an important prevalence of subclinical vitamin D deficiency in our at-risk population. In fact, many clinicians frequently report that many healthy people are probably misclassified as deficient in vitamin D.

The main strengths to be considered for our study include a large sample for whom information on 25(OH)D concentrations was available, and the use of the Liaison test, which assesses total 25(OH)D concentrations (i.e., vitamins D2 and D3). However, our study has an important limitation that must be commented on. First, our sample was shaped by an at-risk population instead of healthy subjects, and most of the patients included were women; therefore, the study did not fulfil the criteria for representativeness of the population. However, we examined vitamin D levels in patients with no vitamin D supplementation. Therefore, the prevalence of hypovitaminosis D observed probably reflects the true figure for the general population. Second, the amount of time that our patients spent in the sun was not recorded. We did, however, use the season as a proxy for sun exposure. Third, lifestyle factors such as physical activity and diet, or pathological conditions, were not considered, and they are factors that should be considered in future studies.

CONCLUSIONS

In La Rioja, Spain, hypovitaminosis D was highly prevalent among PHC users, especially in the older age groups and during the winter months. In order to prevent vitamin D deficiency, health-care advice should be focused on ensuring an optimal vitamin D status through adherence to healthy lifestyles, including a complete and varied diet rich in vitamin D food sources and promotion of outdoor activities, especially during the winter months when sunlight hours tend to decrease.

REFERENCES

- Holick MF. The vitamin D deficiency pandemic: Approaches for diagnosis, treatment and prevention. *Rev Endocr Metab Disord* 2017;18:153-65.
- Palacios C, Gonzalez L. Is vitamin D deficiency a major global public health problem? *J Steroid Biochem Mol Biol* 2014;144:138-45.
- Roth DE, Abrams SA, Aloia J, Bergeron G, Bourassa MW, Brown KH, et al. Global prevalence and disease burden of vitamin D deficiency: a road-map for action in low- and middle-income countries. *Ann NY Acad Sci* 2018;1430(1).
- Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357:266-81.
- Hilger J, Friedel A, Herr R, Rausch T, Roos F, Wahl D A, et al. A systematic review of vitamin D status in populations worldwide. *Br J Nutr* 2014;111:23-45.
- Spiro A, Buttriss JL. Vitamin D: An overview of vitamin D status and intake in Europe. *Nutr Bull* 2014;39:322-50.
- Kimlin MG. Geographic location and vitamin D synthesis. *Mol Aspects Med* 2008;29:453-61.
- Institute of Medicine. Food and Nutrition Board. Dietary Reference Intakes for Calcium and Vitamin D. Washington, DC: National Academy Press; 2010.
- Uday S, Höglér W. Nutritional Rickets and Osteomalacia in the Twenty-first Century: Revised Concepts, Public Health, and Prevention Strategies. *Curr Osteoporos Rep* 2017;15:293-302
- Christakos S, Dhawan P, Verstuyf A, Verlinden L, Carmeliet G. Vitamin D: Metabolism, molecular mechanism of action, and pleiotropic effects. *Physiol Rev* 2015;96:365-408.
- Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Guidelines for preventing and treating vitamin D deficiency and insufficiency revisited. *J Clin Endocrinol Metab* 2012;97:1153-8.
- Basit S. Vitamin D in health and disease: a literature review. *Br J Biomed Sci* 2013;70:161-72.
- Autier P, Boniol M, Pizot C, Mullie P. Vitamin D status and ill health: a systematic review. *Lancet Diabetes Endocrinol* 2014;2:76-89.
- Kheiri B, Abdalla A, Osman M, Ahmed S, Hassan M, Bachuwa G. Vitamin D deficiency and risk of cardiovascular diseases: a narrative review. *Clin Hypertens* 2018;24:9.
- Nair R, Maseeh A. Vitamin D: The "sunshine" vitamin. *J Pharmacol Pharmacother* 2012;3:118-26.
- Muscogiuri G, Altieri B, Annweiler C, Balercia G, Pal HB, Boucher BJ, et al. Vitamin D and chronic diseases: the current state of the art. *Arch Toxicol* 2017;91:97-107.
- González-Molero I, Morcillo S, Valdés S, Pérez-Valero V, Botas P, Delgado E, et al. Vitamin D deficiency in Spain: a population-based cohort study. *Eur J Clin Nutr* 2011;65:321-8.
- Rigueira García AI. Recomendaciones sobre suplementos de vitamina D y calcio para las personas adultas en España. *Rev Esp Salud Publica* 2012;86:461-82.
- Navarro Valverde C, Quesada Gómez JM. Vitamin D deficiency in Spain. Reality or myth? *Rev Osteoporos Metab Miner* 2014;6:5-10.
- Togo A, Espadas Maciá D, Blanes Segura S, Sivó Díaz N, Villalba Martínez C. Is there vitamin D deficiency in children in a sunny Mediterranean city? *An Pediatr* 2016;84:163-9.
- González-Padilla E, Soria López A, González-Rodríguez E, García-Santana S, Mirallave-Pescador A, Groba Marco M del V, et al. High prevalence of hypovitaminosis D in medical students in Gran Canaria, Canary Islands (Spain). *Endocrinol Nutr* 2011;58:267-73.
- Vaqueiro M, Bare M, Anton E, Andreau E, Moya A, Sampere R, et al. Hypovitaminosis D associated to low sun exposure in the population over 64 years old. *Med Clin (Barc)* 2007;129:287-91.
- Díaz-López A, Jardí C, Villalobos M, Serrat N, Basora J, Arija V. Prevalence and risk factors of hypovitaminosis D in pregnant Spanish women. *Sci Rep* 2020;10:15757.
- Quesada-Gómez JM, Díaz-Curiel M, Sosa-Henríquez M, Malouf-Sierra J, Nogues-Solan X, Gomez-Alonso C, et al. Low calcium intake and inadequate vitamin D status in postmenopausal osteoporotic women. *J Steroid Biochem Mol Biol* 2013;136:175-7.
- Niño Martín V, Pérez Castrillón JL. Niveles de vitamina D en población mayor de 65 años. *REEMO* 2008;17:1-4.
- Almirall J, Vaqueiro M, Baré ML, Anton E. Association of low serum 25-hydroxyvitamin D levels and high arterial blood pressure in the elderly. *Nephrol Dial Transplant* 2010;25:503-9.
- Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Evaluation, Treatment, and Prevention of Vitamin D Deficiency: an Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab* 2011;96:1911-30.
- Institute of Medicine (US) Committee to Review Dietary Reference Intakes for Vitamin D and Calcium, Ross AC, Taylor CL, Yaktine AL, Del Valle HB. Dietary Reference Intakes for Calcium and Vitamin D. National Academies Press (US); 2011.
- Pilz S, Zittermann A, Trummer C, Theiler-Schwetz V, Lerchbaum E, Keppel MH, et al. Vitamin D testing and treatment: A narrative review of current evidence. *Endocr Connect* 2019;8:27-43.
- Valverde NC, Gómez QJ, Manuel J. Deficiencia de vitamina D en España. ¿Realidad o mito? La 25-hidroxivitamina D el marcador del estatus corporal de vitamina D. *Rev Osteoporos Metab Min* 2014;S5-10.
- Instituto Nacional de Estadística (INE). Anuario Estadístico de España; 2019. Available from: https://www.ine.es/prodyser/pubweb/anuario19/anu19_completo.pdf
- Lai AT, Lin IC, Yang YW, Wu MF. Climate change and human health. *J Intern Med* 2012;23:343-50.

33. Serrano MA, Cañada J, Moreno JC, Gurrea G. Solar ultraviolet doses and vitamin D in a northern mid-latitude. *Sci Total Environ* 2017;574:744-50.
34. Anic GM, Albanes D, Rohrmann S, Kanarek N, Nelson WG, Bradwin G, et al. Association between serum 25-hydroxyvitamin D and serum sex steroid hormones among men in NHANES. *Clin Endocrinol (Oxf)* 2016;85:258-66.
35. Zhao D, Ouyang P, de Boer IH, Lutsey PL, Farag YMK, Guallar E, et al. Serum vitamin D and sex hormones levels in men and women: The Multi-Ethnic Study of Atherosclerosis (MESA). *Maturitas* 2017;96:95-102.
36. Rodríguez Sangrador M, Beltrán de Miguel B, Cuadrado Vives C, Moreiras Tuni O. Comparative analysis of vitamin D status and solar exposition habits in adolescent and elderly Spanish women. The Five Countries Study (OPTIFORD Project). *Nutr Hosp* 26:609-13.
37. Bhutto A, Morley JE. The clinical significance of gastrointestinal changes with aging. *Curr Opin Clin Nutr Metab Care* 2008;11:651-60.
38. Holman DM, Berkowitz Z, Guy GP, Hawkins NA, Saraiya M, Watson M. Patterns of sunscreen use on the face and other exposed skin among US adults. *J Am Acad Dermatol* 2015;73:83-92.e1.
39. Pereda CA, Nishishinya MB, Roldan EJA. 25-Hydroxyvitamin D serum levels in rheumatic female patients in southeast Spain: The paradigm of daily optimal sunshine levels and inadequate vitamin D status. *Endocrinol Diabetes Nutr* 2019;66:181-7.
40. Dawson-Hughes B, Mithal A, Bonjour JP, Boonen S, Burckhardt P, Fuleihan GEH, et al. IOF position statement: Vitamin D recommendations for older adults. *Osteoporos Int* 2010;21:1151-4.
41. Gómez de Tejada Romero MJ, Sosa Henríquez M, Del Pino Montes J, Jódar Gimeno E, Quesada Gómez JM, Cancelo Hidalgo MJ, et al. Documento de posición sobre las necesidades y niveles óptimos de vitamina D. *Rev Osteoporos Metab Min* 2011;3:53-64.