

# **Nutrición Hospitalaria**



**Beneficios de la suplementación  
con micronutrientes sobre el  
estado nutricional, el  
metabolismo energético y el  
bienestar subjetivo**

**Benefits of micronutrient  
supplementation on nutritional  
status, energy metabolism, and  
subjective wellbeing**

10.20960/nh.03788

12/28/2021

## **Benefits of micronutrient supplementation on nutritional status, energy metabolism, and subjective wellbeing**

*Beneficios de la suplementación con micronutrientes sobre el estado nutricional, el metabolismo energético y el bienestar subjetivo*

Silvia Maggini<sup>1</sup>, Veronika Óvári<sup>1</sup>, Inmaculada Ferreres Giménez<sup>2</sup>, and María Gloria Pueyo Alamán<sup>2</sup>

<sup>1</sup>Bayer Consumer Care AG. Basel, Switzerland. <sup>2</sup>Bayer Hispania Consumer Health. San Joan Despí. Barcelona, Spain

**Correspondence:** Veronika Óvári. Bayer Consumer Care AG. Peter Merian-Strasse 84. 4052 Basel, Switzerland  
e-mail: veronika.ovari@bayer.com

*Acknowledgements: this article was funded by Bayer Consumer Care, AG, including the services of the medical writer who drafted and edited the manuscript (Deborah Nock, Medical WriteAway, Norwich, UK), with full review and approval by all authors.*

*Conflict of interest: the authors declare no conflict of interest.*

### **ABSTRACT**

The human body, particularly the brain, requires energy, stored in the form of adenosine triphosphate. Energy metabolism during cellular respiration is dependent on the presence of multiple micronutrients, which act as essential components, coenzymes, or precursors at every stage. An adequate supply of multiple micronutrients is vital for efficient energy production. However, micronutrient intakes below the recommended dietary allowance are common, even in industrialized

countries. Intakes of vitamins A, D, E, folate, iron, zinc, and selenium are suboptimal across all age groups. Suboptimal micronutrient levels have been shown to contribute to low energy levels, physical and mental fatigue, and impaired cognitive performance and wellbeing – symptoms frequently present in the general population. When supplemented in combination in well-conducted trials, multiple micronutrients ± coenzyme Q10 reduced oxidative stress in chronic fatigue syndrome; in healthy people they increased cerebral blood-flow hemodynamic response, energy expenditure, and fat oxidation; reduced mental and physical fatigue; improved the speed and accuracy of cognitive function during demanding tasks; and reduced stress. The results from these clinical trials suggest that even in industrialized countries, where adults might be assumed to have a healthy, balanced diet, there is a rationale to supplement with multiple micronutrients, including coenzyme Q10, to improve nutritional status, support energy metabolism, and improve subjective wellbeing.

**Keywords:** Coenzyme Q10. Cognitive performance. Energy metabolism. Fatigue. Multiple micronutrient supplementation.

## **RESUMEN**

El cuerpo humano, particularmente el cerebro, requiere energía, almacenada en forma de adenosina trifosfato. El metabolismo de la energía durante la respiración celular depende de la presencia de múltiples micronutrientes, que actúan como componentes esenciales, coenzimas o precursores en cada etapa. Un aporte adecuado de múltiples micronutrientes es vital para una producción eficiente de energía. Sin embargo, la ingesta de micronutrientes inferior a la recomendada es frecuente, incluso en los países industrializados. Las ingestas de vitaminas A, D, E, folato, hierro, zinc y selenio son subóptimas en todos los grupos de edad. Se ha demostrado que las

situaciones subóptimas en relación con diversos micronutrientes contribuyen a tener niveles bajos de energía, fatiga física y mental, y deterioro del rendimiento cognitivo y el bienestar, síntomas presentes a menudo en la población general. Sin embargo, cuando se suplementa en ensayos bien controlados, con una combinación de diversos micronutrientes  $\pm$  coenzima Q10, se constata una reducción del estrés oxidativo en el síndrome de fatiga crónica y, en las personas sanas, se observa un aumento de la respuesta hemodinámica del flujo sanguíneo cerebral, el gasto energético y la oxidación de la grasa; una reducción de la fatiga mental y física; una mejora de la velocidad y la precisión de la función cognitiva durante la realización de tareas exigentes, y una reducción del estrés. Los resultados de estos ensayos clínicos sugieren que, incluso en los países industrializados, donde se podría suponer que los adultos tienen una dieta saludable y equilibrada, hay motivos para complementarla con múltiples micronutrientes, incluida la coenzima Q10, con el fin de mejorar el estado nutricional, respaldar el metabolismo energético y mejorar el bienestar subjetivo.

**Palabras clave:** Coenzima Q10. Rendimiento cognitivo. Metabolismo energético. Fatiga. Suplementación con varios micronutrientes.

## INTRODUCTION

Low energy, physical and mental fatigue, and consequent adverse effects on cognitive performance (1) frequently occur in the general population. Most people who go to their doctor with an unexplained complaint (particularly females) present with fatigue (2). Micronutrient intakes below the recommended dietary allowance (RDA) are also common, even in industrialized countries (3-5). But are there any links between these two factors? And is there a rationale and evidence to indicate that supplementation with multiple

micronutrients can help to restore micronutrient levels and thereby have a beneficial effect on energy, cognition, and hence wellbeing?

## **MULTIPLE MICRONUTRIENTS ARE ESSENTIAL FOR ENERGY METABOLISM**

To stay alive, grow and reproduce, all parts of the human body require energy for processes such as cell division, homeostasis, active transport (i.e., moving molecules against concentration gradients), and transmission of nerve impulses. Even planning complex tasks and making decisions require a lot of energy, and the brain is the largest consumer of energy in the body (6).

During energy production, ingested macronutrients are broken down into glucose, fatty acids, and/or amino acids and utilized by cells to form energy, primarily stored in adenosine triphosphate (ATP) (7). The body's preferred dietary source for synthesizing ATP is glucose, which undergoes a series of metabolic reactions mainly in the mitochondria, collectively known as cellular respiration (Fig. 1). The brain, which is unable to store energy and therefore requires a continuous supply of glucose, consumes approximately 20 % of glucose-derived energy at rest, with acute dynamic variations corresponding to neuronal signaling in the brain. Metabolic regulation is critical to neuronal function.

Every stage of energy metabolism is dependent on the presence of multiple micronutrients, which act as essential components, coenzymes, or precursors (Fig. 1). These include B vitamins and vitamin C, iron, magnesium, zinc, copper, manganese, and sulfur, as well as coenzyme Q10 (CoQ10), a driver of ATP formation. CoQ10, which has antioxidant properties, also helps to regulate reactive oxygen species and reduce oxidative stress (8,9), known to play a role in chronic fatigue (10). In addition to energy production, B vitamins have vital roles in many aspects of brain function, such as DNA/RNA synthesis and/or repair, and the synthesis of numerous neurochemicals and signaling molecules (Fig. 2) (11). Other

micronutrients also play a role in brain function, including zinc in receptor binding and the formation of metalloproteins, calcium in signal transduction and the membrane potential, and zinc and magnesium in neurotransmission (12).

### **SUBOPTIMAL NUTRITION CAN AFFECT ENERGY METABOLISM - AND IS COMMON IN THE GENERAL POPULATION**

Even mild deficiencies in micronutrients in 'healthy' adults may decrease enzymatic activity and impair energy production, resulting in lack of energy and general fatigue (13). Suboptimal levels of B vitamins and the inability to meet metabolic demands lead to poor mood and irritability, and may also have a negative impact on the ability to concentrate, general mental processes, and ultimately cognition (11,14) (Fig. 2). Several factors can affect micronutrient stores, including dietary intake, physical and mental exertion (multiple micronutrients are necessary to replace the energy that is consumed), illness, seasonal demands (such as lower vitamin D levels in the winter), and factors specific to women (e.g., menstruation leads to low iron levels and reports of tiredness in 71 % of women (15)) (13).

Ideally, a sufficient and balanced diet should cover daily micronutrient requirements. Yet a healthy diet can be difficult to achieve and is dependent on a wide range of social, economic, educational, ethnic, and cultural factors. Thus, suboptimal (i.e., below the RDA) micronutrient intakes are surprisingly common – even in industrialized countries (Fig. 2). Reported suboptimal micronutrient intakes vary according to age, but in Europe include vitamins A, D, E, folate, iron, zinc, and selenium across all age groups, vitamin C and copper in adolescents and adults, and B vitamins, calcium, and copper in older adults (16). In a recent evaluation of healthy blood donors in Italy, only a low proportion had adequate plasma levels of folate and vitamin B12, but a high proportion had elevated levels of the potentially toxic amino acid, homocysteine (17).

## **SUPPLEMENTATION WITH MULTIPLE MICRONUTRIENTS MAY IMPROVE NUTRITIONAL STATUS AND SUPPORT ENERGY METABOLISM**

Even when used alone (mostly in trials conducted in women (13)), micronutrients reduced fatigue (vitamins B1 and D, iron), increased activity and feelings of being energetic (vitamin B1), increased muscle endurance (zinc), improved exercise tolerance (magnesium) and physical capacity/performance (CoQ10 (18)), and improved physical recovery (CoQ10 (19)). They also had a beneficial impact on wellbeing and the feeling of being clearheaded and composed (vitamin B1), and on symptoms of seasonal affective disorder, including depression (vitamin D). Thus, it is logical to include such vitamins and minerals in multiple micronutrient supplementation (MMS) to improve/support several aspects of mental and physical wellbeing.

Due to their strong interrelationship in metabolic pathways, an adequate supply of multiple micronutrients is vital for efficient energy production. However, as seen in figure 2, deficiencies of multiple micronutrients are prevalent in the general population. Hence, there is a strong rationale to use a supplement containing a combination of micronutrients, particularly B vitamins, to improve energy levels. Several clinical trials have demonstrated that MMS can improve nutritional status (20-22). Benefits of MMS included increased energy levels, reductions in physical and mental fatigue, improvements in concentration, mental stamina, and mood, and reductions in feelings of depression, anxiety, and stress (13).

One prospective open-label trial (10) and three double-blind, randomized, controlled trials (23-25) have evaluated the effects of MMS  $\pm$  CoQ10 on energy, cognition, and wellbeing (Fig. 3). The formulation (Supradyn<sup>®</sup>, Bayer) differed slightly within each trial, but mostly contained the water-soluble vitamins in amounts that were three times their RDA (10,23-25).

The open-label study was carried out in women with chronic fatigue syndrome (CFS) (10). Oxidative stress (due to diminished antioxidant capacity and/or decreased activity of antioxidant enzymes) plays a prominent role in CFS, and can lead to fatigue, pain, gastrointestinal problems, and cognitive impairment (10). Thus, treatment with a micronutrient supplement could improve symptoms in CFS patients. In women with CFS, MMS for two months was prospectively evaluated to determine objective parameters of antioxidant capability, including superoxide dismutase (SOD) activity, and self-assessment of improvement (10). Results indicated that MMS significantly reduced fatigue and SOD activity (suggesting that antioxidant activity improved with supplementation). There were also significant improvements in sleep disorders, autonomic nervous system symptoms (e.g., dizziness, anxiety, etc.), frequency and intensity of headaches, and subjective feelings of infection.

The other three randomized, double-blind, placebo-controlled studies evaluated the effects of MMS  $\pm$  CoQ10 in healthy subjects (23-25).

In healthy females (n = 216), the effects of MMS (3RDA) for two months on indicators of energy (fatigue and cognitive function) were compared with placebo (23). Supplementation reduced fatigue and improved the speed and accuracy of multitasking, and the authors suggested that MMS could ultimately help in the completion of demanding tasks. It is likely that the addition of CoQ10 (essential for ATP production) to MMS could have further beneficial effects.

In another study in healthy women (n = 97), the effects of acute and chronic administration of MMS + CoQ10 (1RDA) or a higher dose of MMS with no CoQ10 (3RDA) were compared to placebo (24). During the cognitive task performance, only the single dose of MMS + CoQ10 was able to increase the cerebral blood-flow hemodynamic response in the frontal cortex during tasks that activate this brain area. Supplementation with the higher dose led to a dose-dependent increase in total energy expenditure after acute (60 minutes) and chronic (8 weeks) supplementation, and fat oxidation after a single

dose. Both supplements significantly increased vitamin D3 levels and decreased homocysteine levels compared with placebo after 8 weeks. However, there were no interpretable effects on mood or cognitive performance with either dose of supplement compared with placebo. To assess the impact of MMS + CoQ10 (3RDA) on metabolically challenging exercise and cognitive tasks, supplementation for 28 days was compared to placebo in healthy young women *and* men with moderate activity levels (25). Supplementation was found to increase energy expenditure during cognitive tasks even after a single dose, and this effect was sustained when measured a day after chronic administration had concluded; the observed effects were greater in males. Carbohydrate oxidation and energy expenditure during exercise also increased in males - the first time this has been demonstrated. In females, subjective mental and physical tiredness were reduced during exercise, and stress ratings were lower after cognitive tasks. The effects on subjective mental tiredness during exercise in males and females were noted after acute assessment (commencing 45 min after intake) but were greatest after chronic intake. Ferritin levels were higher in males and females after 28 days supplementation in the subset who provided blood samples. Males were found to have sufficient levels at baseline, while females were marginally deficient. The study is important because it shows that even a single dose of MMS + CoQ10 can modulate energy expenditure and subjective tiredness and highlights the cumulative and sustained effect of chronic supplementation.

## **CONCLUSIONS**

Suboptimal (i.e., below the RDA) micronutrient intakes are common, even in industrialized countries. This can have a direct impact on energy levels, cognitive performance, and general wellbeing. Suboptimal nutrition, combined with the interrelationship between numerous vitamins and minerals in metabolic pathways, suggests that supplementation with multiple micronutrients is justified in the

general population. Several trials have evaluated the effects of MMS  $\pm$  CoQ10 in healthy women and men and in women with CFS, and have shown that it has clinically-proven beneficial effects on energy, cognitive performance, and wellbeing.

Nutrición  
Hospitalaria

## REFERENCES

1. Slimani M, Znazen H, Bragazzi NL, Zguira MS, Tod D. The Effect of Mental Fatigue on Cognitive and Aerobic Performance in Adolescent Active Endurance Athletes: Insights from a Randomized Counterbalanced, Cross-Over Trial. *J Clin Med* 2018;7(12):510. DOI: 10.3390/jcm7120510
2. Koch H, van Bokhoven MA, ter Riet G, van der Weijden T, Dinant GJ, Bindels PJ. Demographic characteristics and quality of life of patients with unexplained complaints: a descriptive study in general practice. *Qual Life Res* 2007;16(9):1483-9. DOI: 10.1007/s11136-007-9252-y
3. Rippin HL, Hutchinson J, Jewell J, Breda JJ, Cade JE. Adult Nutrient Intakes from Current National Dietary Surveys of European Populations. *Nutrients* 2017;9(12). DOI: 10.3390/nu9121288
4. Elmadfa I, Meyer A, Nowak V, Hasenegger V, Putz P, Verstraeten R, et al. European Nutrition and Health Report. *Forum Nutr* 2009; Ann Nutr Metab 2009;55(Suppl 2):1-40. DOI: 10.1159/isbn.978-3-8055-9353-3
5. Cowan AE, Jun S, Tooze JA, Eicher-Miller HA, Dodd KW, Gahche JJ, et al. Total Usual Micronutrient Intakes Compared to the Dietary Reference Intakes among U.S. Adults by Food Security Status. *Nutrients* 2019;12(1):38. DOI: 10.3390/nu12010038
6. Harris JJ, Jolivet R, Attwell D. Synaptic energy use and supply. *Neuron* 2012;75(5):762-77. DOI: 10.1016/j.neuron.2012.08.019
7. Da Poian AT, El-Bacha T, Luz MRMP. Nutrient Utilization in Humans: Metabolism Pathways. *Nature Education* 2010;3(9):11.
8. Sangsefidi ZS, Yaghoubi F, Hajiahmadi S, Hosseinzadeh M. The effect of coenzyme Q10 supplementation on oxidative stress: A systematic review and meta-analysis of randomized controlled clinical trials. *Food Science & Nutr* 2020;8(4):1766-76. DOI: 10.1002/fsn3.1492

9. Akbari A, Mobini GR, Agah S, Morvaridzadeh M, Omid A, Potter E, et al. Coenzyme Q10 supplementation and oxidative stress parameters: a systematic review and meta-analysis of clinical trials. *Eur J Clin Pharmacol* 2020;76(11):1483-99. DOI: 10.1007/s00228-020-02919-8
10. Maric D, Brkic S, Mikic AN, Tomic S, Cebovic T, Turkulov V. Multivitamin mineral supplementation in patients with chronic fatigue syndrome. *Med Sci Monit* 2014;20:47-53. DOI: 10.12659/MSM.889333
11. Kennedy D. B vitamins and the brain: Mechanisms, dose and efficacy—a review. *Nutrients* 2016;8(2):68. DOI: 10.3390/nu8020068
12. Huskisson E, Maggini S, Ruf M. The role of vitamins and minerals in energy metabolism and well-being. *J Int Med Res* 2007;35:277-89. DOI: 10.1177/147323000703500301
13. Wishart K. Increased micronutrient requirements during physiologically demanding situations: Review of the current evidence. *Vitamin Miner* 2017;6(3):1-16. DOI: 10.4172/2376-1318.1000166
14. Adan RAH, van der Beek EM, Buitelaar JK, Cryan JF, Hebebrand J, Higgs S, et al. Nutritional psychiatry: Towards improving mental health by what you eat. *Eur Neuropsychopharmacol* 2019;29(12):1321-32. DOI: 10.1016/j.euroneuro.2019.10.011
15. Schoep ME, Nieboer TE, van der Zanden M, Braat DDM, Nap AW. The impact of menstrual symptoms on everyday life: a survey among 42,879 women. *Am J Obstet Gynecol* 2019;220(6):569.e561-569.e567. DOI: 10.1016/j.ajog.2019.02.048
16. Maggini S, Pierre A, Calder PC. Immune function and micronutrient requirements change over the life course. *Nutrients* 2018;10(10):1531. DOI: 10.3390/nu10101531

17. Bortolus R, Filippini F, Udali S, Rinaldi M, Genesini S, Gandini G, et al. B vitamin blood concentrations and one-carbon metabolism polymorphisms in a sample of Italian women and men attending a unit of transfusion medicine: a cross-sectional study. *Eur J Nutr* 2021;60(5):2643-54. DOI: 10.1007/s00394-020-02448-1
18. Rosenfeldt F, Hilton D, Pepe S, Krum H. Systematic review of effect of coenzyme Q10 in physical exercise, hypertension and heart failure. *Biofactors* 2003;18(1-4):91-100. DOI: 10.1002/biof.5520180211
19. Ylikoski T, Piirainen J, Hanninen O, Penttinen J. The effect of coenzyme Q10 on the exercise performance of cross-country skiers. *Mol Aspects Med* 1997;18(Suppl):S283-90. DOI: 10.1016/S0098-2997(97)00038-1
20. Blumberg JB, Frei BB, Fulgoni VL, Weaver CM, Zeisel SH. Impact of Frequency of Multi-Vitamin/Multi-Mineral Supplement Intake on Nutritional Adequacy and Nutrient Deficiencies in U.S. Adults. *Nutrients* 2017;9(8):849. DOI: 10.3390/nu9080849
21. Bailey RL, Fulgoni VL, 3rd, Keast DR, Dwyer JT. Examination of vitamin intakes among US adults by dietary supplement use. *J Acad Nutr Diet* 2012;112(5):657-63.e654. DOI: 10.1016/j.jand.2012.01.026
22. Linus Pauling Institute. Micronutrient supplements; 2018 [accessed 22 March 2021]. Available from: <https://lpi.oregonstate.edu/mic/micronutrient-inadequacies/remedy#micronutrient-supplements>
23. Haskell CF, Robertson B, Jones E, Forster J, Jones R, Wilde A, et al. Effects of a multi-vitamin/mineral supplement on cognitive function and fatigue during extended multi-tasking. *Hum Psychopharmacol* 2010;25(6):448-61. DOI: 10.1002/hup.1144
24. Kennedy D, Stevenson E, Jackson P, Dunn S, Wishart K, Bieri G, et al. Multivitamins and minerals modulate whole-body

energy metabolism and cerebral blood-flow during cognitive task performance: a double-blind, randomised, placebo-controlled trial. *Nutr Metab (Lond)* 2016;13:11. DOI: 10.1186/s12986-016-0071-4

25. Dodd FL, Kennedy DO, Stevenson EJ, Veasey RC, Walker K, Reed S, et al. Acute and chronic effects of multivitamin/mineral supplementation on objective and subjective energy measures. *Nutr Metab (Lond)* 2020;17(1):16. DOI: 10.1186/s12986-020-00435-1

Nutrición  
Hospitalaria




Fig. 1. Multiple micronutrients are required for efficient energy metabolism, and are essential to support every step of cellular respiration.

Suboptimal (<RDA) micronutrient intakes are common	<ul style="list-style-type: none"> <li>Even in industrialized countries, <b>micronutrient intakes are often insufficient</b><sup>1</sup></li> <li>Need to achieve RDA – <b>higher levels</b> may be required to ensure <b>optimum micronutrient levels</b><sup>2</sup></li> </ul>
A combination of all micronutrients is needed to generate energy	<ul style="list-style-type: none"> <li>Insufficient micronutrient intake could decrease enzymatic activity;<sup>2,3</sup></li> <li>Leads to <b>impaired energy metabolism</b> and inability to meet metabolic demands</li> <li>Resulting <b>tiredness, fatigue</b> and <b>low energy</b> affect 5-45% of the general population<sup>2-4</sup></li> <li>Even <b>mild micronutrient deficiencies in 'healthy' adults</b> can reduce energy levels and cause fatigue<sup>2</sup></li> </ul>
Suboptimal levels of B vitamins in particular affect wellbeing	<ul style="list-style-type: none"> <li>Deficiency is associated with cognitive impairment, depression, dementia<sup>5</sup></li> <li>Supplementing with multiple micronutrients can <b>improve general mood</b> and <b>wellbeing</b><sup>6-15</sup></li> </ul>
During the performance of cognitive tasks, overall energy expenditure increases <sup>17</sup>	<ul style="list-style-type: none"> <li>Micronutrients have roles in <b>brain function</b>:<sup>3</sup></li> <li><b>Directly</b>, via neurotransmitter synthesis, receptor binding, membrane ion pump function, etc.</li> <li><b>Indirectly</b>, via involvement in <b>energy metabolism</b>, modulation of cerebral blood supply, etc.</li> </ul>

Suboptimal micronutrients levels	Impact	Deficiencies in the general population <sup>1-18,19</sup>
Folic acid, vitamin C, iron, zinc, electrolytes	Fatigue	✓
Vitamin C, vitamins B1, B2, and B6, iron, zinc, calcium	Reduced physical fitness/performance	✓ (not vitamins B1 & B2 or calcium)
Vitamin C, iron, zinc, copper	Diminished work capacity	✓ (not copper)
Zinc	Decreased cardiorespiratory function	✓
Vitamin C	Impaired connective tissue, shortness of breath	✓
Vitamin C, iron, zinc, magnesium, electrolytes	Muscle weakness	✓
Vitamin C, folic acid	Anemia	✓
Iron	Wellbeing	✓ (particularly women)
Vitamin D	Increased mood disorders (e.g., seasonal affective disorder, depression)	✓
Iron	Mental development	✓ (including children, Elmadfa 2009)

**Rippin 2017 (Europe):** (country-dependent) deficiencies in vitamin D, folic acid, iron (mostly premenopausal-age females), calcium (especially older people), iodine, potassium (mostly females), sodium (some females only).  
**Elmadfa 2009 (Europe):** deficiencies in vitamins A, C, D, E, B2, B6, and B12, folic acid, iron, selenium, and zinc.  
**Cowan 2019 (USA):** deficiencies in vitamins A, B6, B12, C, D, E, and K, magnesium and potassium (males and females), iron (adolescent and premenopausal females), phosphorous, selenium, and zinc (males).  
RDA, recommended dietary allowance

- Elmadfa et al. *Forum Nutr* 2009;62:1-405
- Wishart. *Vitamin Miner* 2017;6:1-16
- Huskisson et al. *J Int Med Res* 2007;35(3):277-89
- Ricci et al. *J Occup Environ Med* 2007;49(11):1-10
- Kennedy. *Nutrients* 2016;8(2):68
- Kaplan et al. *Psychol Bull* 2007;133:747-60
- Haskell et al. *Hum Psychopharmacol* 2010; 25:448-61
- Kennedy et al. *Psychopharmacology (Berl)* 2010;211:55-68
- Harris et al. *Hum Psychopharmacol* 2011;26:560-7
- Stough et al. *Hum Psychopharmacol* 2011;26:470-6
- Pipingas et al. *Appetite* 2013;69:123-36
- White et al. *Nutrients* 2015;7:9005-17
- Macpherson et al. *Evid Based Complement Alternat Med* 2016;3092828
- Kennedy et al. *Hum Psychopharmacol* 2011;26:338-47
- Sarris et al. *Evid Based Complement Alternat Med* 2012;809653
- Jatunen et al. *Int J Food Sci Nutr* 2014;65:507-14
- Seematter et al. *Am J Physiol Endocrinol Metab* 2000;279:E799-805
- Rippin et al. *Nutrients* 2017;9(12)
- Cowan et al. *Nutrients* 2019;12(1)

Fig. 2. Multiple micronutrients are necessary to support metabolic energy production. Suboptimal nutrition, which can have adverse effects on energy, cognition, and wellbeing, is common even in industrialized countries.

	Maric 2014	Haskell 2010	Kennedy 2016	Dodd 2020
Study details	Prospective, open study ♀ CFS (n=38; 18–50 years) MMS (1 RDA) (2 months supplementation) SOD activity, self-rating scales	Double-blind RCT ♀ Healthy, occasional fatigue (n=216; 25–50 years) MMS (3RDA) vs. PBO (62 days supplementation) CT	Double-blind RCT ♀ Healthy (n=97; 25–49 years) MMS (1RDA) + CoQ10 or MMS (3RDA) vs. PBO (56 days supplementation) Assessment 60 min after first intake (acute) and on the day of the final intake (chronic) CT, calorimetry, cerebral blood flow	Double-blind RCT ♀ ♂ Healthy, physically active people (n=82; 18–35 years) MMS (3RDA) + CoQ10 vs. PBO (28 days supplementation) Assessment 45 min after first intake (acute) and a day after final intake (chronic) CT, PE
Stress	↓ SOD activity (oxidative stress)	—	NA	↓ acute & chronic During CT ♀
Mental and physical tiredness	↓ Fatigue, sleep disorders	↓ Physical	— Mental	↓ acute & chronic Physical ♀ Mental (during PE) ♀ ♂
Energy expenditure (kcal/min)	NA	NA	↑ acute & chronic (3RDA)	↑ acute & chronic During CT ♀ ♂ Carbohydrate (during PE) ♂
Cerebral blood flow	NA	NA	↑ acute Frontal cortex (1RDA + CoQ10)	NA
Fat or carbohydrate oxidation	NA	NA	↑ acute Fat (3RDA)	↑ acute & chronic During PE ♂
Performance of cognitive tasks/brain energy delivery	NA	↑ Alertness, accuracy, speed	— (either supplement)	NA
Micronutrient status	NA	↓ HCY (subgroup)	↑ chronic Vit D3, CoQ10      ↓ chronic HCY	↑ chronic Ferritin ♀ ♂
Adverse events reported	✗	✗	✗	✗

♀ female; ♂ male; — no significant change; CFS, chronic fatigue syndrome; CT, cognitive tasks; HCY, homocysteine; MMS, multiple micronutrient supplement; NA, not assessed; PBO, placebo; PE, physical exercise; RCT, randomized controlled trial; RDA, recommended daily allowance; SOD, superoxide dismutase.

Fig. 3. Overview of the effects of MMS ± CoQ10 (Supradyn) on factors of energy, cognition, and nutritional status.