



Original/*Obesidad*

Prevalence of overweight and obesity among secondary school children aged 14 to 18 years (China)

Lijun Zhu, Yan Chen, Lingling Ding, Daoxia Guo, Linghong Wang, Xiaohua Ren, Qijun Gu, Zhonghua Nie, Lianping He, Yuelong Jin and Yingshui Yao

*School of Public Health, Wannan Medical College. Wuhu, Anhui 241002 China.**

Abstract

Background: Prevalence of overweight and obesity in adolescents has been increasing during recent decades. The present study aimed to assess the prevalence of obesity in 14-18 years old Anhui Secondary school children, and provide more comprehensive guidelines for intervention programs.

Methods: This cross-sectional study was conducted among 4788 children in secondary school, and data from the routine health survey was analyzed. Trained health workers took anthropometric measures, height and weight, at the school. Definition of children overweight and obesity children was based on the international age- and sex-specific body mass index reference standards proposed by the International Obesity Task Force and National Center for Health Statistics.

Results: According to the IOTF standard the prevalence of overweight (including obesity) and obesity revealed decreasing trend with age in male (overweight, from 17.3% at 14-y to 5.3% at 18-y; obesity, from 3.8% at 14-y to 2.3% at 18-y). Depending on the NCHS references used, the overall prevalence of overweight (including obesity) and obesity in male aged 14 to 18 decreases from 16.4% to 11.7% and from 4.6% to 2.8%, respectively.

Conclusion: The overweight (including obesity) and obesity from 14 to 18 years old have a decreasing trend, and more attention should be taken to male students in China. These findings suggest secondary school and government need to design appropriate guidelines to keep healthy for adolescents in China.

(*Nutr Hosp.* 2015;31:2006-2010)

DOI:10.3305/nh.2015.31.5.8532

Key words: *Overweight. Obesity. Students. China.*

Correspondence: Yingshui Yao.
School of Public Health, Wannan Medical College.
No.22 Road Wenchangxi, Yijiang district,
Wuhu city, Anhui.
241002 China.
E-mail: yingshuiyao@163.com

Recibido: 26-XII-2014.
Aceptado: 13-I-2015.

**Lijun Zhu, Yan Chen and Lingling Ding are equal contribute to the work.*

PREVALENCIA DE SOBREPESO Y OBESIDAD ENTRE NIÑOS DE ESCUELA SECUNDARIA DE 14 A 18 AÑOS (CHINA)

Resumen

Antecedentes: La Prevalencia de sobrepeso y obesidad en adolescentes ha ido en aumento durante los últimos decenios. El presente estudio pretende evaluar la prevalencia de la obesidad en los niños de la escuela secundaria de 14 a 18 años de edad Anhui, y proporcionar directrices más amplias para los programas de intervención.

Métodos: Este estudio transversal se llevó a cabo en 4788 niños de escuela secundaria, y la fecha de la encuesta de salud de rutina fue analizado. Los trabajadores sanitarios capacitados tomaron medidas antropométricas, la altura y el peso, en la escuela. La definición de los niños con sobrepeso y obesidad basada en la edad y sexo del índice de masa corporal estándar de referencia propuesto por el Grupo de trabajo internacional sobre obesidad y el Centro Nacional de estadísticas de salud.

Resultados: De acuerdo a la IOTF Standard la prevalencia de sobrepeso (obesidad) y la obesidad reveló disminución tendencia con la edad en varones (sobrepeso, del 17,3% en 14-y a 5,3% en 18-y; la obesidad, from 3.8% en 14-y al 2,3% en 18-y). Dependiendo de la NCHS referencias utilizadas, la prevalencia global de sobrepeso (obesidad) y la obesidad en hombres de entre 14 y 18 años de edad disminuye a partir de un 16,4% a 11,7% y del 4,6% al 2,8%, respectivamente.

Conclusión: El sobrepeso (obesidad) y la obesidad de diciembre de los 14 a los 18 años de edad tienen una tendencia a la disminución, y más atención debe ser tomado para estudiantes varones en China. Estos hallazgos sugieren que la escuela secundaria y el Gobierno deben formular directrices apropiadas para mantener saludables a los adolescentes en China.

(*Nutr Hosp.* 2015;31:2006-2010)

DOI:10.3305/nh.2015.31.5.8532

Palabras clave: *Sobrepeso y obesidad. Estudiantes. China.*

Introduction

In recent decades, there were growing evidence of nutrition research and policy shift focused on overweight and obesity. Because of the established health risks and substantial increases in prevalence, obesity has become a major global health challenge¹. Many studies have documented the prevalence of overweight and obesity is growing progressively²⁻⁴. These previous studies point to obesity in adolescents is one of the principal health problems both in the developed and developing countries⁵⁻⁷, and obesity is also a risk factor for development of other diseases, such as cardiovascular complications⁸, diabetes⁹, hypertension¹⁰.

Some reports show that the prevalence of overweight among Chinese adolescents is increasing¹¹ and high, which in the urban population and particular in coastal big cities has reached the same level as that of some developed countries. However, the fact that the prevalence of obesity was low in most of the inland cities¹². In addition, environmental factors¹³, life-style preference¹⁴ and mental health¹⁵ play important roles in the rising prevalence of obesity worldwide. However, there is a paucity of recent data on the prevalence of obesity on Wannan area adolescents. The present study aimed to assess the prevalence of overweight and obesity in Anhui secondary school adolescents.

In this study, we desired to estimate the physical health of adolescents based on two references [the International Obesity Task Force standard (IOTF) and National Center for Health Statistics (NCHS)]

Methods

Subjects and methods

Participants

A cross-sectional study was performed among 6347 students in south of Anhui, China. Then we choose 4788 students who aged from 14 to 18 years old. This study consisted of 2360 male and 2428 female was recruited in 2013. Subjects were selected through multiple-cluster sampling. All subjects agreed to provide their personal information regarding the purpose and the procedures of our study, and written informed consent. Local ethics committee also approved and supported our study.

Anthropometric measurements

Weighing of the students was done, while they had light clothes on and they were bare-footed, measured to the nearest 0.1 kg on an electronic scale. The height of each student was also measured in standard manner, allowing for an error of 0.1 cm. All anthropometric data were collected by trained staff and supervised by

two school nurses. Body mass index was computed using the following standard equation: $BMI = \text{Weight} / \text{height squared in meter}$.

Definitions

Overweight and obesity are defined on BMI cut off points, which are gender and aged special. (1) The International Obesity Task Force standard are based on health related adult definitions of overweight ($\geq 25 \text{ kg/m}^2$) and obesity ($\geq 30 \text{ kg/m}^2$) but are adjusted to specific age and sex categories for adolescents. (2) The National Center for Health Statistics references are based on the different BMI at the age of 14-18 for classification of adolescents, respectively¹⁶.

Ethical consideration

The questionnaire was anonymous, and the participants were assured of data confidentiality. The protocol for human research of local and international guidelines on ethics was followed.

Statistical analysis

Descriptive statistic were used to characterize the sample in different age and sex by Excel software. A line graph was draw for the prevalence of overweight/obesity among subjects by age.

Results

The anthropometric characteristics of the sample, such as the mean (\pm SD) of weight, height, and calculated BMI are shown in Table I. A total of 4788 participants were evaluated in this study, 2360 male and 2428 female, aged 14-18 years. The prevalence of overweight and obesity among adolescents in China is described in Table II. The IOTF standard showed the trend in the prevalence of overweight (including obesity) and obesity tended to decrease with age in male. The prevalence of overweight (including obesity) among adolescent aged 14 to 18 year decreased from 17.3% to 5.3%, and the prevalence of obesity from 3.8% to 2.3%. Depending on the NCHS references used, the overall prevalence of overweight (including obesity) and obesity in male aged 14 to 18 decreases from 16.4% to 11.7% and from 4.6% to 2.8%, respectively. Using both international standards, the prevalence of overweight (including obesity) and obesity was fluctuant in female. Figure (1 y 2) showed that the overall proportion of overweight and obesity decreased as the age increased. The highest prevalence of overweight (17.3%) and obesity (4.6%) was reported at age 14.

Discussion

In the present study, we used two references [the International Obesity Task Force standard (IOTF) and National Center for Health Statistics (NCHS)], which are widely developed worldwide, to define the overweight

and obesity among secondary school students. However, the prevalence of obesity and overweight was lower than that of previous study¹⁷. An interesting observation was that the prevalence of overweight (including obesity) and obesity have a decrease with age in male, which is similar to the research by O,Musaiger A and Al-Mannai, M¹⁸.

Table I
Mean(±SD) of height, weight and BMI of secondary school students by age

		14	15	16	17	18
Male	N	452	720	815	316	57
	Height(cm)	169.15±7.20	172.51±5.90	173.92±5.75	173.85±5.16	175.02±4.70
	Weight(kg)	57.42±12.42	60.03±10.95	62.24±11.06	61.62±9.92	63.00±8.82
	BMI(kg/m ²)	19.96±3.59	20.13±3.32	20.53±3.18	20.36±3.04	20.55±2.66
Female	N	405	798	831	345	49
	Height(cm)	160.80±5.83	162.10±5.41	162.13±4.81	161.89±5.10	162.47±4.34
	Weight(kg)	50.83±9.07	50.96±7.26	51.34±7.48	51.45±7.20	52.18±8.07
	BMI(kg/m ²)	19.64±3.25	19.4±2.63	19.53±2.75	19.64±2.70	19.8±3.23




Fig. 1.—The prevalence of overweight for secondary school students by age (%).




Fig. 2.—The prevalence of obesity for secondary school students by age (%).

Table II
Prevalence of overweight and obesity in secondary school students by sex and age

Sexual	Reference		Prevalence (%)					Total
	Age		14	15	16	17	18	
Male	N		452	720	815	316	57	2360
	Overweight	NCHS	74(16.4)	92(12.8)	86(10.6)	22(7.0)	1(1.8)	275(11.7)
	Obesity	NCHS	21(4.6)	21(2.9)	18(2.2)	5(1.6)	0(0)	62(2.8)
	Overweight	IOTF	78(17.3)	106(14.7)	106(13.0)	30(9.5)	3(5.3)	323(13.7)
	Obesity	IOTF	17(3.8)	18(2.5)	15(1.8)	5(1.6)	0(0)	55(2.3)
Female	N		405	798	831	345	49	2428
	Overweight	NCHS	51(12.6)	38(4.8)	25(3.0)	12(3.5)	1(2.0)	127(5.2)
	Obesity	NCHS	11(2.7)	9(1.1)	13(1.6)	3(0.9)	1(2.0)	37(1.5)
	Overweight	IOTF	43(10.6)	36(4.5)	26(3.1)	14(4.1)	1(2.0)	120(4.9)
	Obesity	IOTF	6(1.5)	7(0.9)	10(1.2)	3(0.9)	1(2.0)	27(1.1)

Some factors that may be associated with the increasing prevalence of obesity among children and adolescent in China have been investigated. Chen, T.J et al found that spending more hours on watching television, playing computer games and surfing the internet were significantly associated with increased risk of obesity among Chinese youth¹⁹. Ahn, et al.²⁰ reported that nutrition transition was accelerating and the result of this trend was a rapid increase in obesity and chronic diseases.

Our finding of this study is similar to those reported in Italy, which reported that the trend in the prevalence of overweight (including obesity) among girls from 9-y to 15-y-old strongly was decreasing²¹. The possible reason maybe those secondary school children have exam stress and inadequate sleep time, and unhealthy eating behavior. Thus, further investigation should be taken to confirm whether there were relationship between exam stress and overweight. Our limitation maybe that samples size in 18 year old children is little small, and lack of data of type of learning, sleep time and test anxiety scale. We should collect more information in future study.

Our study showed that a comparison between two international standards to estimate for overweight and obesity among Anhui adolescents. A suitable defining selection of reference standard, therefore, should be considered when interpreting the prevalence of obesity among school children. The findings of this study raise the need for an intervention of school program to improve the health status and life equality of Chinese children and adolescents, especially in secondary school.

Conclusions

The study shows that the trend in overall prevalence of overweight (including obesity) and obesity is de-

creasing, especially in male should be given more attention in China. The study suggests secondary school and government need to design appropriate guidelines to keep healthy for adolescents in China.

Acknowledgement

This research was supported by the National Natural Science Foundation of China (81072367), the Anhui Provincial Natural Science Foundation (090413126 and 1308085MH135), the key Project University of Youth Talents fund of Anhui Province (2013SQRL-056ZD), and Wannan Medical key scientific research projects Engagement Fund (WK2013Z01)

Conflict of Interest

None declared

References

1. Ng, M., T. Fleming, M. Robinson, B. Thomson, N. Graetz, C. Margono, et al., Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: A systematic analysis for the global burden of disease study 2013. *Lancet* 2014. 384(9945): 766-81.
2. Castetbon, K., [recent prevalence of child and adolescent overweight and obesity in france and abroad.]. *Arch Pediatr* 2014.
3. Kotanidou, E.P., M.G. Grammatikopoulou, B.E. Spiliotis, C. Kanaka-Gantenbein, M. Tsigga and A. Galli-Tsinopoulou, Ten-year obesity and overweight prevalence in greek children: A systematic review and meta-analysis of 2001-2010 data. *Hormones (Athens)* 2013. 12(4): 537-49.
4. Taheri, F., T. Kazemi, T. Chahkandi, K. Namakin, M. Zardast and B. Bijari, Prevalence of overweight, obesity and central obesity among elementary school children in birjand, east of iran, 2012. *J Res Health Sci* 2013. 13(2): 157-61.

5. Wang, Y., C. Monteiro and B.M. Popkin, Trends of obesity and underweight in older children and adolescents in the united states, brazil, china, and russia. *Am J Clin Nutr* 2002. 75(6): 971-7.
6. Jan Mohamed, H.J., R.W. Yap, S.L. Loy, S.A. Norris, R. Biesma and J. Aagaard-Hansen, Prevalence and determinants of overweight, obesity, and type 2 diabetes mellitus in adults in malaysia. *Asia Pac J Public Health* 2014.
7. He, L., X. Ren, Y. Qian, Y. Jin, Y. Chen, D. Guo, et al., Prevalence of overweight and obesity among a university faculty and staffs from 2004 to 2010 in wuhu, china. *Nutr Hosp* 2014. 29(5): 1033-7.
8. Lavie, C.J., P.A. McAuley, T.S. Church, R.V. Milani and S.N. Blair, Obesity and cardiovascular diseases: Implications regarding fitness, fatness, and severity in the obesity paradox. *J Am Coll Cardiol* 2014. 63(14): 1345-54.
9. Garg, S.K., H. Maurer, K. Reed and R. Selagamsetty, Diabetes and cancer: Two diseases with obesity as a common risk factor. *Diabetes Obes Metab* 2014. 16(2): 97-110.
10. Vernooij, J.W., Y. van der Graaf, F.L. Visseren, W. Spiering and S.s. group, The prevalence of obesity-related hypertension and risk for new vascular events in patients with vascular diseases. *Obesity (Silver Spring)* 2012. 20(10): 2118-23.
11. Seo, D.C. and J. Niu, Trends in underweight and overweight/obesity prevalence in chinese youth, 2004-2009. *Int J Behav Med* 2014. 21(4): 682-90.
12. Ji, C.Y. and C. Cooperative Study on Childhood Obesity: Working Group on Obesity in, The prevalence of childhood overweight/obesity and the epidemic changes in 1985-2000 for chinese school-age children and adolescents. *Obes Rev* 2008. 9 Suppl 1: 78-81.
13. Velasquez-Melendez, G., L.L. Mendes and C.M. Padez, Built environment and social environment: Associations with overweight and obesity in a sample of brazilian adults. *Cad Saude Publica* 2013. 29(10): 1988-96.
14. Grundy, S.M., Multifactorial causation of obesity: Implications for prevention. *Am J Clin Nutr*, 1998. 67(3 Suppl): 563S-72S.
15. Lazarevich, I., M.E. Irigoyen-Camacho and C. Velazquez-Alva Mdel, Obesity, eating behaviour and mental health among university students in mexico city. *Nutr Hosp* 2013. 28(6): 1892-9.
16. Cole, T.J., M.C. Bellizzi, K.M. Flegal and W.H. Dietz, Establishing a standard definition for child overweight and obesity worldwide: International survey. *BMJ* 2000. 320(7244): 1240-3.
17. Ji, C.Y., J.L. Sun and T.J. Chen, [dynamic analysis on the prevalence of obesity and overweight school-age children and adolescents in recent 15 years in china]. *Zhonghua Liu Xing Bing Xue Za Zhi* 2004. 25(2): 103-8.
18. A, O. M., M. Al-Mannai and R. Tayyem, Prevalence of overweight and obesity among female adolescents in jordan: A comparison between two international reference standards. *Pak J Med Sci* 2013. 29(2): 678-9.
19. Chen, T. J., B. Modin, C.Y. Ji and A. Hjern, Regional, socioeconomic and urban-rural disparities in child and adolescent obesity in china: A multilevel analysis. *Acta Paediatr* 2011. 100(12): 1583-9.
20. Ahn, S., H. Zhao, M.L. Smith, M.G. Ory and C.D. Phillips, Bmi and lifestyle changes as correlates to changes in self-reported diagnosis of hypertension among older chinese adults. *J Am Soc Hypertens* 2011. 5(1): 21-30.
21. Lazzeri, G., S. Rossi, A. Pammolli, V. Pilato, T. Pozzi and M.V. Giacchi, Underweight and overweight among children and adolescents in tuscany (italy). Prevalence and short-term trends. *J Prev Med Hyg* 2008. 49(1): 13-21.