



Original/*Pediatría*

Energy expenditure and intake comparisons in Chilean children 4-5 years attending day-care centres

Gabriela Salazar¹, Fabián Vásquez¹, María P. Rodríguez¹, Ana M. Andrade¹, María A. Anziani¹, Fernando Vio¹ and Williams Coward²

¹Institute of Nutrition and Food Technology (INTA), University of Chile, Santiago (Chile). ²MRC Human Nutrition Research, Cambridge, United Kingdom.

Abstract

Introduction: the doubly labelled water (DLW) method has an accuracy of 1% and within-subject precision of 5-8%, depending on subject's age and environments issues. Energy intake assessment is prone to errors (>15-20%) depending in the method utilized.

Objective: to quantify DLW methodology errors in four to five year olds that could affect the comparison with energy intake.

Methods: energy expenditure (TEE, by DLW), was assessed during 14 days in 18 preschool children, who attended eight hours daily to day-care centres. Energy intake was determined by a combined method: food weighing during weekdays and recall after leaving the Centre (17h to sleep time) plus 24 h recall, during the weekend. Several assumptions affecting DLW total error were assessed to determine their influence in the comparison to energy intake (i.e. background variability, space ratio, proportion of water subject to fractionation, food quotient value).

Results: the individual mean energy expenditure was $1\,373 \pm 177$ kcal and the energy intake ($1\,409 \pm 161$ kcal). The overall difference between intake and expenditure was 42.9 kcal/day (limits of agreement + 259.1 to -112.3 kcal/day). TEE measurement error only explained a minor quantity (2.4%), between both measurements, and the observed mean isotope dilution space was 1.030 ± 0.010 confirming the value utilized in adults studies.

Conclusions: energy expenditure data is similar to other studies in preschool children. The small difference found between energy intake and expenditure may be attributed to the applied energy intake methodology, the homogeneous diet at care centres during the week-days and the lower DLW methodology error.

(Nutr Hosp. 2015;32:1067-1074)

DOI:10.3305/nh.2015.32.3.9263

Key words: *Doubly labelled water. Energy expenditure. Energy intake. Preschool children. Day-care centres.*

Correspondence: Gabriela Salazar.
Institute of Nutrition and Food Technology (INTA),
University of Chile, El Libano 5524, Santiago (Chile).
Email gsalazar@inta.uchile.cl

Recibido: 13-V-2015.

Aceptado: 28-VI-2015.

COMPARACIÓN ENTRE EL GASTO Y EL CONSUMO DE ENERGÍA EN NIÑOS CHILENOS DE 4-5 AÑOS ASISTENTES A JARDINES INFANTILES

Resumen

Introducción: el método del agua doblemente marcada (ADM) tiene una precisión del 1% y en un mismo sujeto es de 5-8%, dependiendo de la edad y el entorno del sujeto. La evaluación de la ingesta energética es propensa a errores (> 15-20%), dependiendo del método utilizado.

Objetivo: cuantificar los errores metodológicos del ADM en niños de 4-5 años que podrían afectar la comparación con la ingesta de energía.

Métodos: el gasto de energía (GTE, por ADM), se evaluó durante 14 días en 18 preescolares, asistentes a guarderías infantiles. La ingesta energética se determinó mediante un método combinado: pesaje de alimentos durante los días de la semana y registro después de salir del centro (17 horas en adelante), además de un recordatorio de 24 horas, durante un día del fin de semana.

Resultados: el promedio individual del gasto energético total fue 1373 ± 177 kcal y la ingesta de energía (1409 ± 161 kcal). La diferencia global entre la ingesta y el gasto fue 42,9 kcal/día. El error de medición del GET explicó una variación del 2,4%, entre ambas mediciones, y el espacio de dilución de isótopos fue 1030 ± 0.010 , confirmando el valor utilizado en los estudios de adultos.

Conclusiones: los datos de GET fueron similares a otros estudios realizados en niños en edad preescolar. La pequeña diferencia encontrada entre la ingesta y el gasto energético se puede atribuir a la metodología de la ingesta de energía aplicada, la dieta homogénea en los centros de atención, durante los días de la semana, y el bajo error metodológico del ADM.

(Nutr Hosp. 2015;32:1067-1074)

DOI:10.3305/nh.2015.32.3.9263

Palabras claves: *Agua doblemente marcada. Gasto energético. Ingesta de energía. Preescolares. Guarderías infantiles.*

Abbreviations

DLW: doubly labelled water.
TEE: total energy expenditure.
FQ: Food Quotient.

Introduction

Near five hundred thousand children receive food at school (twice) or day-care centre (three times), in Chile. The doubly labelled water method (DLW) has been extensively used for the measurement of total energy expenditure (TEE) since the 1980's¹⁻⁶ and much of the data is now incorporated into the formulation of human energy requirements⁷. In Chile, DLW methodology has been used to measure energy requirements in both infants⁸ and preschool children^{9,10,11}.

It is reported that in general, the DLW method has a relative accuracy of 1% and within-subject precision of 5-8%¹² but errors of the method should be assessed in the age groups and environments in which it is applied. In Chile, TEE has been measured in infants and preschool age children⁸⁻¹⁰ as these age group are relevant in public health nutrition due to the increasing prevalence of obesity in children¹³⁻¹⁵. Validated measurement of energy intake in preschool children is also clearly important for the development of policy recommendations to help prevent obesity from early age¹⁶. As known TEE must equal energy intake in individuals to achieve energy balance, thus the technique may be used to validate the assessment of energy intake by food weighing or questionnaires¹⁷.

In most reports, energy intake is underestimated whatever the method used for intake measurements^{18,19}; Kipnis *et al.* found that the 24 hours food recall method may underestimate energy intake by 12-14% in adult population^{20,21}. However, in studies where a person, other than the subject, is responsible for recording dietary intake, energy intake may often correspond to DLW determined energy expenditure²². In the present study, we analyze some of the errors derived from the approximations and assumptions in the DLW method (involving isotopic background variation, goodness of fit of the data to the model, isotope dilution spaces, fractionation and respiratory quotient), in the context of a comparison between energy intake and energy expenditure in Chilean preschool children, who attend day-care centres.

Materials and methods

Subjects

The study was carried out on 15 children of low socio-economic status aged 3-5 years. They were selected from public day-care centres in Santiago, Chile, according to the following criteria: birth weight 3000-4000 g., normal nutritional status, no congenital disease and

without medication or illness at the time of the study. Children entered the study when their parents had been informed about the experimental protocol and had given written consent. The Ethical Committee of the Institute of Nutrition and Food Technology approved the study. A description of the subjects is provided in table I.

DLW technique

To test the effect of isotope dose two different doses were administered: dose H (high), 0.2 g/kg deuterium oxide (99.9%) and 2.0 g/kg 10% H₂¹⁸O for nine children, and dose L (low), 0.08 g/kg deuterium oxide (99.9%) and 1.55 g/kg 10% H₂¹⁸O for six children. To assess the significance of baseline variation, seven urine samples of 6 ml were collected daily for a week, after dosing. One urine sample was taken every day, for eight consecutive days. Urine samples were immediately sealed in individual sterile tubes and refrigerated at -18 to -22 °C.

Hydrogen was equilibrated with water in the presence of a Pt-Al catalyst at 20 °C for the deuterium analysis²³. Oxygen-18 labelled samples were equilibrated with 4% CO₂ in He and placed in a thermo-regulated bath at 25 °C for 15 hours²⁴. After equilibration, isotopic content of the samples was measured in a HYDRA IRMS mass spectrometer (Europa Scientific, Crewe, United Kingdom). Measurement precision was 0.4-0.6 ppm for deuterium analysis and 0.2-0.3 ppm for oxygen-18. Slopes and intercepts of the regressions of log enrichment above base line versus time were calculated with a computer program adjusting least minimal squares (LMS) to the data. Propagation of errors in the measurement of CO₂ production was calculated from Cole TJ *et al.* and Ritz P *et al.*^{25,26}.

Uncorrected CO₂ production rate was corrected by isotope fraction, assuming that a constant proportion of total water losses were fractionated²⁷. After this correction, the CO₂ production rate was converted into energy expenditure by assuming a fixed respiratory quotient or a calculate one²⁸.

Anthropometry

The children were weighed, at the day care-centres, the day that the DLW protocol started, then at mid time (4-5 days) and finally on the 13-14 days using a standardised SECA scale (model H W 60 with scale up to 60 kg and precision of 0.01 g, Vogel and Halke, Hamburg, FRG).

Dietary intake

Food weighing

Food intake was measured twice during the TEE study period by standard procedures: by weighing all

Table I
Description of subjects participating in the study

Subjects	Sex ¹	Age (y)	Weight (Kg)	Height (cm)	BMP ² (Kg/m ²)	ZWH ³	ZHA ⁴	Birth Weight (g)	Birth Height (cm)
<i>Higher Dose</i>									
1	M	5.2	20.2	110.3	16.7	0.8	-0.2	3.5	50.5
2	M	4.8	18.6	106.6	16.4	0.6	-0.6	3.9	53.0
3	F	5.4	19.1	111.3	15.4	0.2	0.1	3.2	51.0
4	F	5.1	15.8	102.1	15.1	-0.1	-1.7	3.2	47.0
5	F	4.1	16.0	102.3	15.3	0.0	-0.1	3.8	49.0
6	M	3.8	14.8	96.1	16.0	0.1	-1.0	3.7	52.0
7	F	4.2	17.1	103.0	16.1	-0.1	0.1	3.1	48.5
8	M	3.3	15.2	99.3	15.4	-0.3	0.6	3.2	54.0
9	F	3.3	13.4	91.7	15.9	0.1	-1.1	3.5	50.0
Mean		4.4	16.7	102.5	15.8	0.1	-0.4	3.5	50.6
SD		0.8	2.2	6.4	0.5	0.4	0.7	0.3	2.2
<i>Lower dose</i>									
10	M	5.2	19.4	109.8	16.1	0.4	-0.5	3.1	45.6
11	M	4.9	18.3	106.3	16.2	0.4	-0.7	3.3	51.0
12	M	5.0	17.3	105.6	15.6	0.0	-1.0	3.2	49.0
13	M	5.1	16.7	106.5	14.7	-0.6	-1.1	3.4	51.0
14	M	3.6	18.0	104.8	16.4	0.6	1.4	3.2	50.0
15	M	3.9	15.9	97.5	16.7	0.5	-0.9	3.5	52.0
Mean		4.6	17.6	105.1	16.0	0.2	-0.5	3.3	49.8
SD		0.7	1.2	4.1	0.7	0.4	0.9	0.2	2.3

¹Sex: M: Males. F: Females. ²BMI: body mass index. ³ZWH: Z-score Weight for Height. ⁴ZHA: Z-score Height for Age.

the food ingested by the child at the day-care centres (breakfast, lunch and snack). All food was weighed before being eaten by the child, and any left overs were weighed again. A trained nutritionist using a Scaltec scale (Hamburg, Germany) with a precision of 0.1 g performed the weighing.

Recall data

On the same days that food was weighed at the day care-centres, a dietary recall was used for breakfast and the remaining periods (after 17:00 hours). The mother recorded in a notebook all the food ingested by the children at home. Next morning, the nutritionist in charge interviewed the mother to determine the type of food, portion sizes and preparation of the food. The same procedure was repeated during one day of the weekend (either Saturday or Sunday) and then on the next Monday morning. The data obtained from the food intake recall at home, plus the food ingested at the day care-centre and determined by food weighing

was combined in a single week weighed average daily value. Finally, the weekly food energy intake was calculated from the dietary data using in house food intake software with typical Chilean foods values²⁹.

Respiratory quotient

The respiratory quotient of the diet (FQ) was calculated from the measured dietary intake data as described by Black AE *et al.*³⁰.

Statistical Methods

Descriptive statistics including minimum, maximum, and frequency distribution were derived for all variables. The Shapiro Wilk goodness-of-fit test and homogeneity of variance test were performed for continuous variables. Normal variables were expressed as mean \pm standard deviation. Student's t test was used to test for significance of differences.

Subsequently, the results were compared using CO₂ production fractionation and TEE (kcal) and per kg of weigh, using individual FQ (TEEind) versus TEEfix using RQ value = 0.85, with repeated measures (parametric analysis). The significance level was set at p < 0.05. Data was analysed using STATA 12.0 (StataCorp. 2011. Stata Statistical Software: Release 12. College Station, TX: StataCorp LP.).

Results

Assumptions effects

Natural abundance variation in the correlation between ²H and ¹⁸O and its relation to the relative amounts of the isotope doses given, contribute to imprecision in the final CO₂ production estimate^{31,32}. In this study, seven background samples collected on successive days prior to the DLW study were measured and errors arising from this source were calculated related to the goodness of fit of the data to the model²⁶.

The percentage contribution of each of these errors to CO₂ production rate and their combined values are shown in table II.

The component of error associated with background variation tended to be relatively larger (3.76%) for the low dose group compared to 2.99% in the high dose. But neither this, the goodness of fit error or the total error was significantly different between groups (Table II).

It is now usual in calculations to use values for the isotope dilution spaces that are normalised to population means. This is justified if analytical error contributes most to this component of overall variance²⁵ and of course the mean value used, typically 1,0297, needs to correspond to the true value for the study population. The effect of deviations in the values in the ratio of measured space from deuterium dilution to oxygen-18 dilution, compared to an externally derived, assumed population mean (1,0297) is shown in table III. The observed mean isotope dilution space was 1,030 ± 0.010, no different from the customarily assumed population, thus mean CO₂ production values

Table II
Background variability error covariance of both isotopic background values.
Total error of the fit and overall error in 16 subjects

Subject	CO ₂ (moles)	Background variability error (%)	Goodness of fit error (%)	Total error (%)
<i>Higher Dose</i>				
1	10,76	1,11	2,97	3,18
2	13,34	1,45	1,55	2,13
3	10,74	2,64	3,69	4,54
4	10,01	1,35	3,76	4
5	9,79	1,53	2,48	2,92
6	13,02	4,43	6,62	7,97
7	10,97	2,98	4,83	5,68
8	11,09	5,05	3,43	6,1
9	9,89	3,71	4,07	5,51
Mean	11,1	2,7	3,7	4,7
SD	1,3	1,5	1,4	1,8
<i>Lower dose</i>				
10	15,87	2,26	1,21	2,56
11	14,05	6,61	2,47	7,05
12	12,49	1,18	3,81	3,99
13	12,1	1,87	3,22	3,72
14	12,3	5,09	2,68	5,75
15	9,68	5,97	2,85	6,62
Mean	12,7	3,8	2,7	4,9
SD	2,1	2,3	0,9	1,8

calculated by each of these procedures produces no relative bias (Table III).

The effects of smaller (0,20) or greater (0,40) proportion of water assumed to be subject to fractionation as compared to the value of 0,30 applied in this and other studies on children²⁷ are shown in table IV. The effect of the magnitude of fractionation assumptions largely depends upon the relative difference between the slopes of the isotope disappearance curves (the smaller the difference the greater the effect). In practice, larger differences will be found when water turnover rates are high. Thus, the magnitude of effects recorded in one set of circumstances (country, climate etc.) cannot be applied in others. In this study, the effects were small. An overestimate of 1,8% in CO₂ was found when a factor of 0,20 is utilized, and a lower value of CO₂ (-1,9%), if the opposite situation is assumed (0,40).

Assessment of subject's diet produced a different value from the typical utilized in many studies³⁰. For example, in Europe, a factor of 0,85 has been obtained from individual diet results^{18,19}. This present study, mean FQ value was also 0,85. Thus when mean energy expenditures were calculated using the fixed value of 0,85 (TEE_{fix}) for FQ and the individually determined FQ (TEE_{ind}), the values were the same (Table V).

Comparison of TEE and energy intake

The total energy intake was calculated as a weighted average of weekday and weekend energy intake. The individual mean energy expenditure (1373+177 kcal) and energy intake (1409+161 kcal) are shown in table VI.

Discussion

DLW methodology has been used as a gold standard reference method to assess the validity of energy intake. In general, the measurement of energy intake underestimates the real needs, although the difference found in children under six years of age, is lower compared to older children. In fact, this difference increases over age and sex, affecting the outcome and it is worse in girls³³⁻³⁵.

In the present study, energy intake in children was evaluated by weighing all food ingested by the child at the day-care centre and a dietary recall at home. The overall difference between intake and expenditure was 42,9 kcal/day (limits of agreement + 259,1 to -112,3 kcal/day).

At the individual level, absolute agreement between intake and expenditure would not be expected from

Table III
Effects of fractionated water proportion (0,2, 0,3 and 0,4)^a on CO₂ production

Subject	CO ₂ _{0.2} (mols)	CO ₂ _{0.3} (mols)	CO ₂ _{0.4} (mols)
1	10.01	9.82	9.63
2	12.55	12.31	12.12
3	10.68	10.44	10.22
4	9.31	9.13	8.96
5	8.99	8.76	8.53
6	12.27	12.01	11.83
7	10.11	9.87	9.63
8	10.16	9.89	9.62
9	9.14	8.94	8.74
10	14.91	14.70	14.49
11	13.18	13.00	12.81
12	11.67	11.48	11.29
13	11.35	11.11	10.92
14	11.45	11.16	10.92
15	8.96	8.77	8.57
Mean	10.98	10.76	10.55
SD	1.73	1.73	1.73

^aP < 0,05 Repeated measures ANOVA.

Table IV
Observed dilution spaces ratio and values of CO₂ production for observed (o) ratio and normalized ratio (n)

Subject	Observed Ratio	CO ₂ _o mole/d	CO ₂ _n mole/d	CO ₂ _o /CO ₂ _n
1	1.029	10.8	10.79	1.001
2	1.038	12.31	12.66	0.972
3	1.046	9.65	10.44	0.924
4	1.047	8.99	10.01	0.90
5	1.040	8.76	9.24	0.948
6	1.029	13.21	12.71	1.039
7	1.013	12.30	11.56	1.064
8	1.032	11.48	11.59	0.991
9	1.017	11.11	10.58	1.050
10	1.024	11.42	11.17	1.022
11	1.028	11.27	11.17	1.009
12	1.033	9.89	10.06	0.983
13	1.026	8.31	8.16	1.018
14	1.021	12.24	11.79	1.038
15	1.023	9.7	9.43	1.029
Mean	1.030	10.76	10.76	0.999
SD	0.010	1.48	1.26	0.047

Table V
Effects of fractionated water proportion (0,2, 0,3 and 0,4)^a on normalized CO₂ production

Subject	CO _{2 0,2} (mole/d)	CO _{2 0,3} (mole/d)	CO _{2 0,4} (mole/d)
1	11.14	10.79	10.76
2	12.84	12.66	12.47
3	10.68	10.44	10.20
4	10.79	10.62	10.45
5	9.47	9.24	9.01
6	12.99	12.78	12.56
7	11.75	11.56	11.36
8	11.78	11.59	11.39
9	10.76	10.58	10.39
10	11.36	11.17	10.98
11	11.41	11.17	10.94
12	10.33	10.06	9.79
13	8.36	8.16	7.96
14	12.02	11.79	11.56
15	9.62	9.43	9.24
Mean	11.02	10.80	10.60
SD	1.25	1.25	1.25

^aP < 0,05 Repeated measures ANOVA.

a single measurement; but at a group level, the measurement of TEE should validate the measurement of habitual energy intake¹⁸. In a recent review, a similar comparisons in adult studies reported biases ranging from -59 to +24% but in most cases the intake measurement underestimated expenditure³⁶.

In infants and small children, data is less consistent. Lanigan *et al.*, observed a non-significant bias (+7,3%) for weighed intake compared to DLW of in infants of 6-24 months old^{37,38}; Davies in 1,5-4,5 year old children found a non-significant bias of 4% in the comparison of energy intake and expenditure³⁹. Livingstone found that utilising diet history in children of 3-5 years old, produced a 14% significant overestimation at 3 years of age and 11% at 5 years, compared to energy expenditure²⁷.

The validity of the average agreement between the measures in the present study is supported by our calculations on the relative importance of RQ and fractionation assumptions. By extension of the calculations used for table V it can be shown that to increase the mean TEE by 2,4% to match the mean intake would require the use of RQ=0,874 rather than 0,85. Similarly, the amount of water fractionated would need to decrease to 16% in order to explain a 2,4% of the difference. Reported data on children this age elsewhere does not substantiate this value²⁷.

The correlation of intake and expenditure is adequate (r=0,68) which agrees with other authors^{37,39,40}.

Table VI
TEE (kcal) and per kg of weight using individual FQ (TEE_{ind}^a) versus TEE_{fix}^b, using fixed RQ value = 0,85

Subject	Calc FQ	TEE _{ind} ^a (kcal/d)	TEE _{ind} ^b kcal/kg/d	TEE _{fix} ^a (kcal/d)	TEE _{fix} ^b kcal/kg/d	Energy intake ^c week	Energy intake ^c weekend	Energy intake weighed
1	0.88	1337	66.3	1373	68.1	1235	1214	1272
2	0.87	1539	79.8	1566	81.2	1513	1678	1560
3	0.80	1195	61.2	1227	62.9	1604	1080	1454
4	0.87	1247	79.9	1269	78.3	1311	1359	1324
5	0.82	1146	70.3	1114	69.0	1395	1099	1310
6	0.87	1650	83.1	1680	84.6	1685	1420	1609
7	0.87	1536	84.8	1564	86.3	1485	1274	1424
8	0.87	1434	80.5	1460	81.9	1244	1903	1432
9	0.86	1401	82.4	1413	83.2	1255	1449	1310
10	0.77	1568	105.5	1452	97.7	1750	1441	1662
11	0.86	1420	82.9	1433	83.7	1678	1163	1531
12	0.86	1247	81.8	1258	82.6	1377	1129	1306
13	0.85	1057	79.2	1057	79.2	1142	1241	1170
14	0.85	1557	86.5	1557	86.5	1565	1739	1615
15	0.83	1256	78.9	1234	77.5	1337	1050	1255
Mean	0.85	1373	80.2	1377	80.2	1438	1349	1416
SD	0.03	177	9.9	180	8.5	189	258	152

^{a,b,c}P < 0,05 Repeated measures ANOVA.

Although the difference range between individual differences can vary from -7% to +10% from the real value, the adequacy of these results may be attributed to energy intake method and the homogeneous diet that preschool children ate during the week-days at the day care-centres.

Irrespective of this potential bias, we also examined the relationship between the individual errors predicted in table II. TEE measurement error only explains a minor quantity of the 2,4% value between both measurements, thus TEE error would appear to have a small effect on this difference.

In summary, the main factor influencing the comparison between TEE and energy intake could be strongly related to energy intake measurement factors. In this study the errors in intake measurement could only come from the home recordings, where food intake is more variable, particularly at weekends³³. In the Framingham Study the association between energy intake in the 3-4 year old age group, predicted the intake in the subsequent period (5-6 year olds) where 57,1% to 85,7% of the children remained in the energy intake top two quintiles. Sixty percent of the 93,3% children with the highest intake at baseline (3-4 year olds) were still in the top two quintiles at 7-8 years of age³⁵. Thus, the validity of energy intake assessment methods is a very important tool for the study and prevention of childhood weight excess. In this paper, the comparison to DLW to a mixed energy intake methodology prove to be successful and may be applicable to similar places where subjects eat near 75% of their daily energy intake. As a general conclusion, energy intake determination is crucial and requires to be valid especially at preschool age, due to its significant association to the later presence of obesity and future interventions^{41,42}.

Acknowledgements

We thank the participation of the children, their parents and the day-care centres for the facilities provided for this study. The INTA Stable Isotopes Laboratory specially thanks Dr. Andy Coward, who gave scientific and technical support for many years and is present today although no longer with us.

The National Council for Research, Science and Technology (Chile), under grant FONDECYT 1010692 and FONDECYT Postdoctoral 3140344.

Conflict of Interest

None the authors has a conflict of interest.

References

1. Ball EJ, O'Connor J, Abbott R, Steinbeck KS, Davies PS, Wishart C, *et al.* Total energy expenditure, body fatness,

- and physical activity in children aged 6-9 y. *Am J Clin Nutr* 2001;74:524-528.
2. Schoeller DA. Recent advances from application of doubly labeled water to measurement of human energy expenditure. *J Nutr* 1999;129:1765-1768.
3. Coward WA. Contributions of the doubly labeled water method to studies of energy balance in the Third World. *Am J Clin Nutr* 1998;68:962S-969S.
4. Davies PS. Energy requirements for growth and development in infancy. *Am J Clin Nutr* 1998;68: 939S-943S.
5. Goran MI, Sun M. Total energy expenditure and physical activity in prepubertal children: recent advances based on the application of the doubly labeled water method. *Am J Clin Nutr* 1998;68:944S-949S.
6. DeLany JP, Harsha DW, Kime JC, Kumler J, Melancon L, Bray GA. Energy expenditure in lean and obese prepubertal children. *Obes Res* 1995;3:67-72.
7. FAO/WHO/UNU. Human Energy Requirements. Report of a Joint FAO/WHO/UNU, 2001. Expert Consultation. Roma.
8. Salazar G, Vio F, Garcia C, Aguirre E, Coward WA. Energy requirements in Chilean infants. *Arch Dis Child Fetal Neonatal* 2000;83:F120-F123.
9. Vásquez F, Salazar G, Andrade M, Vásquez L, Díaz E. Energy balance and physical activity in obese children attending day-care centres. *Eur J Clin Nutr* 2006;60:1115-1121.
10. Velasquez M, Salazar G, Vio F, Díaz Z N, Anziani G A. Validation of equations to assess body composition using anthropometric data in Chilean preschool children. *Rev Med Chile* 2008;136:433-441.
11. Butte NF, Wong WW, Lee JS, Adolph AL, Puyau MR, Zakeri IF. Prediction of energy expenditure and physical activity in preschoolers. *Med Sci Sports Exerc* 2001;4:1216-1226.
12. Schoeller DA. Validation of habitual energy intake. *Public Health Nutr* 2002;5:883-888.
13. Chile. National Department of Health (MINSAL). [Online]. [2011] [cited June 06, 2014]. Available from: URL:<http://www.minsal.cl>
14. Kain J, Uauy R, Vio F, Albala C. Trends in overweight and obesity prevalence in Chilean children: comparison of three definitions. *Eur J Clin Nutr* 2002;56:200-204.
15. Albala C, Vio F, Kain J, Uauy R. Nutrition transition in Latin America: the case of Chile. *Nutr Rev* 2001;59:170-176.
16. DeLany JP. Role of energy expenditure in the development of pediatric obesity. *Am J Clin Nutr* 1998;68:950S-955S.
17. Schoeller DA. Uses of stable isotopes in the assessment of nutrient status and metabolism. *Food Nutr Bull* 2002;23:17-20.
18. Black AE, Prentice AM, Goldberg GR. Measurements of total energy expenditure provide insights into the validity of dietary measurements of energy intake. *J Am Diet Assoc* 1993;93:572-579.
19. Black AE & Cole T. Within –and between– subject variation in energy expenditure measured by the doubly-labelled water technique: implications for validating reported dietary energy intake. *Eur J Clin Nutr* 2000;54:386-394.
20. Kipnis V, Subar AF, Midthune D, Freedman LS, Ballard-Barbash R, Troiano RP, *et al.* Structure of dietary measurement error: results of the OPEN biomarker study. *Am J Epidemiol* 2003;158:14-21.
21. Subar AF, Dodd KW, Guenther PM, Kipnis V, Midthune D, McDowell M, *et al.* The food propensity questionnaire: concept, development, and validation for use as a covariate in a model to estimate usual food intake. *J Am Diet Assoc* 2006;106:1556-1563.
22. Hill RJ, Davies PS. The validity of self-reported energy intake as determined using the doubly labelled water technique. *Br J Nutr* 2001;85:415-430.
23. Scrimgeour CM, Rollo MM, Mudambo SM, Handley LL, Prosser SJ. A simplified method for deuterium/hydrogen isotope ratio measurements on water samples of biological origin. *Biol Mass Spectrom* 1993; 22:383-387.
24. Prosser SJ, Brookes ST, Linton A, Preston T. Rapid, automated analysis of ¹³C and ¹⁸O of CO₂ in gas samples by continuous-flow, isotope ratio mass spectrometry. *Biol Mass Spectrom* 1991;20:724-730.

25. Coward WA, Ritz P, Cole TJ. Revision of calculations in the doubly labelled water method for measurement of energy expenditure in humans. *Am J Physiol* 1994;267:E805-E807.
26. Cole TJ, Coward WA. Precision and accuracy of doubly labeled water energy expenditure by multipoint and two-point methods. *Am J Physiol* 1992;263:E965-E973.
27. Livingstone MB, Prentice AM, Coward WA, Strain JJ, Black AE, Davies PS, *et al.* Validation of estimates of energy intake by weighed dietary record and diet history in children and adolescents. *Am J Clin Nutr* 1992;56:29-35.
28. Elia M. Energy equivalents of CO₂ and their importance in assessing energy expenditure when using tracer techniques. *Am J Physiol* 1991;260:E75-E88.
29. Schmidt-Hebbel H, Pennachiotti I. Chemical Composition of Chilean Foods, 1985. University of Chile. 6th edition.
30. Black AE, Prentice AM, Coward WA. Use of food quotients to predict respiratory quotients for the doubly-labelled water method of measuring energy expenditure. *Hum Nutr Clin Nutr* 1986;40: 381-391.
31. Ritz P, Cole TJ, Couet C, Coward WA. Precision of DLW energy expenditure: contributions of natural abundance variations. *Am J Physiol* 1996;270:E164-E169.
32. Ritz P, Cole TJ, Davies PSW, Goldberg GR, Coward WA. Interactions between 2H and 18O natural abundance variations and DLW measurements of energy expenditure. *Am J Physiol* 1996;271: E302-E308.
33. Sepp H, Lennernas M, Pettersson R, Abrahamsson L. Children's nutrient intake at preschool and at home. *Acta Paediatr* 2001;90:483-491.
34. Sichert-Hellert W, Kersting M, Manz F. Changes in time-trends of nutrient intake from fortified and non-fortified food in German children and adolescents-15 year results of the DONALD study. Dortmund Nutritional and Anthropometric Longitudinally Designed Study. *Eur J Nutr* 2001;40:49-55.
35. Singer MR, Moore LL, Garrahe EJ, Ellison RC. The tracking of nutrient intake in young children: the Framingham Children's Study. *Am J Public Health* 1995;85:1673-1677.
36. Trabulsi J, Troiano RP, Subar AF, Sharbaugh C, Kipnis V, Schatzkin A, *et al.* Precision of the doubly labeled water method in a large-scale application: evaluation of a streamlined-dosing protocol in the Observing Protein and Energy Nutrition (OPEN) study. *Eur J Clin Nutr* 2003;57:1370-1377.
37. Lanigan JA, Wells JC, Lawson MS, Lucas A. Validation of food diary method for assessment of dietary energy and macronutrient intake in infants and children aged 6-24 months. *Eur J Clin Nutr* 2001;55:124-129.
38. Lanigan JA, Wells JC, Lawson MS, Cole TJ, Lucas A. Number of days needed to assess energy and nutrient intake in infants and young children between 6 months and 2 years of age. *Eur J Clin Nutr* 2004;58:745-750.
39. Davies PS, Coward WA, Gregory J, White A, Mills A. Total energy expenditure and energy intake in the preschool child: a comparison. *Br J Nutr* 1994;72:13-20.
40. McGloin AF, Livingstone MB, Greene LC, Webb SE, Gibson JM, Jebb SA, *et al.* Energy and fat intake in obese and lean children at varying risk of obesity. *Int J Obes Relat Metab Disord* 2002;26:200-207.
41. Deheeger M, Akrouf M, Bellisle F, Rossignol C, Rolland-Cachera MF. Individual patterns of food intake development in children: a 10 months to 8 years of age follow-up study of nutrition and growth. *Physiol Behav* 1996;59:403-407.
42. Salazar G, Vasquez F, Concha F, Rodríguez MP, Berlanga MR, Rojas J. *et al.* Pilot nutrition and physical activity intervention for preschool children attending daycare centres (JUNJI); primary and secondary outcomes. *Nutr Hosp* 2014;29:1004-1012.