



Original/*Valoración nutricional*

Subscapular and triceps skinfolds reference values of Hispanic American children and adolescents and their comparison with the reference of Centers for Disease Control and Prevention (CDC)

María Dolores Marrodán¹, Marisa González-Montero de Espinosa¹, Ángel Herráez^{1,2}, Emma Laura Alfaro³, Ignacio Felipe Bejarano³, María Margarita Carmenate⁴, Consuelo Prado⁴, Delia Beatriz Lomaglio⁵, Noemí López-Ejeda², Antonio Martínez⁶, María Soledad Mesa¹, Betty Méndez Pérez⁷, Juana María Meléndez⁸, Susana Moreno Romero¹, José Luis Pacheco¹, Vanesa Vázquez⁶ and José E. Dipierri³

¹Grupo de Investigación Epinut. Departamento de Zoología y Antropología Física. Facultad de Ciencias Biológicas. Universidad Complutense de Madrid, España. ²Departamento de Biología de Sistemas, Universidad de Alcalá, España. ³Instituto de Biología de la Altura. Universidad Nacional de Jujuy, Argentina. ⁴Departamento de Biología. Facultad de Ciencias. Universidad Autónoma de Madrid, España. ⁵Centro de Estudios de Antropología Biológica. Facultad de Ciencias Exactas y Naturales. Universidad Nacional de Catamarca, Argentina. ⁶Museo de Antropología Montané. Facultad de Biología, Universidad de la Habana, Cuba. ⁷Instituto de Investigaciones Económicas y Sociales. Universidad Central de Venezuela. ⁸Centro de Investigación en Alimentación y Desarrollo (CIAD), Hermosillo, México.

Abstract

Introduction: the assessment of the skinfold thickness is an objective measure of adiposity. Therefore, it is a useful tool for nutritional diagnosis and prevention of metabolic risk associated with excess fat in childhood and adolescence.

Objective: to provide percentiles of subscapular and triceps skinfolds for Hispanic American schoolchildren and compare them with those published by the Centers for Disease Control and Prevention (CDC) from United States, that it have been commonly used as a reference in most of these countries.

Methods: subscapular and triceps skinfolds were measured in 9.973 schoolchildren 4-19 aged from Spain, Argentina, Cuba, Venezuela and Mexico with Holtain caliper with 0.2 mm accuracy. Percentiles were obtained with the LMS statistical method and were presented in tables divided in stages of 6 months and in curves graphics. The difference between Hispanic American and CDC mean values were provided for P3, P50 and P97 in mm and also were graphically represented.

Results: skinfolds measurements obviously increased with age in both sexes but, in boys, this increase is much more marked in highest percentiles between 8 and 13 years; this maximum is reached earlier than what occurs in CDC reference. In both sexes, all percentiles

VALORES DE REFERENCIA PARA LOS PLIEGUES ADIPOSOS SUBESCAPULAR Y TRICIPITAL DE NIÑOS Y ADOLESCENTES HISPANOAMERICANOS Y SU COMPARACIÓN CON LA REFERENCIA DE LOS CENTROS PARA EL CONTROL Y PREVENCIÓN DE ENFERMEDADES (CDC)

Resumen

Introducción: la evaluación del grosor de los pliegues subcutáneos es una medida objetiva de la adiposidad. Es por tanto una herramienta útil para el diagnóstico nutricional y la prevención del riesgo metabólico asociado al exceso de grasa en la infancia y adolescencia.

Objetivo: proporcionar valores percentilares de los pliegues adiposos subcutáneos subescapular y tricipital para escolares hispanoamericanos y compararlos con los valores publicados por los Centros para el Control y Prevención de Enfermedades (CDC) de los Estados Unidos, que comúnmente se emplean como referencia en estos países.

Métodos: se midió el pliegue subescapular y tricipital en 9.973 escolares entre 4 y 19 años procedentes de España, Argentina, Cuba, Venezuela y México con un calibre Holtain de 0,2 mm de precisión. Los percentiles fueron calculados mediante el método estadístico LMS y presentados en tablas divididas en intervalos de seis meses y en gráficos de curvas. La diferencia entre los valores medios hispanoamericanos y los valores del CDC se muestran para el P3, P50 y P97 en mm, y también gráficamente.

Resultados: las medidas de los pliegues subcutáneos se incrementan obviamente con la edad pero, en niños, este incremento es mucho más marcado en los percentiles superiores entre los 8 y 13 años; este máximo es alcanzado antes que en la referencia del CDC. En ambos sexos, todos los percentiles analizados fueron superiores en los escolares hispanoamericanos, exceptuando el P97 por

Correspondence: María Dolores Marrodán Serrano. U.D. Antropología Física, Facultad de Ciencias Biológicas, Universidad Complutense de Madrid, España. c/ José Novais 12, planta 8, despacho 5. CP. 28040 Madrid. E-mail: marrodan@ucm.es

Recibido: 15-VIII-2015.
Aceptado: 6-IX-2015.

analyzed in Hispanic American schoolchildren were higher than the CDC reference except P97 up to 10 or 13 years that was notably smaller.

Conclusions: the skinfolds percentiles of Hispanic American children and adolescents differ from CDC that are usually used as reference. The values of subscapular and triceps skinfolds provided in this study, could be applied to populations of a similar ethnic background, especially in comparative studies of body composition.

(*Nutr Hosp.* 2015;32:2862-2873)

DOI:10.3305/nh.2015.32.6.9775

Key words: *Adiposity. Anthropometry. Spain. Latin-america. Childhood. Adolescence.*

Abbreviations

CDC: Centers for Disease Control and Prevention.

BMI: Body Mass Index.

IOTF: International Obesity Task Force.

NHANES: National Health and Nutrition Examination Survey.

WHO: World Health Organization.

Introduction

The prevalence of obesity shows a positive secular trend in all age groups in developed or industrialized and developing countries in all continents^{1,2}. Based on the trend of obesity observed in America between 1985 and 2005, a prevalence of 15% obesity and 40% of overweight² was estimated for 2010 on this continent. In developing countries, as is the case in many Latin American countries, obesity –in addition– coexists with poverty, malnutrition with overweight or double burden of malnutrition, setting a new nutritional paradigm³.

However, significant interpopulation differences in the prevalence of obesity⁴, the distribution of adiposity by sex before and after puberty⁵ and density of fat-free mass were detected which have been interpreted as conditioned by ethnicity⁶. Interethnic differences have also been observed between diverse assessment methods in human body composition, such as bioelectrical impedance analysis and dual-energy X-ray absorptiometry⁷ and in the distribution of the visceral and subcutaneous fatty compartments of the body, variations that may be attenuated or masked by obesity⁸. It is therefore important to define the extent of adiposity in children from different ethnic groups.

The validity of the Body Mass Index (BMI) as an indicator of body fat and cardiometabolic risk in young population has also been questioned because it has limitations in distinguishing fat from lean mass⁹. Due to its low cost and non-invasive procedure, evaluation of skinfolds is one of the most objective anthropometric measurements to assess adiposity for its high compatibility with other methods of direct measure-

ment of body fat mass¹⁰. Excess adiposity assessed by skinfolds successfully associates in adolescents with increased blood lipids (triglycerides and cholesterol) and insulin resistance, markers for increased risk of metabolic syndrome¹¹.

Conclusiones: los percentiles de pliegues adiposos de los niños y adolescentes hispanoamericanos difieren de la referencia del CDC. Los valores del pliegue subescapular y tricéptico proporcionados en este estudio podrían ser aplicados en poblaciones de similar origen étnico, especialmente en estudios comparativos de la composición corporal.

(*Nutr Hosp.* 2015;32:2862-2873)

DOI:10.3305/nh.2015.32.6.9775

Palabras clave: *Adiposidad. Antropometría. España. Latinoamérica. Infancia. Adolescencia.*

ment of body fat mass¹⁰. Excess adiposity assessed by skinfolds successfully associates in adolescents with increased blood lipids (triglycerides and cholesterol) and insulin resistance, markers for increased risk of metabolic syndrome¹¹.

Between 2005 and 2008 a multicenter collaborative project was conducted to collect anthropometric data in children and adolescents from different localities of Spain, Venezuela, Argentina, Cuba and Mexico. Based on these data, skinfold centiles of biceps, triceps, subscapular and suprailiac folds were calculated using the LMS method.

Objective

In order to assess the extent of interethnic differences in adiposity in large-scale studies, the aim of the present study was to provide percentiles of subscapular and triceps skinfolds for Hispanic American young people and compare them with the values obtained from the Centers for Disease Control and Prevention (CDC)¹² elaborated for United States children and adolescents that commonly have been used as a reference in most of these Hispanic American countries.

Methods

Origin and composition of the sample

The sample consisted of 9973 healthy students (4964 boys, 5009 girls) between 4 and 19 years old, without obvious pathologies at the time of measurement, attending public schools of middle and low socioeconomic level at different locations in Argentina (Catamarca and Jujuy), Cuba (Havana), Spain (Madrid), Mexico (Hermosillo), and Venezuela (Caracas and Merida).

The participants' date of birth was obtained from their national identity document or the School Registry and the decimal age was calculated¹³. Data were grouped by gender and in 31 age groups with an interval of half a year.

Measurements

After obtaining informed consent from parents or guardians and abiding by the rules of Helsinki¹⁴, anthropometric measurements were performed between 2005-2008 by trained personnel, with approved instruments and in accordance with techniques recommended by the International Biological Programme¹⁵.

Body weight (kg) was measured with a lever balance (100 g precision) in light clothing, and height (cm) with a vertical anthropometer (1 mm accuracy). Based on these measurements, BMI was calculated and using the criteria of the International Obesity Task Force (IOTF) developed by Cole *et al.*^{16,17}. The prevalence of thinness, overweight and obesity were determined by sex in the following age groups: 4-8 years, 9-12 years and > 12 years.

Skinfolds were measured with a Holtain constant pressure adipometer and 2 tenths of millimeter accuracy. Instruments were calibrated at the beginning of each anthropometric session. With the subject's arm extended and relaxed, the skinfold was taken at the mesobraquial region in the acromial-radial midline with the thumb of our left hand a pinch of skin and adipose tissue over the triceps not including muscle tissue. The subscapular skinfold was taken by holding the adipose tissue at the inferior angle of the scapula at its vertebral border, obliquely downward and outward at an angle of 45° with the horizontal line passing through the inferior border of the scapula¹⁸.

Statistical Analysis

The dispersion of raw data was analyzed and outliers were deleted using as cutoff ± 4 SD; this criterion eliminated 84 cases. The LMS method was applied to calculate percentiles, as it summarizes the changing distribution of the anthropometric measurements according to age using the L, M and S curves representing the skewness (λ), median (μ) and coefficient of variation (σ), respectively. The LMS method uses the Box-Cox transformation to adjust the distribution of anthropometric data to a normal distribution, essentially minimizing the effects of asymmetry¹⁹. The L, M and S parameters were calculated according to the method of penalized maximum likelihood¹⁹. The values of L, M and S centiles were used to calculate according to the following formula^{20,21}:

$$C = M * 1 * LSZ * 1/L$$

where L, M and S were the values calculated for each age and Z was the corresponding percentile needed (3, 5, 10, 25, 50, 75, 90, 95 and 97). The data processing was performed using the LMS ChartMaker Pro (The Institute of Child Health, London) software²². Q tests for fit were used to assess the global goodness of fit of models.

The 3, 50 and 97 percentiles of triceps and subscapular skinfold thickness were graphically compared with the respective values of the CDC reference¹² also calculated using the same LMS method.

To examine the discrepancy in the calculated percentiles by age and sex, with respect to the CDC reference¹², differences in millimeters and percentage were calculated using the following formula²¹:

$$100 \log (\text{centile of the reference} / \text{calculated centile})$$

Results

Table I shows the nutritional status of the Hispanic American sample according to the classification criteria of the IOTF. As can be seen, while the proportion of normally nourished schoolchildren remained constant along the ontogenetic period analyzed, thinness decreased as increased the percentage of students who were included in the categories of overweight or obese.

Figures 1 (a,b) and 2 (a,b) represent the percentiles of subscapular and triceps folds for girls and boys. The




Fig. 1.a.—Subscapular skinfold thickness percentiles representation for Hispanic American girls from 4 to 19 years.




Fig. 1.b.—Subscapular skinfold thickness percentiles representation for Hispanic American boys from 4 to 19 years.

Table I
Nutritional status distribution of the sample by age group

| | Age (years) | | | Total |
|-------------------|-------------|------|------|-------|
| | 4-8 | 9-12 | >12 | |
| Thinness (%) | 8.3 | 6.7 | 6.8 | 7.1 |
| Normal weight (%) | 71.7 | 69.2 | 69.9 | 70.0 |
| Overweight (%) | 14.5 | 18.5 | 19.1 | 17.9 |
| Obesity (%) | 5.5 | 5.6 | 4.2 | 4.9 |

numerical values of these measurements and LMS parameters are shown in tables II to V for both sexes and for age groups ranges of 6 months from 4 to 19 years old. Subscapular skinfolds measurements increased with age in both sexes but, in boys, this increase is much more marked between 8 and 13 years in top percentiles (90, 95 and 97).

Compared to the CDC reference this increase is reached earlier and then it experiences a decline as from 13 years in both sexes (Figs. 3a, 3b). A similar trend is observed in triceps skinfold in which the extreme

Table II
Subscapular skinfold percentiles and L, M and S values of hispanic american girls

| Age (years) | P3 | P5 | P10 | P25 | P50 | P75 | P90 | P95 | P97 | L | M | S |
|-------------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|------|
| 4 | 3.96 | 4.21 | 4.65 | 5.60 | 7.12 | 9.54 | 13.30 | 17.01 | 20.52 | -0.74 | 7.12 | 0.39 |
| 4.5 | 3.86 | 4.11 | 4.55 | 5.50 | 7.04 | 9.48 | 13.27 | 17.02 | 20.55 | -0.71 | 7.04 | 0.40 |
| 5 | 3.77 | 4.02 | 4.47 | 5.42 | 6.97 | 9.44 | 13.28 | 17.05 | 20.61 | -0.69 | 6.97 | 0.41 |
| 5.5 | 3.71 | 3.96 | 4.40 | 5.37 | 6.94 | 9.44 | 13.34 | 17.16 | 20.75 | -0.67 | 6.94 | 0.41 |
| 6 | 3.66 | 3.92 | 4.38 | 5.36 | 6.96 | 9.52 | 13.49 | 17.39 | 21.02 | -0.64 | 6.96 | 0.42 |
| 6.5 | 3.66 | 3.92 | 4.39 | 5.40 | 7.05 | 9.69 | 13.78 | 17.77 | 21.48 | -0.62 | 7.05 | 0.43 |
| 7 | 3.71 | 3.98 | 4.46 | 5.51 | 7.23 | 9.98 | 14.23 | 18.35 | 22.16 | -0.59 | 7.23 | 0.44 |
| 7.5 | 3.80 | 4.09 | 4.59 | 5.70 | 7.50 | 10.40 | 14.85 | 19.14 | 23.07 | -0.57 | 7.50 | 0.44 |
| 8 | 3.91 | 4.21 | 4.74 | 5.91 | 7.81 | 10.85 | 15.51 | 19.96 | 24.00 | -0.54 | 7.81 | 0.45 |
| 8.5 | 4.01 | 4.32 | 4.88 | 6.10 | 8.09 | 11.27 | 16.09 | 20.64 | 24.75 | -0.52 | 8.09 | 0.45 |
| 9 | 4.11 | 4.44 | 5.02 | 6.30 | 8.37 | 11.67 | 16.61 | 21.23 | 25.34 | -0.49 | 8.37 | 0.45 |
| 9.5 | 4.23 | 4.57 | 5.19 | 6.52 | 8.68 | 12.09 | 17.14 | 21.79 | 25.87 | -0.47 | 8.68 | 0.45 |
| 10 | 4.37 | 4.73 | 5.38 | 6.77 | 9.03 | 12.54 | 17.68 | 22.32 | 26.33 | -0.44 | 9.03 | 0.45 |
| 10.5 | 4.54 | 4.92 | 5.59 | 7.05 | 9.40 | 13.02 | 18.21 | 22.81 | 26.72 | -0.41 | 9.40 | 0.45 |
| 11 | 4.72 | 5.12 | 5.83 | 7.36 | 9.79 | 13.50 | 18.71 | 23.24 | 27.02 | -0.38 | 9.79 | 0.45 |
| 11.5 | 4.91 | 5.33 | 6.07 | 7.67 | 10.19 | 13.96 | 19.16 | 23.58 | 27.21 | -0.35 | 10.19 | 0.44 |
| 12 | 5.10 | 5.54 | 6.32 | 7.98 | 10.56 | 14.37 | 19.52 | 23.80 | 27.26 | -0.32 | 10.56 | 0.44 |
| 12.5 | 5.27 | 5.73 | 6.54 | 8.25 | 10.89 | 14.71 | 19.76 | 23.88 | 27.15 | -0.28 | 10.89 | 0.43 |
| 13 | 5.42 | 5.89 | 6.73 | 8.48 | 11.15 | 14.96 | 19.88 | 23.81 | 26.89 | -0.25 | 11.15 | 0.42 |
| 13.5 | 5.53 | 6.02 | 6.87 | 8.65 | 11.34 | 15.12 | 19.89 | 23.63 | 26.53 | -0.21 | 11.34 | 0.41 |
| 14 | 5.60 | 6.10 | 6.97 | 8.78 | 11.48 | 15.20 | 19.83 | 23.39 | 26.11 | -0.18 | 11.48 | 0.41 |
| 14.5 | 5.65 | 6.16 | 7.04 | 8.88 | 11.58 | 15.25 | 19.74 | 23.13 | 25.69 | -0.14 | 11.58 | 0.40 |
| 15 | 5.68 | 6.20 | 7.10 | 8.96 | 11.67 | 15.30 | 19.66 | 22.92 | 25.35 | -0.10 | 11.67 | 0.40 |
| 15.5 | 5.71 | 6.24 | 7.16 | 9.05 | 11.77 | 15.38 | 19.65 | 22.78 | 25.10 | -0.06 | 11.77 | 0.39 |
| 16 | 5.73 | 6.28 | 7.22 | 9.14 | 11.89 | 15.49 | 19.67 | 22.71 | 24.94 | -0.02 | 11.89 | 0.39 |
| 16.5 | 5.75 | 6.31 | 7.28 | 9.23 | 12.01 | 15.60 | 19.72 | 22.68 | 24.82 | 0.02 | 12.01 | 0.39 |
| 17 | 5.75 | 6.32 | 7.32 | 9.31 | 12.12 | 15.70 | 19.75 | 22.63 | 24.70 | 0.06 | 12.12 | 0.39 |
| 17.5 | 5.73 | 6.32 | 7.34 | 9.37 | 12.20 | 15.77 | 19.75 | 22.55 | 24.54 | 0.11 | 12.20 | 0.39 |
| 18 | 5.71 | 6.32 | 7.36 | 9.43 | 12.28 | 15.83 | 19.74 | 22.45 | 24.38 | 0.15 | 12.28 | 0.38 |
| 18.5 | 5.70 | 6.32 | 7.39 | 9.49 | 12.37 | 15.91 | 19.75 | 22.39 | 24.25 | 0.19 | 12.37 | 0.38 |
| 19 | 5.69 | 6.33 | 7.43 | 9.57 | 12.47 | 16.00 | 19.78 | 22.35 | 24.15 | 0.24 | 12.47 | 0.38 |




Fig. 2.a.—Triceps skinfold thickness percentiles representation for Hispanic American girls from 4 to 19 years.




Fig. 2.b.—Triceps skinfold thickness percentiles representation for Hispanic American boys from 4 to 19 years.

Table III
Subscapular skinfold percentiles and L, M and S values of hispanic american boys

| Age (years) | P3 | P5 | P10 | P25 | P50 | P75 | P90 | P95 | P97 | L | M | S |
|-------------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|------|
| 4 | 3.55 | 3.72 | 4.01 | 4.60 | 5.49 | 6.79 | 8.62 | 10.25 | 11.67 | -0.95 | 5.49 | 0.29 |
| 4.5 | 3.55 | 3.72 | 4.03 | 4.66 | 5.63 | 7.09 | 9.18 | 11.10 | 12.83 | -0.92 | 5.63 | 0.31 |
| 5 | 3.54 | 3.73 | 4.05 | 4.73 | 5.79 | 7.40 | 9.78 | 12.04 | 14.11 | -0.89 | 5.79 | 0.33 |
| 5.5 | 3.54 | 3.74 | 4.08 | 4.81 | 5.95 | 7.73 | 10.44 | 13.07 | 15.54 | -0.86 | 5.95 | 0.35 |
| 6 | 3.56 | 3.77 | 4.13 | 4.91 | 6.15 | 8.11 | 11.16 | 14.20 | 17.13 | -0.83 | 6.15 | 0.37 |
| 6.5 | 3.60 | 3.81 | 4.20 | 5.03 | 6.38 | 8.54 | 11.96 | 15.46 | 18.88 | -0.80 | 6.38 | 0.39 |
| 7 | 3.65 | 3.88 | 4.29 | 5.18 | 6.64 | 9.01 | 12.84 | 16.81 | 20.76 | -0.77 | 6.64 | 0.40 |
| 7.5 | 3.71 | 3.96 | 4.40 | 5.35 | 6.92 | 9.52 | 13.77 | 18.23 | 22.70 | -0.74 | 6.92 | 0.42 |
| 8 | 3.78 | 4.04 | 4.51 | 5.53 | 7.22 | 10.04 | 14.69 | 19.62 | 24.59 | -0.71 | 7.22 | 0.44 |
| 8.5 | 3.85 | 4.12 | 4.62 | 5.70 | 7.50 | 10.53 | 15.55 | 20.88 | 26.24 | -0.68 | 7.50 | 0.45 |
| 9 | 3.90 | 4.19 | 4.71 | 5.84 | 7.75 | 10.96 | 16.28 | 21.92 | 27.56 | -0.65 | 7.75 | 0.46 |
| 9.5 | 3.95 | 4.25 | 4.78 | 5.97 | 7.96 | 11.32 | 16.87 | 22.70 | 28.50 | -0.62 | 7.96 | 0.47 |
| 10 | 3.99 | 4.30 | 4.86 | 6.09 | 8.16 | 11.63 | 17.34 | 23.28 | 29.11 | -0.60 | 8.16 | 0.47 |
| 10.5 | 4.04 | 4.36 | 4.94 | 6.21 | 8.35 | 11.93 | 17.75 | 23.72 | 29.48 | -0.57 | 8.35 | 0.48 |
| 11 | 4.10 | 4.43 | 5.03 | 6.34 | 8.54 | 12.19 | 18.06 | 23.98 | 29.60 | -0.55 | 8.54 | 0.48 |
| 11.5 | 4.17 | 4.51 | 5.12 | 6.47 | 8.72 | 12.41 | 18.25 | 24.02 | 29.41 | -0.53 | 8.72 | 0.48 |
| 12 | 4.24 | 4.59 | 5.22 | 6.59 | 8.86 | 12.55 | 18.29 | 23.84 | 28.92 | -0.50 | 8.86 | 0.47 |
| 12.5 | 4.31 | 4.66 | 5.30 | 6.69 | 8.97 | 12.62 | 18.18 | 23.44 | 28.16 | -0.48 | 8.97 | 0.47 |
| 13 | 4.37 | 4.73 | 5.38 | 6.77 | 9.05 | 12.63 | 17.97 | 22.89 | 27.22 | -0.46 | 9.05 | 0.46 |
| 13.5 | 4.45 | 4.81 | 5.46 | 6.86 | 9.12 | 12.62 | 17.71 | 22.29 | 26.23 | -0.44 | 9.12 | 0.45 |
| 14 | 4.53 | 4.90 | 5.56 | 6.97 | 9.21 | 12.63 | 17.47 | 21.71 | 25.29 | -0.42 | 9.21 | 0.44 |
| 14.5 | 4.65 | 5.02 | 5.69 | 7.11 | 9.34 | 12.69 | 17.30 | 21.25 | 24.51 | -0.39 | 9.34 | 0.43 |
| 15 | 4.79 | 5.17 | 5.86 | 7.30 | 9.54 | 12.83 | 17.26 | 20.96 | 23.96 | -0.36 | 9.54 | 0.42 |
| 15.5 | 4.96 | 5.36 | 6.06 | 7.54 | 9.81 | 13.08 | 17.38 | 20.89 | 23.69 | -0.33 | 9.81 | 0.41 |
| 16 | 5.14 | 5.56 | 6.29 | 7.82 | 10.12 | 13.39 | 17.60 | 20.97 | 23.61 | -0.29 | 10.12 | 0.40 |
| 16.5 | 5.32 | 5.76 | 6.52 | 8.09 | 10.44 | 13.73 | 17.86 | 21.10 | 23.60 | -0.26 | 10.44 | 0.39 |
| 17 | 5.48 | 5.93 | 6.72 | 8.34 | 10.74 | 14.04 | 18.11 | 21.23 | 23.61 | -0.21 | 10.74 | 0.39 |
| 17.5 | 5.59 | 6.06 | 6.88 | 8.55 | 11.00 | 14.30 | 18.30 | 21.31 | 23.58 | -0.17 | 11.00 | 0.38 |
| 18 | 5.68 | 6.17 | 7.01 | 8.73 | 11.21 | 14.51 | 18.43 | 21.35 | 23.51 | -0.12 | 11.21 | 0.38 |
| 18.5 | 5.74 | 6.24 | 7.11 | 8.87 | 11.39 | 14.68 | 18.53 | 21.34 | 23.41 | -0.07 | 11.39 | 0.37 |
| 19 | 5.78 | 6.30 | 7.20 | 9.00 | 11.54 | 14.83 | 18.61 | 21.32 | 23.30 | -0.02 | 11.54 | 0.37 |

Table IV
Triceps skinfold percentiles and L, M and S values of hispanic american girls

| Age (years) | P3 | P5 | P10 | P25 | P50 | P75 | P90 | P95 | P97 | L | M | S |
|-------------|------|------|------|-------|-------|-------|-------|-------|-------|------|-------|------|
| 4 | 5.40 | 5.94 | 6.87 | 8.69 | 11.17 | 14.23 | 17.57 | 19.87 | 21.50 | 0.23 | 11.17 | 0.37 |
| 4.5 | 5.17 | 5.71 | 6.62 | 8.41 | 10.87 | 13.90 | 17.20 | 19.48 | 21.09 | 0.23 | 10.87 | 0.37 |
| 5 | 4.96 | 5.49 | 6.39 | 8.16 | 10.59 | 13.59 | 16.87 | 19.12 | 20.72 | 0.23 | 10.59 | 0.38 |
| 5.5 | 4.78 | 5.30 | 6.19 | 7.95 | 10.37 | 13.35 | 16.61 | 18.85 | 20.43 | 0.23 | 10.37 | 0.38 |
| 6 | 4.65 | 5.17 | 6.06 | 7.82 | 10.23 | 13.22 | 16.48 | 18.72 | 20.30 | 0.24 | 10.23 | 0.39 |
| 6.5 | 4.58 | 5.10 | 5.99 | 7.77 | 10.21 | 13.22 | 16.51 | 18.78 | 20.37 | 0.24 | 10.21 | 0.39 |
| 7 | 4.58 | 5.11 | 6.02 | 7.83 | 10.32 | 13.40 | 16.76 | 19.07 | 20.70 | 0.24 | 10.32 | 0.40 |
| 7.5 | 4.66 | 5.20 | 6.15 | 8.01 | 10.59 | 13.76 | 17.23 | 19.60 | 21.28 | 0.24 | 10.59 | 0.40 |
| 8 | 4.78 | 5.34 | 6.32 | 8.26 | 10.92 | 14.21 | 17.78 | 20.24 | 21.97 | 0.24 | 10.92 | 0.40 |
| 8.5 | 4.90 | 5.49 | 6.50 | 8.51 | 11.25 | 14.64 | 18.31 | 20.82 | 22.59 | 0.24 | 11.25 | 0.40 |
| 9 | 5.03 | 5.64 | 6.68 | 8.74 | 11.56 | 15.03 | 18.77 | 21.33 | 23.13 | 0.24 | 11.56 | 0.40 |
| 9.5 | 5.16 | 5.79 | 6.86 | 8.98 | 11.86 | 15.39 | 19.18 | 21.77 | 23.58 | 0.25 | 11.86 | 0.40 |
| 10 | 5.30 | 5.94 | 7.04 | 9.21 | 12.14 | 15.72 | 19.55 | 22.15 | 23.96 | 0.26 | 12.14 | 0.40 |
| 10.5 | 5.44 | 6.10 | 7.23 | 9.44 | 12.42 | 16.03 | 19.88 | 22.48 | 24.29 | 0.27 | 12.42 | 0.39 |
| 11 | 5.59 | 6.27 | 7.42 | 9.68 | 12.71 | 16.35 | 20.21 | 22.80 | 24.61 | 0.29 | 12.71 | 0.39 |
| 11.5 | 5.74 | 6.44 | 7.63 | 9.93 | 13.00 | 16.68 | 20.55 | 23.14 | 24.94 | 0.30 | 13.00 | 0.38 |
| 12 | 5.90 | 6.61 | 7.82 | 10.17 | 13.29 | 17.00 | 20.89 | 23.49 | 25.28 | 0.32 | 13.29 | 0.38 |
| 12.5 | 6.03 | 6.77 | 8.01 | 10.40 | 13.57 | 17.31 | 21.22 | 23.81 | 25.61 | 0.33 | 13.57 | 0.38 |
| 13 | 6.14 | 6.89 | 8.16 | 10.60 | 13.81 | 17.58 | 21.50 | 24.10 | 25.89 | 0.34 | 13.81 | 0.37 |
| 13.5 | 6.22 | 6.98 | 8.28 | 10.75 | 13.99 | 17.80 | 21.73 | 24.32 | 26.10 | 0.36 | 13.99 | 0.37 |
| 14 | 6.26 | 7.04 | 8.35 | 10.86 | 14.13 | 17.95 | 21.88 | 24.47 | 26.24 | 0.38 | 14.13 | 0.37 |
| 14.5 | 6.28 | 7.07 | 8.39 | 10.93 | 14.22 | 18.05 | 21.98 | 24.56 | 26.32 | 0.39 | 14.22 | 0.37 |
| 15 | 6.28 | 7.08 | 8.43 | 10.99 | 14.30 | 18.15 | 22.07 | 24.64 | 26.40 | 0.41 | 14.30 | 0.37 |
| 15.5 | 6.29 | 7.11 | 8.47 | 11.06 | 14.40 | 18.27 | 22.20 | 24.76 | 26.52 | 0.42 | 14.40 | 0.37 |
| 16 | 6.30 | 7.13 | 8.51 | 11.13 | 14.50 | 18.39 | 22.33 | 24.90 | 26.65 | 0.43 | 14.50 | 0.37 |
| 16.5 | 6.30 | 7.13 | 8.53 | 11.18 | 14.58 | 18.49 | 22.45 | 25.02 | 26.77 | 0.44 | 14.58 | 0.37 |
| 17 | 6.29 | 7.13 | 8.55 | 11.22 | 14.65 | 18.58 | 22.55 | 25.13 | 26.88 | 0.45 | 14.65 | 0.37 |
| 17.5 | 6.28 | 7.13 | 8.56 | 11.26 | 14.72 | 18.68 | 22.67 | 25.26 | 27.02 | 0.46 | 14.72 | 0.37 |
| 18 | 6.27 | 7.14 | 8.59 | 11.32 | 14.82 | 18.81 | 22.83 | 25.43 | 27.20 | 0.47 | 14.82 | 0.37 |
| 18.5 | 6.28 | 7.16 | 8.63 | 11.40 | 14.94 | 18.98 | 23.03 | 25.65 | 27.43 | 0.47 | 14.94 | 0.38 |
| 19 | 6.29 | 7.19 | 8.68 | 11.49 | 15.08 | 19.16 | 23.26 | 25.90 | 27.69 | 0.47 | 15.08 | 0.38 |




Fig. 3.a.—Graphical comparison of percentiles 3, 50 and 97 values of subscapular skinfold thickness between Hispanic American girls (INTER) and the reference of Centers for Disease Control and Prevention (CDC).




Fig. 3.b.—Graphical comparison of percentiles 3, 50 and 97 values of subscapular skinfold thickness between Hispanic American boys (INTER) and the reference of Centers for Disease Control and Prevention (CDC).

Table V
Triceps skinfold percentiles and L, M and S values of hispanic american boys

| Age (years) | P3 | P5 | P10 | P25 | P50 | P75 | P90 | P95 | P97 | L | M | S |
|-------------|------|------|------|------|-------|-------|-------|-------|-------|------|-------|------|
| 4 | 4.21 | 4.59 | 5.23 | 6.51 | 8.30 | 10.58 | 13.16 | 15.00 | 16.32 | 0.05 | 8.30 | 0.36 |
| 4.5 | 4.19 | 4.58 | 5.24 | 6.57 | 8.44 | 10.82 | 13.53 | 15.46 | 16.86 | 0.04 | 8.44 | 0.37 |
| 5 | 4.18 | 4.57 | 5.26 | 6.64 | 8.59 | 11.09 | 13.93 | 15.96 | 17.43 | 0.05 | 8.59 | 0.38 |
| 5.5 | 4.18 | 4.59 | 5.31 | 6.75 | 8.78 | 11.41 | 14.39 | 16.53 | 18.07 | 0.05 | 8.78 | 0.39 |
| 6 | 4.21 | 4.64 | 5.39 | 6.90 | 9.04 | 11.80 | 14.95 | 17.20 | 18.83 | 0.05 | 9.04 | 0.40 |
| 6.5 | 4.28 | 4.73 | 5.51 | 7.10 | 9.36 | 12.28 | 15.62 | 18.01 | 19.73 | 0.06 | 9.36 | 0.41 |
| 7 | 4.37 | 4.85 | 5.67 | 7.35 | 9.74 | 12.83 | 16.37 | 18.89 | 20.72 | 0.07 | 9.74 | 0.41 |
| 7.5 | 4.48 | 4.98 | 5.85 | 7.61 | 10.13 | 13.40 | 17.13 | 19.79 | 21.71 | 0.08 | 10.13 | 0.42 |
| 8 | 4.58 | 5.11 | 6.02 | 7.87 | 10.51 | 13.94 | 17.84 | 20.63 | 22.65 | 0.09 | 10.51 | 0.42 |
| 8.5 | 4.68 | 5.23 | 6.17 | 8.09 | 10.84 | 14.40 | 18.46 | 21.36 | 23.45 | 0.10 | 10.84 | 0.43 |
| 9 | 4.76 | 5.32 | 6.29 | 8.26 | 11.09 | 14.76 | 18.94 | 21.92 | 24.07 | 0.10 | 11.09 | 0.43 |
| 9.5 | 4.80 | 5.37 | 6.35 | 8.37 | 11.25 | 14.99 | 19.26 | 22.30 | 24.50 | 0.10 | 11.25 | 0.43 |
| 10 | 4.82 | 5.40 | 6.40 | 8.44 | 11.37 | 15.17 | 19.51 | 22.61 | 24.84 | 0.10 | 11.37 | 0.43 |
| 10.5 | 4.84 | 5.42 | 6.43 | 8.50 | 11.47 | 15.34 | 19.76 | 22.92 | 25.21 | 0.10 | 11.47 | 0.44 |
| 11 | 4.85 | 5.43 | 6.45 | 8.53 | 11.54 | 15.46 | 19.97 | 23.19 | 25.53 | 0.09 | 11.54 | 0.44 |
| 11.5 | 4.82 | 5.40 | 6.42 | 8.51 | 11.53 | 15.49 | 20.05 | 23.33 | 25.70 | 0.09 | 11.53 | 0.44 |
| 12 | 4.76 | 5.34 | 6.35 | 8.43 | 11.45 | 15.42 | 20.02 | 23.33 | 25.74 | 0.08 | 11.45 | 0.45 |
| 12.5 | 4.68 | 5.25 | 6.25 | 8.31 | 11.32 | 15.29 | 19.90 | 23.24 | 25.67 | 0.07 | 11.32 | 0.45 |
| 13 | 4.58 | 5.14 | 6.12 | 8.16 | 11.13 | 15.08 | 19.69 | 23.03 | 25.48 | 0.07 | 11.13 | 0.46 |
| 13.5 | 4.47 | 5.02 | 5.98 | 7.98 | 10.91 | 14.82 | 19.40 | 22.74 | 25.18 | 0.06 | 10.91 | 0.46 |
| 14 | 4.37 | 4.90 | 5.84 | 7.80 | 10.68 | 14.53 | 19.07 | 22.38 | 24.82 | 0.05 | 10.68 | 0.46 |
| 14.5 | 4.28 | 4.80 | 5.73 | 7.64 | 10.48 | 14.27 | 18.76 | 22.04 | 24.46 | 0.05 | 10.48 | 0.46 |
| 15 | 4.24 | 4.75 | 5.66 | 7.55 | 10.35 | 14.11 | 18.55 | 21.81 | 24.21 | 0.04 | 10.35 | 0.46 |
| 15.5 | 4.25 | 4.76 | 5.66 | 7.55 | 10.33 | 14.07 | 18.50 | 21.75 | 24.14 | 0.04 | 10.33 | 0.46 |
| 16 | 4.30 | 4.81 | 5.72 | 7.61 | 10.39 | 14.12 | 18.55 | 21.79 | 24.18 | 0.04 | 10.39 | 0.46 |
| 16.5 | 4.38 | 4.89 | 5.80 | 7.69 | 10.48 | 14.20 | 18.61 | 21.83 | 24.21 | 0.04 | 10.48 | 0.45 |
| 17 | 4.47 | 4.99 | 5.91 | 7.80 | 10.58 | 14.29 | 18.66 | 21.86 | 24.21 | 0.04 | 10.58 | 0.45 |
| 17.5 | 4.59 | 5.11 | 6.02 | 7.92 | 10.69 | 14.37 | 18.69 | 21.85 | 24.17 | 0.04 | 10.69 | 0.44 |
| 18 | 4.71 | 5.23 | 6.15 | 8.04 | 10.80 | 14.45 | 18.72 | 21.83 | 24.11 | 0.04 | 10.80 | 0.43 |
| 18.5 | 4.84 | 5.36 | 6.29 | 8.18 | 10.92 | 14.53 | 18.74 | 21.80 | 24.03 | 0.04 | 10.92 | 0.43 |
| 19 | 4.98 | 5.51 | 6.43 | 8.32 | 11.05 | 14.62 | 18.77 | 21.77 | 23.97 | 0.04 | 11.05 | 0.42 |

increase is also in top percentiles and being much more pronounced in boys (Figs. 4a, 4b). In general, in the female serie, the values of this study are below those corresponding to the CDC reference for all percentiles whilst in male serie, this values are lower only for 97th percentil.

Table VI represent the difference in mm between percentiles 3, 50 and 97 of the subscapular and triceps skinfolds with respect to the CDC reference. The difference were positive when the CDC reference value was greater than the value of Hispanic American sample

and they were negative when the opposite occurred. In general, the discrepancy between values of both skinfolds in Hispanic American and CDC reference is practically null in P3 values being around zero. For P50 the discrepancy are higher with values around 2 points and for the P97 there is the highest differences with values that ranged from zero in central ages until 9 or 14 in the extreme ages.

In relation to subscapular skinfold, there are generally negative values of the differences what it means that Hispanic American children and adolescent have





Fig. 4.a.—Graphical comparison of percentiles 3, 50 and 97 values of triceps skinfold thickness between Hispanic American girls (INTER) and the reference of Centers for Disease Control and Prevention (CDC).

Fig. 4.b.—Graphical comparison of percentiles 3, 50 and 97 values of triceps skinfold thickness between Hispanic American boys (INTER) and the reference of Centers for Disease Control and Prevention (CDC).

Table VI
Differences in mm between Hispanic American mean values and CDC reference values of skinfolds percentiles 3, 50 and 97 in both sexes

| Age (years) | Subscapular skinfold | | | | | | Triceps skinfold | | | | | |
|-------------|----------------------|-------|-------|-------|-------|-------|------------------|-------|-------|-------|-------|-------|
| | Boys | | | Girls | | | Boys | | | Girls | | |
| | 3 | 50 | 97 | 3 | 50 | 97 | 3 | 50 | 97 | 3 | 50 | 97 |
| 4 | -0.13 | -0.23 | -2.18 | -0.30 | -1.01 | -6.27 | 1.24 | 0.69 | -0.78 | 0.54 | -0.76 | -2.47 |
| 4.5 | -0.21 | -0.49 | -3.46 | -0.29 | -1.07 | -6.59 | 1.09 | 0.38 | -1.30 | 0.62 | -0.63 | -2.22 |
| 5 | -0.26 | -0.77 | -4.81 | -0.29 | -1.14 | -6.87 | 0.91 | 0.04 | -1.83 | 0.69 | -0.51 | -1.97 |
| 5.5 | -0.35 | -1.05 | -6.27 | -0.31 | -1.23 | -7.12 | 0.73 | -0.32 | -2.40 | 0.75 | -0.40 | -1.71 |
| 6 | -0.45 | -1.36 | -7.81 | -0.37 | -1.37 | -3.36 | 0.52 | -0.74 | -3.01 | 0.79 | -0.32 | -1.44 |
| 6.5 | -0.56 | -1.67 | -9.36 | -0.44 | -1.54 | -7.55 | 0.31 | -1.16 | -3.62 | 0.80 | -0.26 | -1.18 |
| 7 | -0.64 | -1.94 | -10.8 | -0.49 | -1.71 | -7.68 | 0.12 | -1.57 | -4.13 | 0.80 | -0.25 | -0.95 |
| 7.5 | -0.71 | -2.18 | -12.1 | -0.54 | -1.91 | -7.69 | -0.04 | -1.89 | -4.42 | 0.75 | -0.29 | -0.78 |
| 8 | -0.77 | -2.40 | -13.1 | -0.60 | -2.06 | -7.36 | -0.15 | -2.12 | -4.45 | 0.69 | -0.33 | -0.56 |
| 8.5 | -0.81 | -2.57 | -13.7 | -0.63 | -2.13 | -6.52 | -0.23 | -2.25 | -4.15 | 0.64 | -0.32 | -0.15 |
| 9 | -0.81 | -2.68 | -13.7 | -0.64 | -2.14 | -5.26 | -0.27 | -2.25 | -3.53 | 0.60 | -0.28 | 0.39 |
| 9.5 | -0.81 | -2.72 | -13.1 | -0.76 | -2.15 | -3.78 | -0.25 | -2.15 | -2.63 | 0.57 | -0.23 | 1.00 |
| 10 | -0.79 | -2.74 | -12.0 | -0.80 | -2.19 | -2.24 | -0.22 | -2.02 | -1.63 | 0.52 | -0.18 | 1.62 |
| 10.5 | -0.79 | -2.77 | -10.6 | -0.86 | -2.24 | -0.74 | -0.21 | -1.92 | -0.75 | 0.44 | -0.15 | 2.23 |
| 11 | -0.80 | -2.82 | -9.09 | -0.93 | -2.30 | 0.64 | -0.22 | -1.86 | 0.01 | 0.36 | -0.14 | 2.78 |
| 11.5 | -0.82 | -2.87 | -7.45 | -1.01 | -2.37 | 1.89 | -0.22 | -1.80 | 0.68 | 0.27 | -0.14 | 3.25 |
| 12 | -0.85 | -2.89 | -5.85 | -1.07 | -2.39 | 3.03 | -0.23 | -1.75 | 1.19 | 0.20 | -0.15 | 3.67 |
| 12.5 | -0.87 | -2.87 | -4.35 | -1.11 | -2.36 | 4.11 | -0.24 | -1.74 | 1.52 | 0.16 | -0.11 | 4.09 |
| 13 | -0.86 | -2.82 | -3.01 | -1.11 | -2.26 | 5.14 | -0.25 | -1.74 | 1.70 | 0.17 | -0.02 | 4.55 |
| 13.5 | -0.86 | -2.72 | -1.85 | -1.06 | -2.08 | 6.11 | -0.27 | -1.75 | 1.78 | 0.23 | 0.15 | 5.06 |
| 14 | -0.83 | -2.62 | -0.87 | -0.96 | -1.84 | 6.97 | -0.31 | -1.77 | 1.78 | 0.35 | 0.39 | 5.62 |
| 14.5 | -0.82 | -2.52 | -0.11 | -0.83 | -1.57 | 5.91 | -0.34 | -1.81 | 1.73 | 0.51 | 0.69 | 6.20 |
| 15 | -0.82 | -2.47 | 0.42 | -0.68 | -1.29 | 10.3 | -0.40 | -1.89 | 1.59 | 0.69 | 1.01 | 6.75 |
| 15.5 | -0.84 | -2.47 | 0.70 | -0.54 | -1.04 | 8.98 | -0.50 | -2.03 | 1.34 | 0.89 | 1.32 | 7.21 |
| 16 | -0.87 | -2.49 | 0.90 | -0.41 | -0.82 | 9.63 | -0.60 | -2.21 | -1.41 | 1.09 | 1.62 | 7.63 |
| 16.5 | -0.88 | -2.50 | 1.14 | -0.30 | -0.61 | 10.3 | -0.72 | -2.35 | 0.99 | 1.30 | 1.94 | 8.03 |
| 17 | -0.87 | -2.46 | 1.47 | -0.20 | -0.41 | 11.1 | -0.82 | -2.46 | 1.06 | 1.53 | 2.26 | 8.37 |
| 17.5 | -0.80 | -2.35 | 1.96 | -0.09 | -0.19 | 11.9 | -0.92 | -2.50 | 1.35 | 1.75 | 2.56 | 8.67 |
| 18 | -0.70 | -2.15 | 2.64 | 0.01 | 0.01 | 12.7 | -0.99 | -2.48 | 1.85 | 1.97 | 2.83 | 8.89 |
| 18.5 | -0.56 | -1.90 | 3.44 | 0.09 | 0.20 | 13.4 | -1.04 | -2.42 | 2.53 | 2.16 | 3.07 | 9.05 |
| 19 | -0.41 | -1.60 | 4.28 | 0.16 | 0.37 | 14.1 | -1.10 | -2.34 | 3.26 | 2.34 | 3.29 | 9.17 |
| Mean | -0.70 | -2.20 | -4.81 | -0.56 | -1.45 | 1.85 | -0.18 | -1.68 | -2.20 | 0.81 | 0.50 | 3.25 |

higher subscapular adiposity. However, p97 of girls up to 11 years are lower than those in CDC and this discrepancy clearly increase with age rising 14 positive points.

The difference for triceps skinfold between Hispanic American children and adolescent and the CDC reference are less evident than for subscapular skinfold rising smallest values especially in boys for the three percentiles. The greatest difference for triceps skinfold is observed for the P97 in girls up to 11 years.

Discussion

This study was the first to consider a child and adolescent group of Hispanic American origin to establish skinfold percentiles. The relevance of this analysis was based on the finding of an increase in skinfolds at a rate of 0.4 to 0.5 mm per decade over the 1951-2003 period analyzed by Olds²³ using information provided by 154 studies on more than 458,547 children and adolescents in 30 developed countries. This increase in skinfold thickness was consistent with increased BMI in the world and the obesity epidemic^{2,24,25}. However, the prevalence of excess weight (overweight and obesity) observed in the sample of this study was lower than the reported values for the different countries also participating in the Olds²³ study: Argentina²⁶, Cuba²⁷, Mexico²⁸ and Spain²⁹.

The differences found in skinfold percentiles over the years in the Olds study²³ might primarily arise in terms of the methodology used to calculate them. Currently there are a number of statistical methods based on the adjustment of mathematical models controlling kurtosis and/or asymmetry of raw anthropometric data and adequately represent their changes and trends in terms of age. One such tool is the LMS method, which also describes the temporal changes of anthropometric measurements, and provides a set of statistical parameters that enable further biological interpretation of growth and interpopulation comparison^{19,30,31}. The results obtained in the present research are comparable with those of the CDC reference because they were obtained using the same LMS method but there is a methodological difference between both studies that is related to the adipometer used to measure the skinfolds. In all national health surveys conducted in the United States, skinfolds were measured to the nearest 0.5 mm by using Lange calipers, except for the III National Health and Nutrition Examination Survey (NHANES) in which Holtain calipers were used with a sensitivity of 0.2 mm¹². In the present work, however, all the children were measured with a Holtain caliper by highly qualified staff.

The greatest difference between this study and the CDC reference relate to the ethnicity of the populations analyzed but also the age and characteristics of the sample. The ethnic composition of Latin American populations is the convergence in varying degrees, ac-

ording to their geographical position in the vast territory of the Americas, of three parental populations: Amerindian, European and African. The percentage of genetic mixing of these populations, estimated with different molecular markers, varies depending on this geographical distribution and differential parental contribution and is a valuable indicator of population and migration dynamics³². In the CDC reference children were identified from the ethnical-racial point of view as black or white, but the values of percentiles in tables were presented together, as opposed to graphical displays where black boys and girls showed P50 values of triceps skinfold lower than those of white children¹². In the subscapular skinfold these differences were less evident in the graphical display of the publication¹². As for the age of the samples, anthropometric data used in the present study were collected in 2005-2008 whilst the CDC reference mix data from the National Health Examination Survey II and III conducted between 1963 and 1970 and with the NHANES I, II and III which took place between 1971 and 1994. In addition, the children and adolescents included in the present study were schoolchildren without pathologies, attending public schools and of middle socioeconomic conditions, whilst the CDC reference data corresponded to non-prescriptively selected children and adolescents representing the entire civilian, noninstitutionalized population of the United States.

Studies on body composition and particularly skinfold measurement are very scarce in Hispanic American populations and most of the existing ones have been performed in resident or migrant children in the United States. Comparisons between the Hispanic American sample and the CDC reference conducted in this study were consistent with the differences in the thickness of the subscapular, triceps, suprailiac and medial calf skinfolds found between white American children and American children of Mexican origin included in NHANES (1982-1984) who tend to have thicker skinfolds³³; however the measurements were not comparable because of the different calipers used, the age groups made using different criteria and because were not calculated using the LMS method.

The comparison between the hispanic american skinfolds of this study with respect to the CDC reference also indicated that there were differences in the distribution of adipose tissue. Interpopulational differences were observed in the fat distribution in terms of the trunk (main) and extremities (peripheral) model³⁴ and that in adolescents, lower trunk extremity and upper trunk components accounted for 80% of the variance in fat distribution³⁵. This was also evaluated in a oriental sample recruited using a non-random purposive sampling approach consisting of 578 children aged 8-10 years from China, Lebanon, Malaysia and Thailand, after controlling for height and weight there was a significant ethnic difference in biceps, triceps, subscapular, supraspinal, and medial calf skinfolds³⁶. The Chinese and Thai children showed increased fatty

deposition in the trunk compared to the Malays, who in turn had higher values than the Lebanese. Specifically, the ratio of the subscapular to the triceps skinfolds (S/T ratio) in children of Asian Americans was higher with respect to the Mexican, European, and African ones³⁴ and white children from Arizona State, compared to Asian and American children of Hispanic American origin, presented a smaller android/gynoid fat ratio³⁵. The results found in the present study partially confirmed these findings because the boys and girls evaluated showed greater central adiposity, based on the subscapular skinfold, than the children of the CDC reference, especially in 3 and 50 percentiles.

According to Freedman, at similar age stage and BMI-for-age, the body fatness of children and adolescents can differ by up to 5% across racial/ethnic groups⁴. In fact, the interpopulation differences in the content and distribution of body fat, can be shown even within this Hispanic American sample that have a closer ethnic origin compared with the United States sample. In a preliminary study based on 2436 infants and 6-9 year-old children from Spain, Mexico, Cuba, Venezuela and Argentina³⁷ it was revealed that: a) the higher total and relative adiposity corresponded to Mexican and Argentine series and the lower one to Venezuela, leaving Spain and Cuba in an intermediate position; b) with regard to the pattern of distribution of adiposity, the Spanish series featured a more peripheral trend, whereas the Argentine one showed a well pronounced backbone arrangement.

Given this background, Hispanic American children and adolescents evaluated in the present study showed a pattern of distribution of triceps and subscapular adiposity different from the CDC reference. While it has been proposed that theoretically human populations should follow a similar growth pattern in similar environmental conditions, it cannot be excluded that some of the interpopulation differences in this pattern may reflect not only the influence of the environment but also differences in the genetic potential³⁸. The differences found in this research, in addition to ethnic and health characteristics of the samples, could be also attributed to the effect of secular changes in body composition because the most recent part of CDC sample is dated in 1994 whilst the measurements of this study were taken in 2005-2008.

Because BMI is of easy determination and because it has a high correlation with body fat, it is epidemiologically the most frequently used indicator to evaluate excess body fat in children and adolescents²⁵. The most common criteria used to define overweight and obesity based on BMI among children are the proposed by three expert organizations: IOTF^{16,17}, CDC³⁹ and World Health Organization (WHO)⁴⁰. The comparison of the prevalences of nutritional status obtained with these criteria in populations of different ethnic origin⁴¹⁻⁴⁴, including Latin American's⁴⁵⁻⁴⁹, provide different results that can lead to erroneous conclusions. These studies also indicated the need to identify BMI cutoffs that are

properly associated with an increased risk of health problems later in life⁴¹. The sensitivity and specificity of the criteria and cutoff points of BMI as predictors of body fat should be based on comparative analyses of body composition from subcutaneous skinfolds or other direct methods on estimating body fat, but such studies on school children have been scarce to date^{44,49}. The percentiles of triceps and subscapular folds calculated in the present study can contribute to the purposes of such comparative studies.

Conclusions

The percentiles of triceps and subscapular skinfolds of contemporary and healthy Hispanic American children and adolescents from different localities in Spain, Argentina, Cuba, Mexico and Venezuela, differ substantially with respect to the American CDC reference that are commonly used for adiposity determination in these countries. Because adiposity depends on the reference used, the skinfold percentiles calculated in this study could be applied to people of a similar ethnicity, especially in comparative studies of body composition.

Acknowledgments

The authors acknowledge Ms. Estela Chaves, Institute of High Altitude Biology, for her collaboration in the design of tables and figures. This study was supported by funds from Ministry of Science and Technology of Spanish Government (Project GCL2005-03157), Argentinean National Agency for Scientific and Technological Advancement (ANPCyT) (PICTO 2008-00139) and National University of Jujuy in Argentina.

Conflicts of interest

None.

References

1. Uauy R, Albala C, Kain J. Obesity trends in Latin America: transitioning from under- to overweight. *J Nutr*; 2001; 131: 893S-9S.
2. Lobstein T. Prevalence and trends in childhood obesity. In: Crawford D, Robert WJ, Ball K, Brug J (Eds). *Obesity Epidemiology: From Aetiology to Public Health*. Oxford University Press. 2010.
3. Peña M, Bacallao J. La obesidad en la pobreza: un nuevo reto para la salud pública. *Publicación Científica*, 576. Organización Panamericana de la Salud (OPS). 2000.
4. Freedman DS, Wang J, Thornton JC, Mei Z, Pierson RN, Dietz WH, Horlick M. Racial/ethnic Differences in Body Fatness Among Children and Adolescents. *Obesity*; 2008; 16: 1105-11.
5. He Q, Horlick M, Thornton J, Wang J, Pierson RN, Heshka S, Gallagher D. Sex-Specific Fat Distribution Is Not Linear Across Pubertal Groups in a Multiethnic Study. *Obes Res*; 2004; 12: 725-733.

6. Novotny R, Going S, Teegarden D, Van Loan M, McCabe G, McCabe L, *et al.* Hispanic and Asian pubertal girls have higher android/gynoid fat ratio than whites. *Obesity*, 2007; 15: 1565-70.
7. Meyer KA, Friend S, Hannan PJ, Himes HE, Demerath EW, Neumark-Sztainer D. Ethnic variation in body composition assessment in a sample of adolescent girls. *Int J Ped Obes*, 2011; 6: 481-90.
8. Staiano AE, Katzmarzyk PT. Ethnic and sex differences in body fat and visceral and subcutaneous adiposity in children and adolescents. *Int J Obes*, 2012; 36: 1261-9.
9. Freedman DS, Sherry B. The validity of BMI as an indicator of body fatness in childhood. *Pediatrics*, 2009; 124: S23-S34.
10. Marrodán MD, González-Montero M, Morales E. Relationship between physical measures of anthropometry and bioimpedance measures. In: Preedy VR (Edt). *Handbook of Anthropometry: Measures of Human Form in Health and Disease*. Springer. 2012. pp. 459-77.
11. Freedman DS, Serdula MK, Srinivasan SR, Berenson GS. Relation of circumferences and skinfold thicknesses to lipid and insulin concentrations in children and adolescents: the Bogalusa Heart Study. *Am J Clin Nutr*, 1999; 69(2): 308-17.
12. Addo OY, Himes JH. Reference curves for triceps and subscapular skinfold thicknesses for US children and adolescents. *Am J Clin Nutr*, 2010; 91: 635-42.
13. Tanner JM. Standards from birth to maturity for height, weight, height velocity and weight velocity. *Arch Dis Child*, 1966; 41: 613-35.
14. World Medical Association (WMA). *Ethical principles for medical research involving human subjects*. Helsinki: Helsinki Declaration. 2000.
15. Weiner JS, Lourie JA. *Practical Human Biology*. Academic Press. London. 1981.
16. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*, 2000; 320: 1-6.
17. Cole TJ, Flegal KM, Nicholls D, Jackson A. Body mass index cut offs to define thinness in children and adolescents: international survey. *BMJ*, 2007; 335: 194.
18. Cabañas D, Esparza F. *Compendio de Cineantropometría*. CTO Editorial. 2009.
19. Cole TJ, Green PJ. Smoothing reference centile curves: the LMS method and penalized likelihood. *Statist Med*, 1992; 11: 1305-19.
20. Cole TJ. Fitting smoothed centile curves to reference data. *J R Statist Soc*, 1988; 151: 385-418.
21. Cole TJ. The British, American NCHS, and Dutch weight standard compared using the LMS method. *Am J Hum Biol*, 1989; 1: 397-408.
22. Pan H, Cole TJ. LMSchartmaker, a program to construct growth references using the LMS method. 2011. Version 2.43. <http://www.healthforallchildren.co.uk>.
23. Olds TS. One million skinfolds: secular trends in the fatness of Young people 1951-2004. *Eur J Clin Nutr*, 2009; 63: 934-46.
24. Rudolf MC, Levine R, Feltbower R, Connor A, Robinson M. The TRENDS project: development of a methodology to reliably monitor the obesity epidemic in childhood. *Arch Dis Child*, 2006; 91: 309-11.
25. Albala C, Corvalan C. Epidemiology of obesity in children in South America. In: Moreno LA, Pigeot I, Ahrens W (Eds). *Epidemiology of Obesity in Children and Adolescents*. Springer Series on Epidemiology and Public Health, 2011; 2(1): 95-110.
26. Oyhenart EE, Dahinten L, Alba J, Alfaro EL, Bejarano IF, Cabrera GE, *et al.* Estado nutricional infante juvenil en seis provincias de Argentina. Variación regional. *Rev Arg Antropol Biol*, 2008; 10(1): 1-62.
27. Fariñas Rodríguez L, Vázquez Sánchez V, Martínez Fuentes A, Fuentes Smith EL, Toledo Borrero E, Martiati Hendrich M. Evaluación nutricional de niños de 6 a 11 años de Ciudad de La Habana. *Rev Cubana Invest Biomed*, 2011; 30(4): 439-49.
28. Barquera Cervera S, Campos-Nonato I, Rojas R, Rivera J. Obesidad en México: epidemiología y políticas de salud para su control y prevención. *Gac Méd Méx*, 2010; 146: 397-407.
29. Serra-Majem LL, Ribas L, Aranceta J. Epidemiología de la obesidad en España. Resultados del estudio Enkid (1998-2000). In: Serra-Majem LL, Aranceta J (Eds.). *Obesidad Infantil y Juvenil. Estudio Enkid*. Masson, Barcelona. 2001. pp.81-108.
30. Alfaro E; Bejarano IF, Dipierri JE, Quispe Y, Cabrera G. Centílos de peso, talla e índice de masa corporal de escolares jujeños calculados por el método LMS. *Arch Arg Ped*, 2004; 102(6): 434-40.
31. Alfaro EL, Vázquez ME, Bejarano IF, Dipierri JE. The LMS method and weight and height centiles in Jujuy, (Argentina) children. *HOMO*, 2008; 59: 223-34.
32. Wang S, Ray N, Rojas W, Parra MV, Bedoya G, Gallo C, Poletti G, Mazzotti G, Hill K, Hurtado AM, Camrena B, Nicolini H, Klitz W, Barrantes R, Molina JA, Freire NB, Bortolini MC, Salzano FM, Petzl-Erler ML, Tsuneto LT, Dipierri JE, Alfaro EL, Bailliet G, Bianchi NO, Llop E, Rothhammer F, Excoffier L, Ruiz-Linares A. Geographic patterns of genome admixture in latinamerican mestizos. *PLoS Genet*, 2008; 4(3): e1000037.
33. Ryan AS, Martínez GA, Baumgartner RN, Roche AF, Guo S, Chumlea WC, Kuczumarski RJ. Median skinfold thickness distributions and fat-wave patterns in Mexican-American children from the Hispanic Health and Nutrition Examination Survey (HHANES 1982-1984). *Am J Clin Nutr*, 1990; 51(5): 925S-35S.
34. Malina RM, Huang YC, Brown KH. Subcutaneous adipose tissue distribution in adolescent girls of 4 ethnic groups. *Int J Obes Relat Metab Disord*, 1995; 19: 793-7.
35. Hattori K, Becque MD, Katch VL, Katch VL, Rocchini AP, Boileau RA, Slaughter MH, Lohman TG. Fat patterning of adolescents. *Ann Hum Biol*, 1987; 14 (1): 23- 8.
36. Liu A, Byrne NM, Kagawa M, Ma G, Kijboonchoo K, Nasreddine L, Koon Poh B, Ismail MN, Hills AP. Ethnic differences in body fat distribution among Asian pre-pubertal children: a cross-sectional multicenter study. *BMC Public Health*, 2011; 26(11): 500.
37. Romero-Collazos JF, Marrodán Serrano MD, Mesa Santurino MS, Bejarano IF, Alfaro EL, Martínez AJ, *et al.* Grasa corporal y distribución de la adiposidad en escolares latinoamericanos y españoles. In: Gutiérrez-Redomero E, Sanchez-Andrés A, Galera-Olmos V. *Diversidad Humana y Antropología Biológica*. Universidad de Alcalá de Henares. 2010. pp.221-8.
38. Butte NF, Garza C, de Onis M. Evaluation of the Feasibility of International Growth Standards for School-Aged Children and Adolescents. *J Nutr*, 2007; 137(1): 153-7.
39. Kuczumarski RJ, Ogden CL, Guo SS, Grummer-Strawn LM, Flegal KM, Mei Z, Wei R, Curtin LR, Roche AF, Johnson CL. 2000 CDC Growth Charts for the United States: Methods and development. National Center for Health Statistics. *Vital Health Stat*, 2002; 11(246): 1-190.
40. De Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmanna J. Development of a WHO growth reference for school-aged children and adolescents. *WHO Bulletin n°85*. 2007; (9): 660-7.
41. Monasta L, Lobstein T, Cole TJ, Vignerová J, Cattaneo A. Defining overweight and obesity in pre-school children: IOTF reference or WHO standard? *Obes Rev*, 2010; 12: 295-300.
42. So HK, Nelson EAS, Sung RYT, Ng PC. Implications of using World Health Organization growth reference (2007) for identifying growth problems in Hong Kong children aged 6 to 18 years. *Hong Kong Med J*, 2011; 17(3): 174-9.
43. Kulaga Z, Litwin M, Tkaczyk M, Różdzyńska A, Barwicka K, Grajda A, Świąder B, Gurzkowska E, Napieralska HP. The height-, weight-, and BMI-for-age of Polish school-aged children and adolescents relative to international and local growth references. *BMC Public Health*, 2010; 10: 109.
44. De Sousa Lopes HM. Diagnostic accuracy of CDC, IOTF and WHO criteria for obesity classification, in a Portuguese school-aged children population. Maestrado en Saúde Pública. Universidad do Porto. Faculdade de Medicina. Instituto de Ciências Biomédicas Abel Salazar. 2012.
45. Baya Botti A, Pérez-Cueto FJA, Vázquez Monllor PA, Kols-teren PW. International BMI-for-age references underestimate

- thinness and overestimate overweight and obesity in Bolivian adolescents. *Nutr Hosp*, 2010; 25(3): 428-36.
46. Padula G, Salceda S. Comparación entre referencias de las prevalencias de sobrepeso y obesidad, estimadas a través del Índice de Masa Corporal, en niños de Argentina. *ALAN*, 2008; 58(4): 330-5.
 47. Kovalskys I, Rausch Herscovici C, De Gregorio MJ.. Nutritional status of school-aged children of Buenos Aires, Argentina: data using three references. *J Public Health*, 2010; 33(3): 403-11.
 48. Meyer E, Carrillo R, Román EM, Bejarano IF, Dipierri JE. Prevalence of overweight and obesity in students from different altitudinal zones of Jujuy according to three international references (IOTF, CDC and WHO). *Arch Argent Pediatr*, 2013; 111(6): 516-22.
 49. Deurenberg P, Yap M, van Staveren WA. Body mass index and percent body fat: a meta analysis among different ethnic groups. *Int J Obesity Relat Metab Disor*, 1998; 22(12): 1164-8.