



Revisión

Relationship between food insecurity and malnutrition in schoolchildren from low- and middle-income countries — A systematic review

Relación entre inseguridad alimentaria y desnutrición en escolares de países de bajos y medianos ingresos: revisión sistemática

Diana Marcela Valenzuela-Guzmán, Julián Esparza-Romero

Diabetes Research Unit. Centro de Investigación en Alimentación y Desarrollo A.C. Hermosillo, Sonora. Mexico

Abstract

The objective of this review is to study the relationship between food insecurity (FI) and malnutrition in schoolchildren from low- and middle-income countries (LMIC). The review was conducted using the databases PubMed, MEDLINE, CENTRAL, LILACS and SCIELO during the months of March to April 2022 without language or publication date restrictions. The search strategy consisted of combinations of text words and controlled vocabulary (MeSH terms and DeCS) related to "schoolchildren", "low- and middle-income countries" and "food insecurity". Fifteen studies were included in this review. Studies assessing FI and undernutrition in LMIC schoolchildren have indicated that FI is associated with lower height-for-age and higher prevalence of undernutrition overall. Only two studies identified a positive risk association between FI and overweight and obesity, the remaining studies suggested that schoolchildren with FI have a lower risk of overweight and obesity than those without FI. The review suggests a link between FI and undernutrition in schoolchildren from LMIC, with controversial results on overweight and obesity. Comprehensive public health policies should consider contextual and population-specific factors in addressing FI's impact on nutritional status.

Keywords:

Food insecurity. Children. Malnutrition. Vulnerable populations. Low- and middle-income countries.

Resumen

El objetivo de esta revisión fue estudiar la relación entre la inseguridad alimentaria (IA) y la desnutrición en escolares de países de bajos y medianos ingresos (PBMi). La revisión se realizó utilizando las bases de datos PubMed, MEDLINE, CENTRAL, LILACS y ScieLo durante los meses de marzo a abril de 2022 sin restricciones de idioma o fecha de publicación. La estrategia de búsqueda consistió en combinaciones de palabras y vocabulario controlado (términos MeSH y DeCS) relacionados con "escolares", "países de bajos y medianos ingresos" e "inseguridad alimentaria". Quince estudios se incluyeron en esta revisión. Los estudios que evaluaron la IA y la desnutrición en escolares de PBMi mostraron que la IA está asociada con una menor talla para la edad y una mayor prevalencia de desnutrición en general. Solo dos estudios encontraron una asociación de riesgo positiva entre la IA y el sobrepeso y la obesidad; el resto de los estudios sugieren que los escolares con IA tienen menor riesgo de sobrepeso y obesidad que aquellos sin IA. La revisión sugiere una asociación entre la IA y la desnutrición en escolares de PBMi, con resultados contradictorios en el sobrepeso y la obesidad. Las políticas de salud pública deberían considerar factores contextuales y específicos de la población al abordar el impacto de la IA en el estado nutricional.

Palabras clave:

Inseguridad alimentaria. Escolares. Desnutrición. Poblaciones vulnerables. Países de bajos y medianos ingresos.

Received: 05/07/2023 • Accepted: 13/02/2024

Conflicts of interest: the authors declare to be free of conflicts of interest.

Artificial intelligence: the authors declare not to have used artificial intelligence (AI) or any AI-assisted technologies in the elaboration of the article.

Valenzuela-Guzmán DM, Esparza-Romero J. Relationship between food insecurity and malnutrition in schoolchildren from low- and middle-income countries — A systematic review. *Nutr Hosp* 2024;41(4):897-905
DOI: <http://dx.doi.org/10.20960/nh.04847>

Correspondence:

Julián Esparza-Romero. Diabetes Research Unit. Centro de Investigación en Alimentación y Desarrollo, A.C. Gustavo Enrique Astiazarán Rosas Road, 46. 83304 Hermosillo, Sonora. Mexico
e-mail: julian@ciad.mx

INTRODUCTION

Malnutrition affects physical, mental and social development, especially in children (1). The World Health Organization (WHO) defines malnutrition as the product of “deficiencies, excesses and imbalances in a person’s caloric and/or nutrient intake”, this definition includes both undernutrition and overweight and obesity (1). Multiple studies have documented that children with undernutrition have a higher risk of dying from any cause, an increased risk of infectious and respiratory diseases, diarrhea, anemia, as well as poor physical and cognitive development (1,2). On the other hand, overweight and obesity in children are associated with lower self-esteem and an increased risk of skin and orthopedic diseases, as well as a higher risk of developing lipid abnormalities, sleep apnea, cardiovascular disease, type 2 diabetes, and some types of cancer in adulthood (1,2).

The children with the highest risk of malnutrition are those who are in vulnerable environments, that is, they live in rural communities or in low-income households or belong to an indigenous community or ethnic minority (3). Globally, 18.4 % of children between the ages of 5 and 19 were overweight and obese in 2016 (4). Although developed countries have the highest prevalence, low- and middle-income countries (LMICs) have seen a significant increase in their rates of obesity and overweight in the last decade (4). Undernutrition in schoolchildren is rarely reported worldwide, but several local studies have shown that rural and indigenous communities in LMICs are the most affected (2). Malnutrition can occur due to limited access to nutritious food as in food insecurity (FI).

The Food and Agriculture Organization (FAO) defines FI as “the limited or uncertain availability of nutritionally adequate and safe foods, or limited or uncertain ability to acquire acceptable foods in socially acceptable ways” (5). FI is associated with multiple health problems, including a higher risk of asthma, anemia, cognitive problems, anxiety and depression, than people who are food secure (6-8). In the world, 29.3 % of people were moderately or severely food insecure in 2021, with LMICs, such as Africa (57.9 %) and Latin America and the Caribbean (40.6 %), having the highest prevalence (2).

Multiple studies have evaluated the relationship between FI and malnutrition. In the case of adults and adolescents, a higher level of FI is associated with a higher prevalence of overweight and obesity, particularly in women (9,10). In contrast, in pre-school children (< 5 years), higher levels of FI are associated with an increased risk of developing undernutrition (10,11). In relation to schoolchildren (5 to 12 years), higher levels of FI have been associated with lower diet quality (12). Schoolchildren with FI have a lower consumption of fruits, vegetables, fiber, legumes, as well as a higher intake of calories, mainly from saturated fats and added sugars than their peers without FI (13-16). This indicates that schoolchildren with FI have a greater risk of malnutrition than those without FI.

A century ago, being overweight and obese was associated with wealth, but not anymore. In developed countries, poor children are often the most likely to be overweight or obese (17).

Children who are overweight often come from socioeconomically disadvantaged families. In the United States, for example, the rate of overweight in children decreases as educational level and family income increase (18). Sopoede et al. conducted a systematic review whose objective was to evaluate the relationship between FI and malnutrition in children aged 2 to 19 years in the US (19). The authors concluded that there is no evidence showing a relationship between FI and undernutrition. In relation to FI and overweight and obesity, the results were not consistent, but there seems to be a trend towards a higher risk of obesity. For schoolchildren (5-12 years) in LMICs, little is known about the relationship between FI and malnutrition. The purpose of this review is to study the relationship between FI and nutritional status in schoolchildren with characteristics of vulnerability from LMICs.

METHODS

CRITERIA FOR CONSIDERING STUDIES FOR THIS REVIEW

Types of studies

All study designs were considered for inclusion in this review.

Participants

The population was schoolchildren (5 to 12 years) in conditions of vulnerability from LMICs. Vulnerability was considered when they lived in a rural community, had a low socioeconomic level and/or belonged to an Indigenous community or ethnic minority. Studies in which the majority of participants (more than 80 %) are in this age range or present separate data for this age range were included.

Exposure

The review included studies that assessed FI using a validated survey tool.

Comparisons

Schoolchildren from the same environment without FI were considered as a comparison.

Outcome measures

Primary outcomes were BMI-for-age Z-score (BMI-Z), odds ratio (OR)/relative risk (RR) for undernutrition, and OR/RR for overweight and obesity. Other anthropometric measures such as weight, waist circumference, skinfolds, height-for-age Z-score (HAZ), weight-for-

age Z-score (WAZ), weight-for-height Z-score (WHZ) as well as the prevalence/incidence of malnutrition and prevalence/incidence of overweight and obesity were considered secondary outcomes.

SEARCH STRATEGY

The search for articles was performed using the databases PubMed (interface National Library of Medicine), MEDLINE (OVID interface), The Cochrane Library – CENTRAL (OVID interface), LILACS (Virtual Health Library) and SciELO (Scientific Electronic Library Online). The search was conducted during the months of March to April 2022 without language or publication date restrictions. The search strategy consisted of combinations of text words and controlled vocabulary (MeSH terms and DeCS) related to “schoolchildren”, “low- and middle-income countries” and “food insecurity” considering the search platform. The PubMed search strategy can be found in the supplementary material (Supplementary Table I). Following the recommendations of the Cochrane handbook, the bibliographical references of systematic reviews and included studies were considered as additional search sources (20).

SELECTION OF STUDIES

For the article selection process, the Mendeley version 1.19.8 software was used, to which the results of the searches in the different databases were imported for subsequent processing. In the first stage, duplicate results were removed. In the second stage, titles and abstracts were reviewed eliminating articles that clearly did not meet the inclusion criteria. If the inclusion criteria were not well defined in the abstract, the article went to the third

stage. In the third stage, the full text was reviewed to assess whether it met the inclusion criteria. The study selection process is described in the PRISMA flowchart (Fig. 1).

DATA EXTRACTION AND SYNTHESIS

For data extraction, an electronic format (Excel sheet) was used, which collected the title of the article, author, year of publication, type of study, study population, number of participants, measurement tools and cut-off points for FI and malnutrition, in addition to the main results. The information collected from the included studies is presented through a narrative synthesis.

RESULTS

CHARACTERISTICS OF INCLUDED STUDIES

Of the 36 full-text articles that were evaluated, nine did not evaluate the relationship between FI and malnutrition, five did not meet the age criteria, and seven were not conducted in LMICs (Fig. 1). The 15 studies included in the review had a cross-sectional design and were published between 2007 and 2021 (6,21,30-34,22-29). These studies analyzed populations from Brazil (21,31), Ethiopia (23,30), the Philippines (24), Malaysia (25), Venezuela (27), Nicaragua (6), Colombia (34), Bolivia (22), Mexico (26,28,29,33) and Jamaica (32). The sample size ranged from 61 to 7181 schoolchildren, and all studies had vulnerability characteristics.

Nine different tools were used to assess FI, eight of which assessed FI at the household level and one at the personal level. The tools used were Brazilian Food Insecurity Scale (EBIA) (21,31),




Figure 1.

Flowchart of Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA).

Household Food Insecurity Access Scale (HFIAS) (22,23), Food Insecurity Experience Scale (FIES) (24), Radimer/Cornell Hunger and Food Insecurity Instrument (25), USDA Household Food Security Survey Module (HFSSM) (34) and its version short (27), Latin American and Caribbean Food Security Questionnaire (ELCSA) (6,29), Food and Nutrition Technical Assistance (FANTA) (30), Mexican Food Safety Scale (EMSA) (26,28) and 2-items Hunger Vital Sign (HVS) (32,33).

To assess nutritional status, BMI-Z, HAZ, WAZ and WHZ were used. To classify children, 12 studies used the WHO 2007 cut-off points (21,22,31,34,23-30), one the CDC 2000 cut-off points (33), one the Frisancho 2008 cut-off points (6), and one the Cole references specific for sex and age (32).

FOOD INSECURITY AND UNDERNUTRITION

Eight articles evaluated the relationship between FI and undernutrition, the summary of results is shown in table I. Three studies found that higher levels of FI were associated with an increased risk (OR, 1.34 to 2.79) of stunted (measurement by HAZ) in schoolchildren (6,23,27). However, leiri *et al.* could not find a significant association between FI and HAZ, but did find a significant association between FI and lower WAZ (24). In agreement, two studies showed that schoolchildren with FI have a higher risk of low WAZ (OR, 3.0 to 4.8) compared to children without FI (30,34). Two studies found no significant association between FI and anthropometric indices (21,25).

Table I. Summary of studies that evaluated food insecurity and undernutrition

Population (authors, year)	Study design	FI evaluation	Malnutrition assessment	Outcomes
61 Brazilian children aged 5-10 years (Bueno <i>et al.</i> , 2021)	Cross-sectional	Brazilian Food Insecurity Scale (EBIA)	BMI-Z and HAZ Cut-off points WHO 2007	No significant association was found between nutritional status and familial FI
671 Ethiopian children aged 10.9 ± 2.67 years (Geletaw <i>et al.</i> , 2021)	Institution-based quantitative cross-sectional study	Household Food Insecurity Access Scale (HFIAS)	BMI-Z and HAZ Cut-off points WHO 2007	Children from food-insecure households were 2.79 times [AOR. 2.79; 95 % CI: 1.81, 4.31] more likely to be stunted than children from food-secure households
327 Philippine children aged 6-12 years (leiri <i>et al.</i> , 2021)	Community-based, cross-sectional study	Food Insecurity Experience Scale (FIES)	HAZ and WAZ Cut-off points WHO 2007	There was no association between the experience of household FI and HAZ Household income, household size, parental BMI, and the experience of household severe FI were significantly associated with lower WAZ in the adjusted model
167 Malaysian children aged 7-11 (Teh <i>et al.</i> , 2020)	Cross-sectional	Radimer/Cornell Hunger and Food Insecurity Instrument	BMI-Z and HAZ Cut-off points WHO 2007	There were no associations between the food security index and anthropometric indices
1730 Venezuelan children aged 7-12 years (Herrera-Cuenca <i>et al.</i> , 2019)	Cross-sectional	6-item short USDA Household Food Security Survey Module (HFSSM)	BMI-Z and HAZ Cut-off points WHO 2007	We found a significant association between FI and short height in children ($\chi^2 = 8.205$, 2 df, $p = 0.017$)
431 Nicaraguan children aged 3-11 years (Schmeer <i>et al.</i> , 2017)	Cross-sectional	Latin American and Caribbean Food Security Questionnaire (ELCSA)	HAZ Cut-off points Frisancho 2008	Children in mildly food insecure households had 34% higher odds of low height-for-age compared with children living in food secure households
450 Ethiopia children aged 7-14 years (Wolde <i>et al.</i> , 2015)	Cross-sectional survey	Food and Nutrition Technical Assistance (FANTA)	BMI-Z, HAZ and WAZ Cut-off points WHO 2007	Children live in food insecure households are more likely to be stunted, under-weight and wasted than children live in food secure households (AOR = 2.5; 95 % CI, 1.5-5.6; AOR = 3.9; 95 % CI, 1.2-12.0; AOR = 4.8; 95 % CI, 1.7-13.6)
2526 Colombian children aged 5-12 years (Isanaka <i>et al.</i> , 2007)	Cross-sectional survey	USDA Household Food Security Survey Module (HFSSM)	HAZ, WAZ and WHZ Cut-off points WHO 2007	FI children were almost 3 times more likely to be underweight compared with food-secure children ($p = 0.0007$)

FI: food insecurity; BMI-Z: body mass index for age Z-score; HAZ: height-for-age Z-score; WAZ: weight-for-age Z-score; WHZ: weight-for-height Z-score; WHO: World Health Organization.

FOOD INSECURITY AND OVERWEIGHT OR OBESITY

Nine articles evaluated the relationship between FI and overweight or obesity, the summary of results is shown in table II. Two studies conducted in Mexican schoolchildren found that higher levels of FI were associated with an increased risk of overweight and obesity (OR, 2.3) (28,33). However, Murillo-Castillo *et al.* and Shamah-Levy *et al.* found that higher

levels of FI were associated with lower prevalence of overweight and obesity in Mexican schoolchildren (26,29). Similarly, two studies conducted in schoolchildren in Brazil and Jamaica found the same negative association between FI and lower risk of overweight and obesity (OR, 0.78 and OR, 0.65, respectively) (31,32). Bethancourt *et al.* found that FI and hair cortisol concentration were associated with a lower BMI-Z, in turn, FI was associated with a higher percentage of body fat in Bolivian schoolchildren (22).

Table II. Summary of studies that evaluated food insecurity and overweight or obesity

Population (authors, year)	Study design	FI evaluation	Malnutrition assessment	Outcomes
61 Brazilian children aged 5-10 years (Bueno <i>et al.</i> , 2021)	Cross-sectional	Brazilian Food Insecurity Scale (EBIA)	BMI-Z and HAZ Cut-off points WHO 2007	No significant association was found between nutritional status and familial FI
167 Bolivian children aged 6-16 years (Bethancourt <i>et al.</i> , 2021)	Cross-sectional	Household Food Insecurity Access Scale (HFIAS)	BMI-Z Cut-off points WHO 2007	There was a significant positive association between HFIAS score and % body fat (Model 1 $\beta = 0.44$, SE = 0.22, $p = 0.04$) Although neither HFIAS nor hair cortisol concentration were independently associated with BMI-Z, they had a significant negative linear joint effect on BMI-Z ($\beta = -0.08$; SE = 0.04, $p = 0.04$)
167 Malaysian children aged 7-11 (Teh <i>et al.</i> , 2020)	Cross-sectional	Radimer/Cornell Hunger and Food Insecurity Instrument	BMI-Z and HAZ Cut-off points WHO 2007	There were no associations between the food security index and anthropometric indices
100 Mexican children aged 6-12 years (Murillo-Castillo <i>et al.</i> , 2020)	Cross-sectional	Mexican Food Safety Scale (EMSA)	BMI-Z Cut-off points WHO 2007	Adjusted analysis showed a negatively association between food insecurity and overweight and obesity, both in boys and girls (p for trend in logit < 0.01)
105 Mexican children aged 6-12 years (Rosas <i>et al.</i> , 2017)	Descriptive cross-sectional	Mexican Food Safety Scale (EMSA)	BMI-Z Cut-off points WHO 2007	49.4 % of households present FI; among these, 51.61 % of schoolchildren show obesity; 29.03 % overweight; 19.35 % risk of overweight, and 64.52 % cardiometabolic risk
7181 mother-schoolchild (5-11 years) pairs Mexicans (Shamah-Levy <i>et al.</i> , 2017)	Cross-sectional survey	Latin American and Caribbean Food Security Scale (ELCSA)	HAZ, WAZ and WHZ Cut-off points WHO 2007	The logistic regression model revealed a significantly lower prevalence of overweight and obesity among those with mild, moderate, or severe FI ($p < 0.05$)
782 Brazilian children aged 6.9 ± 0.6 years (Vicenzi <i>et al.</i> , 2015)	Cross-sectional school-based	Brazilian Food Insecurity Scale (EBIA)	BMI-Z Cut-off points WHO 2007	After controlling for potential confounders, children with food insecurity had 22 % lower odds of overweight
1674 Jamaican children aged 10-11 years (Dubois <i>et al.</i> , 2011)	Cross-sectional	2-items Hunger Vital Sign (HVS)	BMI-Z Overweight was defined using Cole's age- and sex-specific criteria	Children in FI households had odds of 0.65 (95 % CI: 0.4-0.9) for being overweight or obese in comparison to those in food-secure households
768 Mexican children aged 9-15 years (Ortiz-Hernández <i>et al.</i> , 2007)	Cross-sectional	2-2-items Hunger Vital Sign (HVS)	BMI-Z Cut-off points CDC 2000 references	Children with severe FI were 2.53 times more likely to be overweight than children without FI ($p = 0.002$)

FI: food insecurity; BMI-Z: body mass index for age Z-score; HAZ: height-for-age Z-score; WAZ: weight-for-age Z-score; WHZ: weight-for-height Z-score; WHO: World Health Organization.

DISCUSSION

LMICs are the most affected by FI and malnutrition. Children with FI and malnutrition have less physical, mental and social development in addition to more health problems (1,2,6,8). Knowing the relationship between FI and malnutrition, as well as its determinants, can allow the development of better strategies for its approach. However, evaluating the association between FI and malnutrition in schoolchildren is complicated because both problems are multifactorial.

There are many validated tools to assess FI and nutritional status, further complicating the association between FI and malnutrition. FI can be assessed at a household or personal level. In this review, nine different tools for assessing FI were identified, with eight at the household level and one at the personal level. Household-level surveys include a section to capture children's experience, however, when answered by the head of the household, they identify FI globally (35). Various authors have suggested that parents' perceptions of their children's FI could be imprecise or incomplete (36,37). This may be because not all members of the same household deal with FI in the same way (38). Parents may not be fully aware of their children's experiences or the actions they take to reduce the severity of FI (39). Research suggests that this discrepancy may be due to different ways of reasoning and response styles (36-39).

Studies evaluating the reliability of schoolchildren reporting their own experiences of FI concluded that those aged between 6 and 16 years are capable of doing so (37,40). In the USA, Connell et al. developed the Children Food Security Survey (CFSS), which adapted 9 questions from the HFSSM for use with children aged between 12 and 17 years (40). The CFSS demonstrated acceptable reliability and apparent internal validity (40). A study comparing the diet quality with FI levels, using both the HFSSM and CFSS for measurement, concluded that both surveys concurred in classifying FI as long as the child was ≥ 6 years old (41). However, the author suggested that the CFSS might be more sensitive in detecting differences in micronutrient intake (41).

The studies included in this review used the BMI-Z, HAZ, WAZ and WHZ indices to define malnutrition. Two of the studies included in this review did not find a significant association between FI and anthropometric indices (21,25). Nutritional status is normally defined by anthropometric indices, however, there are other tools that can aid in a more comprehensive evaluation. Nutritional status is defined as the physical condition resulting from the relationship between individual requirements and the consumption, absorption and use of energy and nutrients (1). Therefore, for a comprehensive nutritional assessment, one must consider not only anthropometric parameters, but also macro and micronutrient intake, as well as physical activity levels and the child's medical history. Some studies suggest that the lack of association between FI and malnutrition may be attributed to potential confounding variables, including the child's sex and age, household income level, parents' nutritional status, and ethnicity (42-44).

Of the eight articles in this review that evaluated the association between FI and undernutrition, six of them showed significant associations between higher FI and greater risk of stunted and undernutrition in general. Various theories have been proposed as possible causes of lower HAZ in schoolchildren with FI, such as delayed introduction of supplemental foods, or the introduction of poorly nutritious foods rich in starch and low in protein, vitamin A, iron and zinc (16). On the other hand, an analysis of eating habits in relation to FI levels showed that diet quality decreases with increasing FI. Children with FI have a higher intake of calories from saturated fats and added sugars, as well as a lower intake of fruits, vegetables, and dairy products (12-15,45). These eating patterns suggest an increased risk of obesity in children with FI.

Examining the intriguing relationship between FI and a reduced risk of excess weight, particularly overweight and obesity, reveals a noteworthy pattern in our reviewed studies. Five out of the nine studies investigating the FI-overweight/obesity link reported a lower prevalence of these conditions, as indicated by a lower BMI-Z, among children experiencing FI. This trend, observed predominantly in developing countries, contrasts with patterns in developed nations, where higher FI levels often correlate with increased prevalence of overweight and obesity. The variations observed across different socio-economic contexts, food availability, accessibility, and levels of physical activity might account for these disparities (42-44).

In the global issue of overweight and obesity, it is noteworthy that, while developed countries exhibit the highest rates of these conditions, LMICs have experienced a significant increase in their rates over the past decade (4). This phenomenon is intertwined with the complex relationship between FI and overweight or obesity, as revealed by nine studies analyzed in this review. Of these, five demonstrate that children facing FI have lower prevalences of overweight and obesity. However, these same studies agree that the quality of the diet is compromised, reflected in reduced consumption of fruits and vegetables, as well as increased intake of simple carbohydrates and saturated fats, which over time could contribute to the development of overweight or obesity (22,26,29,31,32).

It is important to note that the apparent lack of association between FI and a higher prevalence of overweight and obesity can be explained by the nutritional transition phase these countries are undergoing, where a dual burden of malnutrition is observed in the same household. One study highlights that FI is associated with stunting in preschoolers only if the mother does not suffer from obesity (29). The disparity in the relationship between FI and overweight or obesity between developed and developing LMICs may be due to divergent trajectories. While in developed countries, available social support services are associated with an increase in overweight/obesity, in developing countries, limited assistance in the form of food subsidies contributes to children from poorer families with FI consuming fewer total calories, thereby decreasing the chances of developing overweight or obesity (32,46,47).

However, it is crucial to note that the association between FI and excess body fat was highlighted in the study by Bethancourt et al., demonstrating a higher percentage of body fat in schoolchildren experiencing FI (22). Similarly, the studies

conducted by Rosas et al. and Ortiz-Hernández et al. found a higher risk of overweight and obesity among schoolchildren facing FI (28,33). The complexities of the relationship between FI and obesity are multifaceted and encompass factors such as the consumption of cheaper and more energy-dense foods, periods of insufficient food leading to overeating when available, and fluctuations in eating habits within families experiencing FI (48).

Schoolchildren living in vulnerable environments are at higher risk of experiencing a high prevalence of FI and malnutrition (2,3). Knowing the social determinants associated with FI and malnutrition will allow the creation of better approach strategies. As a strength, this review shows how FI may be related to malnutrition in schoolchildren (5 to 12 years) from LMICs. A limitation of this review is that the included studies had a cross-sectional design that precludes establishing causal relationships, so more research is recommended to explain how FI is related to malnutrition in schoolchildren through stronger epidemiological designs. Additional investigations may also be conducted to determine if there are other social determinants associated with these factors. These social determinants may include family structure, socioeconomic and educational level, access to resources, among others.

CONCLUSION

The present systematic review addressed the complex relationship between FI and malnutrition in schoolchildren from low- and middle-income countries. Analyzing 15 cross-sectional studies revealed significant variability in the findings. Concerning undernutrition, the majority of studies indicate an association between elevated levels of FI and an increased risk of low HAZ or low WAZ. However, discrepancies surfaced, with certain studies failing to establish a significant link between FI and anthropometric indices. Regarding overweight and obesity, the evidence presented conflicting results, with studies suggesting both positive and negative associations between FI and these health issues. The diverse outcomes underscore the necessity of considering contextual and population-specific factors when interpreting the relationship between FI and nutritional status.

These findings highlight the complexity of the relationship between FI and malnutrition in schoolchildren, suggesting that socioeconomic, cultural, and regional factors may significantly shape this association. The emphasis is placed on comprehensively addressing FI in public health policies, considering the various dimensions of nutritional status in school populations across diverse contexts.

Supplemental Table I. Search strategy in PubMed

Search number	Query	Results
12	#5 and #8 and #11	1,139
11	#9 or #10	14,520
10	Food insecur*[Title/Abstract] or Food secur*[Title/Abstract]	14,395
9	"Food Insecurity"[Mesh:NoExp] or "Food Security"[Mesh:NoExp]	1,254
8	#6 or #7	250,603
7	(((((("Poverty"[Mesh:NoExp]) OR "Child Poverty"[Mesh:NoExp]) OR "Poverty Areas"[Mesh:NoExp]) OR "Indigenous Peoples"[Mesh:NoExp]) OR "Rural Population"[Mesh:NoExp]) OR "Ethnic and Racial Minorities"[Mesh:NoExp]	113,585
6	developing countr*[Title/Abstract] OR developing nation*[Title/Abstract] OR developing population*[Title/Abstract] OR "developing world"[Title/Abstract] OR less developed countr*[Title/Abstract] OR "underdeveloped world"[Title/Abstract] OR middle income countr*[Title/Abstract] OR middle-income nation*[Title/Abstract] OR middle-income population*[Title/Abstract] OR low-income countr*[Title/Abstract] OR low-income nation*[Title/Abstract] OR low-income population*[Title/Abstract] OR lower income countr*[Title/Abstract] OR lower income nation*[Title/Abstract] OR lower income population*[Title/Abstract] OR underserved countr*[Title/Abstract] OR underserved nation*[Title/Abstract] OR underserved population*[Title/Abstract] OR underserved world[Title/Abstract] OR under served countr*[Title/Abstract]	148,889
5	#1 or #2 or #3 or #4	2,491,586
4	Elementary school[Title/Abstract]	8,346
3	school children[Title/Abstract]	24,911
2	Child*[Title/Abstract]	1,579,931
1	"Child"[Mesh:NoExp]	1,840,224

REFERENCES

- OMS. Malnutrición [Internet]. OMS. 2021 [cited 24 January 2022]. Available from: <https://www.who.int/es/news-room/fact-sheets/detail/malnutrition>
- FAO, FIDA, OMS, PMA, UNICEF. El estado de la seguridad alimentaria y la nutrición en el mundo 2022. Adaptación de las políticas alimentarias y agrícolas para hacer las dietas saludables más asequibles. Roma: FAO; 2022. 40 p.
- UNICEF. Estado mundial de la infancia 2019. Niños, alimentos y nutrición: Crecer bien en un mundo en transformación. Nueva York: UNICEF; 2019. 255 p.
- Bentham J, Di Cesare M, Bilano V, Bixby H, Zhou B, Stevens GA, et al. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million children, adolescents, and adults. *Lancet* 2017;390(10064):2627-42. DOI: 10.1016/S0140-6736(17)32129-3
- FAO. Seguridad Alimentaria y Nutricional Conceptos Básicos. 3ra ed. Roma, FAO.; 2011. 8 p.
- Schmeer KK, Piperata BA. Household food insecurity and child health. *Matern Child Nutr* 2017;13(2):1-13. DOI: 10.1111/mcn.12301
- Thomas MMC, Miller DP, Morrissey TW. Food Insecurity and Child Health. *Pediatrics* 2019;144(4):e20190397. DOI: 10.1542/peds.2019-0397
- Pai S, Bahadur K. The Impact of Food Insecurity on Child Health. *Pediatr Clin North Am* 2020;67(2):387-96. DOI: 10.1016/j.pcl.2019.12.004
- Morales-Ruán M del C, Humarán IMG, Shamah-Levy T, Valderrama-Álvarez Z, Melgar-Quirínez H. La inseguridad alimentaria está asociada con obesidad en mujeres adultas de México. *Salud Publica Mex* 2014;56(Supl. 1):54-61.
- Shamah-Levy T, Mundo-Rosas V, A Rivera-Dommarco J. La magnitud de la inseguridad alimentaria en México: su relación con el estado de nutrición y con factores socioeconómicos. *Salud Publica Mex* 2014;56(supl. 1):S79-85.
- Cuevas-Nasu L, Rivera-Dommarco JA, Shamah-Levy T, Mundo-Rosas V, Méndez-Gómez I. Inseguridad alimentaria y estado de nutrición en menores de cinco años de edad en México. *Salud Publica Mex* 2014;56:S47-53.
- Rodríguez LA, Mundo-Rosas V, Méndez-Gómez-Humarán I, Pérez-Escamilla R, Shamah-Levy T. Dietary quality and household food insecurity among Mexican children and adolescents. *Matern Child Nutr* 2017;13:e12372. DOI: 10.1111/mcn.12372
- Landry MJ, van den Berg AE, Asigbee FM, Vandyousefi S, Ghaddar R, Davis JN. Child-report of food insecurity is associated with diet quality in children. *Nutrients* 2019;11(7):1574. DOI: 10.3390/nu11071574
- Lee J, Kubik MY, Fulkerson JA. Diet Quality and Fruit, Vegetable, and Sugar-Sweetened Beverage Consumption by Household Food Insecurity among 8- to 12-Year-Old Children during Summer Months. *J Acad Nutr Diet* 2019;119(10):1695-702. DOI: 10.1016/j.jand.2019.03.004
- Tan ML, Laraia B, Madsen KA, Au LE, Frongillo EA, Ritchie LD. Child Food Insecurity Is Associated with Energy Intake among Fourth- and Fifth-Grade Girls. *J Acad Nutr Diet* 2018;119(10):1722-1731.e2. DOI: 10.1016/j.jand.2018.07.011
- Eicher-Miller HA, Zhao Y. Evidence for the age-specific relationship of food insecurity and key dietary outcomes among US children and adolescents. *Nutr Res Rev* 2018;31(1):98-113. DOI: 10.1017/S0954422417000245
- Manios Y, Androustos O, Katsarou C, Vampouli EA, Kulaga Z, Gurzkowska B, et al. Prevalence and sociodemographic correlates of overweight and obesity in a large Pan-European cohort of preschool children and their families: the ToyBox study. *Nutrition*. 2018;55-56:192-8. DOI: 10.1016/j.nut.2018.05.007
- Ogden CL, Carroll MD, Fakhouri TH, Hales CM, Fryar CD, Li X, et al. Prevalence of Obesity Among Youths by Household Income and Education Level of Head of Household - United States 2011-2014. *Morb Mortal Wkly Rep* 2018;67(6):186-9. DOI: 10.15585/mmwr.mm6706a3
- Spoede E, Corkins MR, Spear BA, Becker PJ, Gunnell Bellini S, Hoy MK, et al. Food Insecurity and Pediatric Malnutrition Related to Under- and Overweight in the United States: An Evidence Analysis Center Systematic Review. *J Acad Nutr Diet* 2021;121(5):952-78. DOI: 10.1016/j.jand.2020.03.009
- Lefebvre C, Glanville J, Briscoe S, Featherstone R, Littlewood A, Marshall C, et al. Chapter 4: Searching for and selecting studies. In: Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page M, et al., editors. *Cochrane Handbook for Systematic Reviews of Interventions* [Internet]. Cochrane; 2022. Available from: www.training.cochrane.org/handbook.
- Bueno MC, Franco JG, da Silva Leal GV, Kirsten VR. Food insecurity and the relationship with social, economic and nutritional factors in rural school students. *Cad Saúde Coletiva* 2021;29(2):153-62.
- Bethancourt HJ, Ulrich MA, Almeida DM, Rosinger AY. Household Food Insecurity, Hair Cortisol, and Adiposity Among Tsimane' Hunter-Forager-Horticulturists in Bolivia. *Obesity (Silver Spring)* 2021;29(6):1046-57. DOI: 10.1002/oby.23137
- Geletaw A, Egata G, Weldegebreal F, Kibr G, Semaw M. Nutritional Status and Associated Factors among Primary Schoolchildren from Pastoral Communities, Mieso-Mulu District, Sitti Zone, Somali Regional State, Eastern Ethiopia: Institution-Based Cross-Sectional Study. *J Nutr Metab* 2021;2021:6630620. DOI: 10.1155/2021/6630620
- leiri MCA, Kosaka S, Tomitsuka E, Umezaki M. Factors Affecting Undernutrition among School Children in Cebu, Philippines. *Ecol Food Nutr* 2020;60(2):182-97. DOI: 10.1080/03670244.2020.1813733
- Teh S, Asma A, Hamid Jan J, Yusof H. Assessment of Food Security, Anthropometric and Cognitive Function among Orang Asli Children in Pahang, Malaysia. *IMJM* 2020;19(3):81-91. DOI: 10.31436/imjm.v19i3.1669
- Murillo-Castillo KD, Frongillo EA, Corella-Madueño MA, Quizán-Plata T. Food Insecurity Was Associated with Lower Fruits and Vegetables Consumption but Not with Overweight and Obesity in Children from Mexican Fishing Communities. *Ecol Food Nutr* 2020;59(4):420-35. DOI: 10.1080/03670244.2020.1737042
- Herrera-Cuenca M, Rodríguez-Arroyo G, Vansant G, Roelants M, Soubry A, Macías-Tomei C, et al. Alcohol drinking, smoking and food insecurity in Venezuelan parents as potential determinants of their children's growth. *An venez nutr* 2019;32(1):13-25. DOI: 10.21203/rs.2.10784/v1
- Rosas T de J, Córdova AP, Villegas JG, Morales N. Evaluación de la Inseguridad Alimentaria y Nutrición de Escolares y sus Familias. *Rev Médica la Univ Veracruzana* 2017;17(1):7-22.
- Shamah-Levy T, Mundo-Rosas V, Morales-Ruan C, Cuevas-Nasu L, Méndez-Gómez-Humarán I, Pérez-Escamilla R. Food insecurity and maternal-child nutritional status in Mexico: cross-sectional analysis of the National Health and Nutrition Survey 2012. *BMJ Open* 2017;7:e014371. DOI: 10.1136/bmjopen-2016-014371
- Wolde M, Berhan Y, Chala A. Determinants of underweight, stunting and wasting among schoolchildren. *BMC Public Health* 2015;15(1):1-9. DOI: 10.1186/s12889-014-1337-2
- Vicenzi K, Henn RL, Weber AP, Backes V, Paniz VMV, Donatti T, et al. Food insecurity and overweight in first grade students in the municipal school system in São Leopoldo, Rio Grande do Sul State, Brazil. *Cad Saude Publica* 2015;31(5):1084-94. DOI: 10.1590/0102-311X00055914
- Dubois L, Francis D, Burnier D, Tatone-Tokuda F, Girard M, Gordon-Strachan G, et al. Household food insecurity and childhood overweight in Jamaica and Québec: a gender-based analysis. *BMC Public Health* 2011;11:199. DOI: 10.1186/1471-2458-11-199
- Ortiz-Hernández L, Acosta-Gutiérrez MN, Núñez-Pérez AE, Peralta-Fonseca N, Ruiz-Gómez Y. En escolares de la Ciudad de México la inseguridad alimentaria se asoció positivamente con el sobrepeso. *Rev Investig Clin* 2007;59(1):32-41.
- Isanaka S, Mora-Plazas M, Lopez-Arana S, Baylin A, Villamor E. Food insecurity is highly prevalent and predicts underweight but not overweight in adults and school children from Bogotá, Colombia. *J Nutr* 2007;137(12):2747-55. DOI: 10.1093/jn/137.12.2747
- Castell GS, Ngo de la Cruz J, Pérez C, Aranceta J. Escalas de evaluación de la inseguridad alimentaria en el hogar. *Rev Española Nutr Comunitaria* 2015;Supl. 1:270-6. DOI: 10.14642/RENC.2015.21.sup1.5074
- Escobar-Alegria JL, Frongillo EA, Fram MS, Pérez-Garay M, Macaudo MM, Billings DL. Parents are not fully knowledgeable of their children's experiences of food-insecurity. *FASEB J* 2012;26. DOI: 10.1096/asebj.26.1_supplement.28.3
- Fram SM, Frongillo EA, Draper CL, Fishbein EM. Development and Validation of a Child Report Assessment of Child Food Insecurity and Comparison to Parent Report Assessment. *J Hunger Environ Nutr* 2013;8(2):128-45. DOI: 10.1080/19320248.2013.790775
- Fram SM, Frongillo EA, Jones SJ, Williams RC, Burke MP, DeLoach KP, et al. Children Are Aware of Food Insecurity and Take Responsibility for Managing Food Resources. *J Nutr* 2011;141:1114-9. DOI: 10.3945/jn.110.135988
- Nalty CC, Sharkey JR, Dean WR. Children's reporting of food insecurity in predominately food insecure households in Texas border colonias. *Nutr J* 2013;12:15. DOI: 10.1186/1475-2891-12-15
- Connell CL, Nord M, Lofton KL, Yadrick K. Food security of older children can be assessed using a standardized survey instrument. *J Nutr* 2004;134(10):2566-72. DOI: 10.1093/jn/134.10.2566
- Jun S, Zeh MJ, Eicher-Miller HA, Bailey RL. Children's dietary quality and micronutrient adequacy by food security in the household and among household children. *Nutrients* 2019;11(5). DOI: 10.3390/nu11050965
- Jackson JA, Smit E, Branscum A, Gunter K, Harvey M, Manore MM, et al. The Family Home Environment, Food Insecurity, and Body Mass

- Index in Rural Children. *Heal Educ Behav* 2017;44(4):648-57. DOI: 10.1177/1090198116684757
43. Kaur J, Lamb MM, Ogden CL. The Association between Food Insecurity and Obesity in Children-The National Health and Nutrition Examination Survey. *J Acad Nutr Diet* 2015;115(5):751-8. DOI: 10.1016/j.jand.2015.01.003
 44. Kral TVE, Chittams J, Moore RH. Relationship between food insecurity, child weight status, and parent-reported child eating and snacking behaviors. *J Spec Pediatr Nurs* 2017;22(2):1-11. DOI: 10.1111/jspn.12177
 45. Fram SM, Ritchie LD, Rosen N, Frongillo EA. Child experience of food insecurity is associated with child diet and physical activity. *J Nutr* 2015;145(3):499-504. DOI: 10.3945/jn.114.194365
 46. Ruiz-Arranz M, Davis B, Stampini M, Winters P, Handa S. More calories or more diversity? An econometric evaluation of the impact of the PROGRESA and PROCAMPO transfer programs on food security in rural Mexico. 2002.
 47. de Bem Lignani J, Sichieri R, Burlandy L, Salles-Costa R. Changes in food consumption among the Programa Bolsa Familia participant families in Brazil. *Public Health Nutr* 2011;14(5):785-92. DOI: 10.1017/S136898001000279X
 48. Alaimo K, Olson CM, Frongillo J. Low Family Income and Food Insufficiency in Relation to Overweight in US Children: Is There a Paradox? *Arch Pediatr Adolesc Med* 2001;155(10):1161-7. DOI: 10.1001/archpedi.155.10.1161