



Trabajo Original

Paciente anciano

Association between the fat mass/fat-free mass ratio and muscle strength, static balance and exercise capacity in older adults: a cross-sectional study

Asociación entre la relación masa grasa/masa libre de grasa y la fuerza muscular, el equilibrio estático y la capacidad de ejercicio en adultos mayores: un estudio transversal

Walter Sepúlveda-Loyola^{1,5}, Yshoner Antonio Silva-Díaz², Mario Molari³, Erikson Alexander Jiménez Torres², Cintya Elisabeth Odar-Rojas², Denilson Teixeira⁴, Vanessa Suziane Probst⁴

¹Faculty of Health and Social Sciences. Universidad de Las Américas. Santiago, Chile. ²Grupo de Investigación en Terapia Física y Rehabilitación (TEFIRE). Instituto de Salud Integral Intercultural (ISI). Facultad de Ciencias de la Salud (FACISA). Universidad Nacional Toribio Rodríguez de Mendoza (UNTRM). Chachapoyas, Perú. ³Postgraduate Program in Physical Education. Universidade Estadual de Londrina (UEL) and Universidade Estadual de Maringá (UEM). Londrina/Maringá, Paraná. Brazil. ⁴Centro de Pesquisa e Pós Graduação em Saúde (CEPPoS). Centro de Ciências da Saúde (CCS). Universidade Estadual de Londrina (UEL). Londrina, Paraná. Brazil. ⁵Centro de Investigación en Ciencias Biológicas y Químicas. Universidad de Las Américas. Santiago, Chile

Abstract

Background: the FM/FFM ratio has been proposed as a new body composition variable. However, the association between FM/FFM ratio and physical function in older adults remains limited. Therefore, the aim of this study was to analyze the association between FM/FFM ratio and muscle strength, static balance, and exercise capacity in older adults.

Methods: this was a cross-sectional study involving 392 older adults from Londrina, Brazil. Subjects underwent body composition analysis by bioelectrical impedance, and assessments for aerobic capacity, muscle strength and static balance including the 6-minutes walking test, handgrip strength and one-legged stance test. Statistical analysis was conducted using correlation tests and lineal regression models adjusted for age and gender. The statistical significance considered was $p < 0.05$.

Results: a total of 392 older adults participated (mean age = 69.77 years; women: 74.7 %). The FM/FFM ratio was significantly associated with muscle strength ($\beta = -22.779$, 95 % CI: -26.741 to -18.818; $p < 0.001$), static balance ($\beta = -14.335$, 95 % CI: -19.980 to -8.690; $p < 0.001$), and exercise capacity ($\beta = -98.937$, 95 % CI: -152.286 to -45.588, $p < 0.001$). After adjusting for age and gender, FM/FFM ratio was an important predictor of muscle strength ($\beta = -4.687$, 95 % CI: -8.646 to -0.728; $p = 0.020$) and static balance ($\beta = -18.361$, 95 % CI: -24.943 to -11.778; $p < 0.001$).

Conclusions: the FM/FFM ratio is an important clinical measure of body composition that is significantly associated with key indicators of physical performance, including muscle strength, static balance, and exercise capacity in older adults.

Keywords:

Body composition.
Muscle strength. Electric impedance. Ageing.

Received: 12/11/2024 • Accepted: 29/03/2025

Funding: This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

Conflict of interest: The authors declare no conflict of interest.

Artificial intelligence: The authors declare not to have used artificial intelligence (AI) or any AI-assisted technologies in the elaboration of the article.

Sepúlveda-Loyola W, Silva-Díaz YA, Molari M, Jimenez Torres EA, Odar-Rojas CE, Teixeira D, Probst VS. Association between the fat mass/fat-free mass ratio and muscle strength, static balance and exercise capacity in older adults: a cross-sectional study. Nutr Hosp 2025;42(3):464-469
DOI: <http://dx.doi.org/10.20960/nh.05616>

Correspondence:

Walter Sepúlveda-Loyola. Faculty of Health and Social Sciences. Universidad de Las Américas. Echaurren, 140. Santiago Centro, Chile
e-mail: wsepulveda@udla.cl

Resumen

Antecedentes: la relación entre masa grasa (MG) y masa libre de grasa (MLG) (MG/MLG) se ha propuesto como nueva variable de la composición corporal. Sin embargo, la asociación entre MG/MLG y otras variables clínicas como la función física en los adultos mayores sigue siendo limitada. Por lo tanto, el objetivo de este estudio fue analizar la asociación entre el MG/MLG y la fuerza muscular, el equilibrio estático y la capacidad de ejercicio en adultos mayores.

Métodos: este fue un estudio transversal que incluyó a 392 adultos mayores de Londrina, Brasil. Se evaluaron la composición corporal mediante la impedancia bioeléctrica y la capacidad aeróbica, la fuerza muscular y el equilibrio estático mediante la prueba de la marcha de 6 minutos, la fuerza de prensión de la mano y la prueba del equilibrio en un solo pie. Se realizó un análisis estadístico utilizando pruebas de correlación y modelos de regresión lineal ajustados por edad y sexo. Se consideró un nivel de significancia estadística de $p < 0,05$.

Resultados: participaron un total de 392 adultos mayores (edad media = 69,77 años; mujeres: 74,7%). El MG/MLG se asoció significativamente con la fuerza muscular ($\beta = -22,779$, IC 95 %: -26,741 a -18,818; $p < 0,001$), el equilibrio estático ($\beta = -14,335$, IC 95 %: -19,980 a -8,690; $p < 0,001$) y la capacidad de ejercicio ($\beta = -98,937$, IC 95 %: -152,286 a -45,588, $p < 0,001$). Tras ajustar la edad y el sexo, MG/MLG se mantuvo como predictor importante de la fuerza muscular ($\beta = -4,687$, IC 95 %: -8,646 a -0,728; $p = 0,020$) y de equilibrio estático ($\beta = -18,361$, IC 95 %: -24,943 a -11,778; $p < 0,001$).

Conclusiones: MG/MLG es una medida clínica relevante de evaluación de la composición corporal que está asociada con indicadores clave del rendimiento físico, incluyendo fuerza muscular, equilibrio estático y capacidad de ejercicio en adultos mayores.

Palabras clave:

Composición corporal.
Fuerza muscular.
Impedancia eléctrica.
Envejecimiento.

INTRODUCTION

As time goes on, the older adult population experiences significant growth worldwide (1). This not only involves an interest in improving life expectancy, but also in addressing and reducing the disabilities that often accompany aging process (2). Longevity comes with a series of complex changes in the body composition, physical (3) and cognitive function (4,5). These changes can contribute to an increased risk of chronic diseases and geriatric syndromes such as sarcopenia or frailty affecting the quality of life and health (4,6).

Changes in body composition, particularly shifts in fat mass (FM) and fat-free mass (FFM), are critical indicators of metabolic health, physical function and overall well-being (6). In addition, FM and FFM are predictors of hospitalization and mortality in older adults (7). Accurate measurement of body composition parameters is essential, especially in populations vulnerable to muscle loss or increased fat mass, such as older adults with chronic diseases or geriatric syndromes (8,9). The relationship between FM and FFM, expressed as FM/FFM ratio has been proposed as new variable of body composition, which currently represents a crucial measurement unit for understanding overall health (6,10). Additionally, this relationship describes two aspects: metabolic load, which refers to the fat accumulated in the body, and metabolic capacity, which encompasses the body's ability to use energy and maintain proper physiological functions (11). In the elderly population, these values reflect a significant increase in FM, accompanied by a decrease in FFM, which have related to the presence of sarcopenic obesity (12).

Scientific evidence suggests that FM/FFM ratio is associated with metabolic and fatty liver clinical variables (12). Among the few studies conducted, the FM/FFM ratio has been related to conditions such as sarcopenia, chronic obstructive pulmonary disease (COPD), osteoporosis, and sarcopenic obesity (13). These comorbidities are often accompanied by reductions in grip strength, higher fracture incidence, increased fall risk, and

decreased exercise capacity (6,14). However, studies specifically examining the association between FM/FFM ratio and physical function in older adults remain limited (6). Despite existing evidence, further research is needed to clarify the clinical significance of FM/FFM ratio in relation to other functional and clinical variables in older adults, helping to establish its relevance for routine clinical assessments. Therefore, the aim of this study was to analyze the association between FM/FFM ratio and muscle strength, static balance, and exercise capacity in older adults.

METHODS

POPULATION STUDY

The present study is a cross-sectional analysis from the study on aging and longevity (ELLO data from 2009 to 2018) from the community of Londrina, Paraná, Brazil (15). The total sample of the EELO Project was 508 subjects, which is representative of the 43,610 citizens over 60 years of age who live in Londrina city. The inclusion and exclusion criteria and characteristics of the population were published previously. It is mentioned that adults aged 60 or older, who present a level 3 or 4 of physical independence (according to the Functional Status Scale), have a normal mental state (scoring more than 17 points on the Mini-Mental State Examination), and reside in an urban community of the city of Londrina, were included. Additionally, adults who exhibited severe dysfunction in the respiratory, neurological, or musculoskeletal systems, including amputations or the use of prosthetics were excluded (15). From the EELO program database, we included older adults of both sexes who have measured the body composition. The research was approved by the Research Ethics Committee of Universidade Norte do Paraná (UNOPAR) for both study designs (PP/0070/09 and PP 1,168,693). The collections and tests were performed at the UNOPAR. All participants signed an informed consent form.

BODY COMPOSITION

FM and FFM were measured using bioelectrical impedance of multi-frequency (Bio-dynamics 310TM; Biodynamics Corp., USA), after 10-hour of fasting. The assessment was performed on the dominant side of the individual with the patient in a supine position, the arms and legs were separated by approximately 30 degrees from a midline. The electrode placement sites were cleaned with alcohol. One electrode was placed in the hand and the other in the foot. FM/FFM ratio was estimated considering a previous study (16).

MUSCLE STRENGTH

Muscle strength was assessed via Hand Grip Strength (HGS) (17,18) assessed in a seated position with a Jamar hydraulic dynamometer (Sammons Preston Inc, Saint Paul, MN). The forearm and wrist were supported on the chair's armrests. Participants completed three trials with each hand, alternating between hands, with a 30-second rest between attempts. The highest value obtained was recorded for analysis.

STATIC BALANCE

Static balance was assessed by the One-Legged Support Test (OLST) (19). Participants stood on one leg on a stable surface with their eyes open. The best result from three attempts using the dominant leg was recorded for analysis and the maximum time of 30 seconds was considered for the test (20).

EXERCISE CAPACITY

Exercise capacity was assessed by the 6-Minute Walk Test (6MWT) (21,22). Participants walked along a 30-meter linear course, marked by two cones, for a duration of six minutes to cover the maximum possible distance. Standardized incentive phrases and information on the remaining time were provided throughout the test. Two assessments were conducted, with a 30-minute rest interval between them (23).

COMORBIDITIES

Charlson's comorbidity index was assessed to quantify the overall burden of comorbidities, which is an index with 19 medical conditions along with the corresponding weights. Comorbidities were weighted and scored using an algorithm proposed by Charlson et al. (24,25).

STATISTICAL ANALYSIS

Descriptive statistics were calculated for all variables, and data normality was assessed using the Kolmogorov-Smirnov test and the graphical procedures (normal probability plot). Continuous variables (age, body mass index, FM, etc.) were presented as

the mean and standard deviation (SD), and categorical variables (gender and prevalence of comorbidities) were presented as the number and percentage. Pearson's correlation coefficients were used to evaluate the linear relationships between the FM/FFM ratio and the dependent variables: muscle strength, static balance, and exercise capacity. To explore the associations between FM/FFM ratio and muscle strength, static balance, and exercise capacity, linear regression models were constructed. First, a crude model (unadjusted) was fitted to assess the univariate associations between the FM/FFM ratio (independent variable) and each outcome variable (HGS, OLST, and 6MWT). Subsequently, multivariable linear regression models were performed, adjusting for potential confounders, including age and gender. Regression coefficients (β) with corresponding 95 % confidence intervals (CIs) and p -values were reported to quantify the strength and significance of the associations. The statistical significance was considered as $p < 0.05$, and the analysis was performed using SPSS software (IBM Co., USA) and GraphPad Prism 6.0 (GraphPad Software, San Diego, CA, USA).

RESULTS

A total of 392 older adults were included in the study. The clinical characteristics of the individuals included in this study are reported in table I. The mean age of the participants was 69.77 ± 6.348 years and 74.7 % were female. The prevalence of diabetes was 63.8 %, arterial hypertension 63.7 %, and dyslipidemia 33.7 %. The average for FM/FFM ratio was 0.551 ± 0.169 .

Table I. Characteristics of the individuals

Variables	Total (n = 392)
Age (years)	69.77 \pm 6.348
Female, n (%)	293 (74.745%)
Comorbidities	
Diabetes	105 (26.786 %)
Arterial hypertension	250 (63.776 %)
Dyslipidemia	132 (33.673 %)
Charlson comorbidity index, score	2.1 \pm 1.7
Body composition	
BMI (kg/m ²)	27.789 \pm 4.665
FM (kg)	24.158 \pm 7.399
FFM (kg)	44.798 \pm 8.757
FM/FFM ratio	0.551 \pm 0.169
Muscle strength, static balance and exercise capacity	
HGS (kg)	26.143 \pm 7.748
OLST (sec)	12.789 \pm 9.899
6MWT (m)	523.23 \pm 91.330

Data are expressed as mean and standard deviation. BMI: body mass index; FM: fat mass; FFM: fat-free mass; HGS: hand grip strength; OLST: one-legged stance test; 6MWT: 6-minute walking test.

**CORRELATION AND REGRESSION ANALYSES
BETWEEN FM/FFM RATIO AND MUSCLE
STRENGTH, STATIC BALANCE AND EXERCISE
CAPACITY**

Pearson's correlation analysis revealed significant negative associations between the FM/FFM ratio and all three physical performance measures: muscle strength (HGS: $r = -0.498$; $p < 0.001$), static balance (OLST: $r = -0.244$; $p < 0.001$), and exercise capacity (6MWT: $r = -0.182$; $p < 0.001$). The correlation between the FM/FFM ratio and muscle strength, static balance and exercise capacity in older adults is presented in figure 1 (A, B and C, respectively).

In the non-adjusted linear regression analysis (Model 1), the FM/FFM ratio was significantly associated with muscle strength (HGS: $\beta = -22.779$, 95 % CI: -26.741 to -18.818, $p < 0.001$, $R^2 = 0.247$), static balance (OLST: $\beta = -14.335$, 95 % CI: -19.980 to -8.690, $p < 0.001$, $R^2 = 0.060$), and exercise capacity (6MWT: $\beta = -98.937$, 95 % CI: -152.286 to -45.588,

$p < 0.001$, $R^2 = 0.033$). After adjusting for age and gender in Model 2, FM/FFM ratio continues to be an important predictor of muscle strength (HGS: $\beta = -4.687$, 95 % CI: -8.646 to -0.728, $p = 0.020$, $R^2 = 0.539$) and static balance (OLST: $\beta = -18.361$, 95 % CI: -24.943 to -11.778, $p < 0.001$, $R^2 = 0.218$). However, the association with exercise capacity was no longer significant (6MWT: $\beta = -32.518$, 95 % CI: -97.965 to 32.929, $p = 0.329$, $R^2 = 0.109$). The associations between the FM/FFM ratio and the dependent variables: muscle strength, static balance, and exercise capacity were presented in table II. In addition, a regression analysis was performed to examine the association between gender and HGS, OLST, 6MWT, and the FM/FFM ratio in older adults. Older women exhibited lower HGS ($\beta = -11.4$, 95 % CI: -12.6 to -10.3, $p < 0.001$), OLST ($\beta = -1.6$, 95 % CI: -3.6 to -0.5, $p < 0.001$), and 6MWT performance ($\beta = -42.7$, 95 % CI: -60.8 to -24.5, $p < 0.001$), as well as a higher FM/FFM ratio ($\beta = 0.2$, 95 % CI: 0.19 to 0.25, $p < 0.001$) compared to older men. Similar results were observed in the model adjusted for age (Supplementary Table I).




Figure 1.
Correlation between FM/FFM ratio and muscle strength, static balance and exercise capacity in older adults.

Table II. Association between FM/FFM ratio and muscle strength, static balance and exercise capacity in older adults

Variable	Muscle strength (HGS)		Static balance (OLST)		Exercise capacity (6MWT)	
	Coef. β (CI: 95 %)	p	Coef. β (CI: 95 %)	p	Coef. β (CI: 95 %)	p
Model 1	-22.779 (-26.741 to -18.818)	< 0.001* $R^2 = 0.247$	-14.335 (-19.980 to -8.690)	< 0.001* $R^2 = 0.060$	-98.937 (-152.286 to -45.588)	< 0.001* $R^2 = 0.033$
Model 2	-4.687 (-8.646 to -0.728)	0.020* $R^2 = 0.539$	-18.361 (-24.943 to -11.778)	< 0.001* $R^2 = 0.218$	-32.518 (-97.965 to 32.929)	0.329 $R^2 = 0.109$

FM: fat mass; FFM: fat-free mass; HGS: hand grip strength; OLST: one-legged stance test; 6MWT: 6-minute walking test. Model 1: unadjusted analysis; Model 2: adjusted model by age and gender. Statistical significance, * $p < 0.05$. Analysis with 392 individuals.

Supplementary Table I. Association between gender and HGS, OLST, 6MWT and FM/FFM ratio in older adults

Variable	FM/FFM ratio		HGS		OLST		6MWT	
	Coef. β (CI: 95 %)	<i>p</i>	Coef. β (CI: 95 %)	<i>p</i>	Coef. β (CI: 95 %)	<i>p</i>	Coef. β (CI: 95 %)	<i>p</i>
Male	Ref.	-	Ref.	-	Ref.	-	Ref.	-
Female Model 1	0.22 (0.19 to 0.25)	< 0.001	-11.4 (-12.6 to 10.3)	< 0.001	-1.6 (3.6 to 0.5)	< 0.001	-42.7 (60.8 to 24.5)	< 0.001
Female Model 2	0.22 (0.20 to 0.25)	< 0.001	-11.9 (-13.1 to 10.9)	< 0.001	-2.7 (4.5 to 0.8)	0.005	-48.9 (66.7 to 31.2)	< 0.001

FM: fat mass; FFM: fat-free mass; HGS: hand grip strength; OLST: one-legged stance test; 6MWT: 6-minute walking test; Ref: reference. Model 1: unadjusted analysis; Model 2: adjusted model by age. Statistical significance, **p* < 0.05. Analysis with 392 individuals.

DISCUSSION

The present study demonstrates a significant association between the FM/FFM ratio and muscle strength, static balance, and exercise capacity in older adults. These findings underscore the importance of assessing body composition and suggest that a higher FM/FFM ratio is strongly associated with reduced physical function in older adults.

The negative correlation between FM/FFM ratio with muscle strength and balance is particularly noteworthy, as both muscle variables are well-established predictor of frailty, disability, and overall quality of life in aging populations (26). We observed that higher FM/FFM ratio was strongly associated with lower muscle strength and poorer static balance, even after adjusting for age and gender. This suggests that the detrimental effects of a higher FM/FFM ratio on these physical performance measures are not merely a function of age or gender differences but may reflect underlying physiological mechanisms such as sarcopenic obesity (27). Although the FM/FFM ratio itself is not a diagnostic criterion for sarcopenic obesity, this index may serve as a proxy, as low fat-free mass commonly associated with sarcopenia can occur alongside either normal or elevated fat mass, indicative of sarcopenic obesity (28). Sarcopenic obesity, defined by the simultaneous presence of increased fat mass and decreased muscle mass, is widely implicated in reduced physical performance and heightened risk of falls and fractures in older adults (29). The observed association between a higher FM/FFM ratio and poorer static balance further supports the idea that elevated body fat, particularly when coupled with lower lean mass, may impair the neuromuscular control needed to maintain stability. This aligns with previous studies that emphasize the adverse impact of increased adiposity on physical function in aging populations (30).

The relationship between body composition and exercise capacity is more complex and influenced by additional factors, which were reported in this and in a current study (31). For this reason, the association between FM/FFM ratio and exercise capacity was no longer significant after adjusting for age and gender in this study. Previous research has also suggested that exercise capacity may be more sensitive to other markers of physical fitness, rather than body composition alone (32).

Our findings underscore the potential value of incorporating the FM/FFM ratio into clinical assessments as an integrated measure of body composition that can provide additional insights into the physical health of older adults. Future studies should explore the mechanisms underlying these associations and investigate the effects of targeted exercise and nutritional interventions on FM/FFM ratio and physical performance. Furthermore, reference values for FM/FFM ratio have been proposed for populations outside of Latin America (33) highlighting the need for studies to establish region-specific reference values and assess their association with clinical outcomes in diverse populations.

Finally, this study has several limitations that should be considered. First, its cross-sectional design limits the findings to an exploratory perspective, precluding conclusions about causality. Second, we did not include additional measurements of physical function, such as the sit-to-stand test or the timed up and go test, commonly used in clinical practice with older adults. However, the study also has notable strengths, including a large sample size and the use of standardized protocols for measurements which has been extensively investigated in older adult populations worldwide, recommended by international guidelines and validated in Brazilian population (17-19,21,22). Future research with a longitudinal design, incorporating other key functional variables and examining the predictive capacity of FM/FFM ratio to identify risks of falls, fractures, hospitalizations, or mortality in older adults, is warranted.

CONCLUSION

In conclusion, FM/FFM ratio is a crucial clinical measure of body composition that is significantly associated with key indicators of physical performance, including muscle strength, static balance, and exercise capacity in older adults. A higher FM/FFM ratio is linked to decreased muscle strength and poorer static balance, even after adjusting for confounders, highlighting the importance of body composition in maintaining functional capacity during aging. These results emphasize the need to incorporate the FM/FFM ratio into daily clinical practice, which will allow for the implementation of personalized programs to reduce the risks of falls and loss of mobility, through exercises focused on increasing muscle mass and improving balance.

REFERENCES

- World Health Organization. Ageing and health [Internet] 2024 [cited 2024 Oct 8]. Available from: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>
- Kakita D, Harada K, Kurita S, Morikawa M, Nishijima C, Fujii K, et al. Impact of fat to muscle ratio with risk of disability on community-dwelling Japanese older adults: A 5-year longitudinal study. *Arch Gerontol Geriatr* 2024;126:105524. DOI: 10.1016/j.archger.2024.105524
- Wannagat W, Nieding G, Tibken C. Age-related decline of metacognitive comprehension monitoring in adults aged 50 and older: Effects of cognitive abilities and educational attainment. *Cogn Dev* 2024;70:101440. DOI: 10.1016/j.cogdev.2024.101440
- Spinedi M, Clark C, Zullo L, Kerkisiek A, Pistis G, Castelao E, et al. Cholesterol-metabolism, plant sterols, and long-term cognitive decline in older people – Effects of sex and APOEε4. *iScience* 2024;27(2):109013. DOI: 10.1016/j.isci.2024.109013
- Wang ZT, Shen XN, Ma YH, Ou YN, Dong PQ, Tan PL, et al. Associations of the Rate of Change in Geriatric Depression Scale with Amyloid and Cerebral Glucose Metabolism in Cognitively Normal Older Adults: A Longitudinal Study. *J Affect Disord* 2021;280:77-84. DOI: 10.1016/j.jad.2020.10.078
- Merchant RA, Seetharaman S, Au L, Wong MWK, Wong BLL, Tan LF, et al. Relationship of Fat Mass Index and Fat Free Mass Index With Body Mass Index and Association With Function, Cognition and Sarcopenia in Pre-Frail Older Adults. *Front Endocrinol (Lausanne)* 2021;12:765415. DOI: 10.3389/fendo.2021.765415
- Travassos A, Rodrigues A, Furlanetto KC, Donária L, Bisca GW, Nellesen AG, et al. Fat-free mass depletion in patients with COPD in Brazil: Development of a new cutoff point and its relation with mortality and extrapulmonary manifestations. *Eur J Clin Nutr* 2014;71(11):1285-90. DOI: 10.1038/ejcn.2017.105
- Sepúlveda-loyola W, Sergio P, Probst S. Mecanismos fisiopatológicos de la sarcopenia en la EPOC. *Revista chilena de enfermedades respiratorias* 2019;35(2). DOI: 10.4067/S0717-73482019000200124
- Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, et al. Sarcopenia: Revised European consensus on definition and diagnosis. *Age Ageing* 2019;48(1):16-31. DOI: 10.1093/ageing/afy169
- He X, Li Z, Tang X, Zhang L, Wang L, He Y, et al. Age- and sex-related differences in body composition in healthy subjects aged 18 to 82 years. *Medicine* 2018;97(25):e11152. DOI: 10.1097/MD.0000000000001152
- Rugila DF, Oliveira JM, Machado FVC, Correia NS, Puzzi VC, Passos NFP, et al. Fat mass to fat-free mass ratio and its associations with clinical characteristics in asthma. *Heart & Lung* 2022;56:154-60. DOI: 10.1016/j.hrtlung.2022.07.006
- Marcelin G, Silveira ALM, Martins LB, Ferreira AVM, Clément K. Deciphering the cellular interplays underlying obesity-induced adipose tissue fibrosis. *Journal of Clinical Investigation* 2019;129(10):4032-40. DOI: 10.1172/JCI129192
- Vijewardane SC, Balasuriya A, Johnstone AM, Myint PK. Impact of age on the prevalence of poor-quality dietary variety, associated lifestyle factors, and body composition profile (low body muscle mass and high body fat mass) in older people residing in Colombo district, Sri Lanka. *Heliyon* 2024;10(5):e27064. DOI: 10.1016/j.heliyon.2024.e27064
- Ishimoto T, Hisamatsu K, Fujimoto T, Matsudaira N, Yamamoto N, Hayashi H, et al. Association between adductor pollicis muscle thickness and low skeletal muscle mass index in community-dwelling older women undergoing outpatient rehabilitation. *Clin Nutr ESPEN* 2024;60:116-21. DOI: 10.1016/j.clnesp.2024.01.016
- Molari M, Fernandes KBP, Marquez A de S, Probst VS, Bignardi PR, Teixeira D de C. Impact of physical and functional fitness on mortality from all causes of physically independent older adults. *Arch Gerontol Geriatr* 2021;97:104524. DOI: 10.1016/j.archger.2021.104524
- Low S, Goh KS, Ng TP, Ang SF, Moh A, Wang J, et al. The prevalence of sarcopenic obesity and its association with cognitive performance in type 2 diabetes in Singapore. *Clinical Nutrition* 2020;39(7):2274-81. DOI: 10.1016/j.clnu.2019.10.019
- Mendes J, Amaral TF, Borges N, Santos A, Padrão P, Moreira P, et al. Handgrip strength values of Portuguese older adults: a population based study. *BMC Geriatr* 2017;17(1):191. DOI: 10.1186/s12877-017-0590-5
- Garcia PA, Sampaio RX, de Moura JA, de Souza PF, da Costa LB, dos Santos Mendes FA. What is the most appropriate handgrip strength testing protocol for sarcopenia screening in older adults with cognitive impairment? *Braz J Phys Ther* 2024;28(4):101104. DOI: 10.1016/j.bjpt.2024.101104
- Pereira C, da Silva RA, de Oliveira MR, Souza RDN, Borges RJ, Vieira ER. Effect of body mass index and fat mass on balance force platform measurements during a one-legged stance in older adults. *Aging Clin Exp Res [Internet]* 2018 [cited 2025 Feb 24];30(5):441-7. Available from: <https://link.springer.com/article/10.1007/s40520-017-0796-6>
- Sepúlveda W, Mella K, Araya-Quintanilla F, Barros J, Molari M, Suziane V. Asociación entre medidas clínicas para el diagnóstico de osteosarcopenia con funcionalidad y mortalidad en adultos mayores: estudio longitudinal. *Nutrición Clínica y Dietética Hospitalaria* 2022;42(3). DOI: 10.12873/423sepulveda
- Frade MC, dos Reis IM, Basso-Vanelli RP, Brandão AF, Jamami M. Reproducibility and Validity of the 6-Minute Stationary Walk Test Associated With Virtual Reality in Subjects With COPD. *Respir Care* 2019;64(4):425-33. DOI: 10.4187/respcare.06237
- Negreiros A, Padula RS, Andrea Bretas Bernardes R, de Moraes MV, Pires RS, Chiavegato LD. Predictive validity analysis of six reference equations for the 6-minute walk test in healthy Brazilian men: a cross-sectional study. *Braz J Phys Ther* 2017;21(5):350-6. DOI: 10.1016/j.bjpt.2017.06.003
- Britto RR, Probst VS, Andrade AFD de, Samora GAR, Hernandez NA, Marinho PEM, et al. Reference equations for the six-minute walk distance based on a Brazilian multicenter study. *Braz J Phys Ther* 2013;17(6):556-63. DOI: 10.1590/S1413-35552012005000122
- Charlson M, Szatrowski TP, Peterson J, Gold J. Validation of a combined comorbidity index. *J Clin Epidemiol* 1994;47(11):1245-51. DOI: 10.1016/0895-4356(94)90129-5
- Charlson ME, Pompei P, Ales KL, Mackenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *J Chronic Dis* 1987;40(5):373-83. DOI: 10.1016/0021-9681(87)90171-8
- Roubenoff R. Origins and Clinical Relevance of Sarcopenia. *Canadian Journal of Applied Physiology* 2001;26(1):78-89. DOI: 10.1139/h01-006
- Tomažič A, Žvanut B, Grbac LV, Jurdana M. Identification of sarcopenic obesity in adults undergoing orthopaedic surgery: Relationship between “a body shape index” (ABSI) and fat-free mass. A cross-sectional study. *PLoS One* 2022;17(6):e0269956. DOI: 10.1371/journal.pone.0269956
- Siervo M, Rubele S, Shannon OM, Prado CM, Donini LM, Zamboni M, et al. Prevalence of sarcopenic obesity and association with metabolic syndrome in an adult Iranian cohort: The Fasa PERSIAN cohort study. *Clin Obes* 2021;11(4). DOI: 10.1111/cob.12459
- Follis S, Cook A, Bea JW, Going SB, Laddu D, Cauley JA, et al. Association Between Sarcopenic Obesity and Falls in a Multiethnic Cohort of Postmenopausal Women. *J Am Geriatr Soc* 2018;66(12):2314-20. DOI: 10.1111/jgs.15613
- Kim S, Leng XI, Kritchevsky SB. Body Composition and Physical Function in Older Adults with Various Comorbidities. *Innov Aging* 2017;1(1). DOI: 10.1093/geroni/igx008
- Gültekin M, Yeğinoğlu G, Bingöl H. The relationship between the exercise capacity and somatotype components, body composition, and quadriceps strength in individuals with coronary artery disease. *Journal of Clinical Medicine of Kazakhstan* 2021;18(6):62-8. DOI: 10.23950/jcmk/11345
- Myers J, McAuley P, Lavie CJ, Despres JP, Arena R, Kokkinos P. Physical Activity and Cardiorespiratory Fitness as Major Markers of Cardiovascular Risk: Their Independent and Interwoven Importance to Health Status. *Prog Cardiovasc Dis* 2015;57(4):306-14. DOI: 10.1016/j.pcad.2014.09.011
- Xiao J, Purcell SA, Prado CM, Gonzalez MC. Fat mass to fat-free mass ratio reference values from NHANES III using bioelectrical impedance analysis. *Clinical Nutrition* 2018;37(6):2284-7. DOI: 10.1016/j.clnu.2017.09.021