

**OR 1052**

**Normative reference values for hand grip dynamometry in Spain. Association with lean mass**

*Valores de normalidad de dinamometría de mano en España. Relación con la masa magra*

Francisco José Sánchez Torralvo<sup>1</sup>, Nuria Porras<sup>1</sup>, José Abuín Fernández<sup>1</sup>, Francisca García Torres<sup>1</sup>, María José Tapia<sup>1</sup>, Fuensanta Lima<sup>1</sup>, Federico Soriguer<sup>1</sup>, Montserrat Gonzalo<sup>1</sup>, Gemma Rojo Martínez<sup>1,2</sup> y Gabriel Olveira<sup>1,2</sup>

<sup>1</sup>Unidad de Gestión Clínica de Endocrinología y Nutrición. IBIMA. Hospital Regional Universitario de Málaga-Universidad de Málaga. Málaga. <sup>2</sup>CIBERDEM - CIBER de Diabetes y Enfermedades Metabólicas Asociadas (CB07/08/0019). Instituto de Salud Carlos III. Madrid

**Recibido:** 19/02/2017

**Aceptado:** 09/10/2017

**Correspondencia:** Gabriel Olveira Fuster. Unidad de Gestión Clínica de Endocrinología y Nutrición. Hospital Civil (Complejo del Hospital Regional Universitario de Málaga). Plaza del Hospital Civil, s/n. Pabellón 1, sótano. 29009 Málaga  
e-mail: gabrielm.olveira.sspa@juntadeandalucia.es

**DOI:** 10.20960/nh.1052

**ABSTRACT**

**Background and objectives:** The objective of this study was to establish reference values for hand grip strength, compare the results obtained with Collin and Jamar type dynamometers and determine their association with anthropometric and lean mass measurements.

**Material and methods:** This cross-sectional population-based study was undertaken in Pizarra (Málaga, Spain). The grip strength of the dominant hand was measured using

Collin and Jamar dynamometers. Skinfolds (triceps, abdominal, biceps of dominant arm and subscapular) were measured, and body composition was estimated. Eight hundred seventeen adults randomly selected from the census were recruited. Dynamometry reference values are presented for the dominant hand, by gender and age groups.

**Results:** No determinations could be made with the Collin dynamometer in 69 women due to the difficulty in grasping the dynamometer. We found significant positive correlations between the measurements with Jamar and Collin dynamometers ( $r = 0.782$ ;  $p < 0.001$ ) and between grip strength and lean mass index (LMI), determined by both dynamometers ( $r = 0.538$ ,  $p < 0.001$  and  $r = 0.462$ ,  $p < 0.001$ , respectively). Malnourished patients according to LMI had significantly lower grip strength than normally nourished patients ( $p < 0.001$  for Jamar;  $p < 0.02$  for Collin).

**Conclusions:** Dynamometry reference values in the Spanish population are presented. We recommend the use of the Jamar type dynamometer *versus* the Collin type dynamometer. Hand grip dynamometry is associated with lean mass, which confirms its usefulness in nutritional assessment.

**Key words:** Hand strength. Reference values. Hand grip. Dynamometry. Nutritional status.

## RESUMEN

**Antecedentes y objetivos:** no existen valores de normalidad en España con el dinamómetro Jamar. El objetivo fue determinar valores de normalidad de fuerza muscular, comparar los resultados obtenidos con los dinamómetros tipo Collin y tipo Jamar entre sí, y determinar su asociación con medidas antropométricas y de masa magra.

**Material y métodos:** estudio transversal de base poblacional en Pizarra (Málaga). Se determinó la fuerza de presión de la mano dominante mediante dinamómetros Collin y Jamar. Se midieron los pliegues cutáneos (tricipital, abdominal, bicipital del brazo dominante y subescapular) y se estimó la composición corporal. Se reclutaron 817 adultos seleccionados aleatoriamente del censo. Se presentan valores de referencia de dinamometría para la mano dominante, por género y grupos de edad.

**Resultados:** no se pudieron realizar determinaciones con el dinamómetro Collin en 69 mujeres debido a la dificultad para agarrar el dinamómetro. Encontramos correlaciones positivas significativas entre las medidas de los dinamómetros Jamar y Collin ( $r = 0,782$ ;  $p < 0,001$ ) y entre la fuerza muscular determinada mediante ambos dinamómetros y el índice de masa magra (IMM) ( $r = 0,538$ ,  $p < 0,001$  y  $r = 0,462$ ,  $p < 0,001$ , respectivamente). Los pacientes desnutridos según IMM presentaron una fuerza muscular significativamente menor a la de los pacientes normonutridos ( $p < 0,001$  para Jamar y  $p < 0,02$  para Collin).

**Conclusiones:** se presentan valores de referencia de dinamometría en población española. Recomendamos el uso del dinamómetro tipo Jamar frente al dinamómetro tipo Collin. La dinamometría de mano se asocia con la masa magra, lo que avala su utilidad en la valoración nutricional.

## INTRODUCTION

There is a great variety of techniques to evaluate the nutritional status of a patient, although there is no single parameter available. Malnutrition implies a decrease in muscle mass, which is reflected in poorer performance on functional tests and alterations in body composition (1-3). This decrease in muscle strength appears before changes in anthropometric measurements and laboratory parameters are observed. Accordingly, the measurement of muscle strength may be a useful tool in screening and assessing malnutrition (4). The American Society for Parenteral and Enteral Nutrition has included the assessment of grip strength by dynamometer as one of the six criteria to define malnutrition (5), encouraging the use of cut-off points in each population by age and gender (6).

There are numerous clinical studies in hospitalized individuals or outpatients (surgical, elderly, oncological, etc.) that demonstrate that decreased grip strength, measured by hand dynamometry, is associated with increased stays, mortality and complications (3,7,8). Similarly, in epidemiological studies performed in different age groups, decreased grip strength is also associated with higher mortality and impaired functionality (3). Some body composition measures, like lean mass, seem to have a close relationship with hand strength as well as with physical function (9).

The hand dynamometer evaluates the isometric force of the hand and the forearm providing a quick, easy to use, and inexpensive method to assess the grip strength and, thus, the nutritional status of patients (10). Currently, the Jamar dynamometer is the most commonly used in clinical practice (4,11-13). Over the last few years, dynamometry reference values have been published in different countries, usually for the Jamar-type dynamometer (4,11,14,15), including a meta-analysis (16).

In Spain, the normative data are from another type of dynamometer (17-19) with no studies using the Jamar dynamometer. Likewise, there are no data comparing the results of the two different types of dynamometers, so it remains unknown if the measurements of both dynamometers are comparable in clinical practice. There are only a few studies relating dynamometry results to body composition (15,20). The objective of this study was to contribute reference values in a sample of subjects belonging to the general population in Spain using the Jamar dynamometer, to compare them with the Collin dynamometer, and to assess the relationship between dynamometry and anthropometric parameters, especially lean mass.

## **MATERIAL AND METHODS**

Our sample comprised a total of 817 healthy adults recruited from the population-based Pizarra study (Malaga) (21). These individuals, aged between 18 and 65 years, were randomly selected from the municipal census. Individuals who were institutionalized, pregnant, or who had severe physical or psychiatric conditions were excluded as were morbidly obese subjects (body mass index [BMI] > 40 kg/m<sup>2</sup>). All subjects gave their written consent, and the study was approved by the Ethics and Clinical Research Committee of the Regional University Hospital of Malaga.

Anthropometric measurements included weight, obtained with a scale adjusted to 0.1 kg (SECA 665, Seca, Germany), and height, using a stadiometer adjusted to 0.01 m (Holtain Ltd., Crosswell, UK), to calculate BMI. Skinfold thicknesses (subscapular, triceps, biceps, abdominal) were measured at standard sites (22) by a single investigator using a plicometer (Holtain Ltd., Crosswell, UK) with the precision of 0.2 mm. Three measurements were taken and the mean was calculated. Arm circumference was measured using a flexible measuring tape to the nearest 0.1 cm at the midpoint of the arm. Lean mass and fat mass percentages were estimated using the Durnin (23) and

Siri (24) equations. Hand dynamometry was performed using a Collin dynamometer (Medizintechnik AS, Germany) and a Jamar dynamometer (Asimow Engineering Co., Los Angeles, CA).

In the case of the Jamar dynamometer, the subjects were instructed to adjust the device so that the grip would be comfortable for their hand to obtain the best performance (25), although most chose to use the second position (3.8 cm). The subjects were instructed to squeeze the dynamometer with the maximum force they could apply after receiving a verbal command (26).

The measurements were taken with the patients sitting in a straight back chair with both feet on the ground, shoulders close to the body in a neutral position, and the elbow flexed at 90° without rotation (27).

Three measurements were obtained in the dominant hand with a rest period of at least one minute between trial (28). There was a minimum ten minute break between Collin and Jamar measurements, in that order.

The mean was calculated and the highest value was used to represent hand grip strength.

### **Statistical analysis**

Data analysis was performed using SPSS Statistics software v22. Descriptive data are shown as means and standard deviations. The Kolmogorov-Smirnov test was used to establish whether the variables followed a normal distribution.

In the hypothesis testing for continuous variables between groups, the Student's t test was used in the variables that followed a normal distribution and a non-parametric test (Mann-Whitney) was used for variables that were not normally distributed. We rejected the null hypothesis with an alpha of 0.05 for two tails. The degree of association between hand dynamometry, BMI and body composition measurements was analyzed using the Pearson's correlation coefficient.

### **RESULTS**

A total of 817 adults, 364 men and 453 women, were studied (Table I).

The measurements were valid for the Jamar dynamometer in all subjects studied, but only in 748 cases for the Collin dynamometer (364 men and 384 women), due to the

difficulty for grasping the dynamometer. A total of 69 women could not apply the necessary force. Their mean age was  $54.6 (\pm 9.2)$  years, significantly higher ( $p < 0.001$ ) than in the group of women who were able to perform the test ( $47.8 \pm 9.2$  years). Tables II and III show the grip strength values for the two dynamometers used, grouped by gender and distributed by age, together with their corresponding percentiles.

A high correlation was found between the data obtained with the Jamar and Collin dynamometers ( $r = 0.782$ ;  $p < 0.001$ ) (Fig. 1). Mean strength was higher in men than in women with both instruments and in all age groups ( $p < 0.001$ ). There was a tendency towards a negative correlation between age and grip strength with the Jamar dynamometer ( $r = -0.67$ ;  $p = 0.58$ ), which was significant with the Collin dynamometer ( $r = -0.143$ ;  $p < 0.001$ ). This significant correlation was observed with the Collin dynamometer in the subjects between the ages of 45 and 60 ( $r = -0.1$ ;  $p = 0.04$ ), and those above 60 years ( $r = -0.22$ ;  $p = 0.02$ ), which was not the case in those under age 45 ( $r = 0.02$ ;  $p = 0.74$ ). With the Jamar type dynamometer, this correlation was significant in the group of subjects under age 45 ( $r = 0.12$ ;  $p = 0.04$ ), in those between the ages of 45 and 60 ( $r = -0.12$ ;  $p = 0.023$ ) and in subjects over 60 years of age ( $r = -0.2$ ;  $p = 0.02$ ).

A positive correlation was found between grip strength using the Jamar dynamometer and BMI ( $r = 0.086$ ;  $p = 0.014$ ). This relationship increased when subjects were classified as normal weight ( $r = 0.268$ ;  $p < 0.001$ ) or overweight ( $r = 0.146$ ;  $p = 0.006$ ). No association was found between BMI and grip strength in obese patients.

The values obtained using the Jamar dynamometer showed a positive correlation with weight ( $r = 0.514$ ;  $p < 0.001$ ), height ( $r = 0.714$ ;  $p < 0.001$ ), and arm circumference ( $r = 0.249$ ;  $p < 0.001$ ), and a negative correlation with fat mass in kg: ( $r = -0.597$ ;  $p < 0.001$ ), triceps skinfold ( $r = -0.497$ ;  $p < 0.001$ ), biceps skinfold ( $r = -0.404$ ;  $p < 0.001$ ) and subscapular skinfold ( $r = -0.209$ ;  $p < 0.001$ ). There was also a significantly positive correlation between muscle strength and lean mass in kg ( $r = 0.774$ ;  $p < 0.001$ ) and lean mass index (LMI) ( $r = 0.538$ ;  $p < 0.001$ ) (Figs. 2 and 3).

Mean values obtained using the Collin dynamometer showed a positive correlation with weight ( $r = 0.434$ ;  $p < 0.001$ ), height ( $r = 0.663$ ;  $p < 0.001$ ) and arm circumference ( $r = 0.206$ ;  $p < 0.001$ ), and a negative correlation with fat mass (kg): ( $r = -0.569$ ;  $p <$

0.001), triceps skinfold ( $r = -0.518$ ;  $p < 0.001$ ), biceps skinfold ( $r = -0.404$ ;  $p < 0.001$ ) and subscapular skinfold ( $r = -0.214$ ;  $p < 0.001$ ). There was also a significantly positive correlation between grip strength and lean mass (kg) ( $r = 0.683$ ;  $p < 0.001$ ) and LMI ( $r = 0.462$ ;  $p < 0.001$ ).

Stratifying our population by European Society of Clinical Nutrition and Metabolism criteria (LMI of  $17 \text{ kg/m}^2$  for men and  $15 \text{ kg/m}^2$  for women [29]), patients considered as malnourished due to low lean mass had significantly lower mean grip strength than normally nourished patients, both with the Jamar dynamometer and with the Collin dynamometer (Table IV).

## DISCUSSION

In this study, we present normative reference values for the Spanish population using a Jamar hand dynamometer, a dynamometer for which there were no previous references in Spain, providing cut-off points to define malnutrition. Our results are similar to those of other studies that have published reference values for the Jamar-type dynamometer in Caucasian populations (30,14,20).

In the total sample, the cut-off points to define malnutrition (5<sup>th</sup> percentile) were 29 kg in men and 14 kg in women for the Jamar dynamometer, although this varied depending on age (Table III). These values are similar to those of other studies (15,20) and may be related to poorer functionality, as some authors suggest (31,32).

Similar to other studies (4,20,27), our values were significantly higher in men than in women for all age groups and negative correlations with age were observed, finding a decrease in grip strength (4). This has been attributed to age-related sarcopenia, since the weight loss that occurs with aging is primarily due to the loss of lean mass, thus leading to loss of grip strength (17). In our sample, no significant differences between the groups of individuals under 45 years and between 45 and 60 years were found, possibly because age-related sarcopenia appears especially in subjects aged 60 and older (4).

As in other studies, a positive association was found between grip strength and anthropometric measures such as weight, BMI, arm circumference, and height (33,11), although the association is weak (14).

A close correlation between dynamometry values and lean mass was also observed. Patients who met criteria for malnutrition according to LMI (29) presented lower hand grip strength. Since malnutrition due to low lean mass is primarily associated with increased morbidity and mortality related to malnutrition (29), dynamometry becomes a measure that provides a clear added value to nutritional assessment.

In subjects with acute or chronic disease, numerous factors may influence decreased muscle strength, including immobilization, decreased intake, inflammation, oxidative stress, electrolyte disturbances, use of drugs (corticosteroids, muscle relaxants, etc.). In this respect, grip strength is an excellent marker of functionality as, unlike weight, it can discriminate between malnourished individuals and those who are simply underweight and share the same BMI (10).

Furthermore, nutritional intervention studies have demonstrated significant improvement in muscle strength in the short and medium term, which also supports hand grip usefulness in patient follow-up (3,34).

We found good correlations between grip strength measured with both dynamometers. Although the Collin dynamometer is significantly more economical, we believe its use may be less suitable than that of the Jamar dynamometer because some people (especially older women) have some difficulty in correctly grasping the device and applying force, so the measurement may not be valid. This may be of special relevance in the hospital setting, where the mean inpatients age is high.

### **Strengths and weaknesses**

Our work was carried out in the context of a population-based epidemiological study with an adequate sample size. Moreover, we measured anthropometric parameters enabling us to relate muscle strength to body composition, especially to lean mass.

Nonetheless, this was a cross-sectional study in which other parameters of functionality were not evaluated, nor was the long-term effect on morbidity and mortality verified. Measurements were only taken in the dominant hand, although some studies suggest that dominance does not affect grip strength (15,33,35). Finally, there was a large percentage of subjects with obesity, which could have partially conditioned the results.

## CONCLUSIONS

We present reference values for hand dynamometry using a Jamar hand dynamometer for a Spanish population, providing cut-off points to define malnutrition. We recommend using the Jamar dynamometer as opposed to the Collin dynamometer in clinical practice. Hand dynamometry is associated with lean mass, which supports its usefulness in nutritional assessment.

## ACKNOWLEDGMENTS

We would like to thank the town of Pizarra and all those who participated in the study.

## REFERENCES

1. Hornby ST, Nunes QM, Hillman TE, Stanga Z, Neal KR, Rowlands BJ, et al. Relationships between structural and functional measures of nutritional status in a normally nourished population. *Clin Nutr* 2005;24(3):421-6.
2. Norman K, Schütz T, Kemps M, Lübke HJ, Lochs H, Pirlich M. The Subjective Global Assessment reliably identifies malnutrition-related muscle dysfunction. *Clin Nutr* 2005;24(1):143-50.
3. Norman K, Stobäus N, González MC, Schulzke J-D, Pirlich M, Symreng T, et al. Hand grip strength: Outcome predictor and marker of nutritional status. *Clin Nutr* 2011;30(2):135-42.
4. Budziareck MB, Pureza Duarte RR, Barbosa-Silva MCG. Reference values and determinants for handgrip strength in healthy subjects. *Clin Nutr* 2008;27(3):357-62.
5. White JV, Guenter P, Jensen G, Malone A, Schofield M, Academy of Nutrition and Dietetics Malnutrition Work Group, et al. Consensus statement of the Academy of Nutrition and Dietetics/American Society for Parenteral and Enteral Nutrition: Characteristics recommended for the identification and documentation of adult malnutrition (undernutrition). *J Acad Nutr Diet* 2012;112(5):730-8.
6. Malone A, Hamilton C. The Academy of Nutrition and Dietetics/The American Society for Parenteral and Enteral Nutrition Consensus Malnutrition Characteristics: Application in Practice. *Nutr Clin Pract* 2013;28(6):639-50.
7. Gale CR, Martyn CN, Cooper C, Sayer AA. Grip strength, body composition, and mortality. *Int J Epidemiol* 2007;36(1):228-35.

8. Rantanen T, Volpato S, Ferrucci L, Heikkinen E, Fried LP, Guralnik JM. Handgrip strength and cause-specific and total mortality in older disabled women: Exploring the mechanism. *J Am Geriatr Soc* 2003;51(5):636-41.
9. Araujo AB, Chiu GR, Kupelian V, Hall SA, Williams RE, Clark R V, et al. Lean mass, muscle strength, and physical function in a diverse population of men: A population-based cross-sectional study. *BMC Public Health* 2010;10(1):508.
10. Vaz M, Thangam S, Prabhu A, Shetty PS. Maximal voluntary contraction as a functional indicator of adult chronic undernutrition. *Br J Nutr* 1996;76:9-15.
11. Mitsionis G, Pakos EE, Stafilas KS, Paschos N, Papakostas T, Beris AE. Normative data on hand grip strength in a Greek adult population. *Int Orthop* 2009;33(3):713-7.
12. Schlüssel MM, Dos Anjos LA, De Vasconcellos MTL, Kac G. Reference values of handgrip dynamometry of healthy adults: A population-based study. *Clin Nutr* 2008;27(4):601-7.
13. Werle S, Goldhahn J, Drerup S, Simmen BR, Sprott H, Herren DB. Age- and gender-specific normative data of grip and pinch strength in a healthy adult Swiss population. *J Hand Surg Eur Vol* 2009;34(1):76-84.
14. Massy-Westropp NM, Gill TK, Taylor AW, Bohannon RW, Hill CL. Hand grip strength: Age and gender stratified normative data in a population-based study. *BMC Res Notes* 2011;4:127.
15. Peters MJH, Van Nes SI, Vanhoutte EK, Bakkers M, Van Doorn PA, Merkies ISJ, et al. Revised normative values for grip strength with the Jamar dynamometer. *J Peripher Nerv Syst* 2011;16(1):47-50.
16. Bohannon RW, Peolsson A, Massy-Westropp N, Desrosiers J, Bear-Lehman J, Rothstein JM, et al. Reference values for adult grip strength measured with a Jamar dynamometer: A descriptive meta-analysis. *Physiotherapy* 2006;92(1):11-5.
17. Lázaro MLM, Penacho Lázaro MA, Losantos FB, Plaza Bayo A. Nuevas tablas de fuerza de la mano para población adulta de Teruel. *Nutr Hosp* 2008;23(1):35-40.
18. Luna-Heredia E, Martín-Peña G, Ruiz-Galiana J. Handgrip dynamometry in healthy adults. *Clin Nutr* 2005;24(2):250-8.
19. Marrodán Serrano MD, Romero Collazos JF, Moreno Romero S, Mesa Santurino MS, Cabañas Armesilla MD, Pacheco Del Cerro JL, et al. Handgrip strength in children and teenagers aged from 6 to 18 years: Reference values and relationship with size

and body composition. *An Pediatr* 2009;70(4):340-8.

20. Dodds RM, Syddall HE, Cooper R, Benzeval M, Deary IJ, Dennison EM, et al. Grip strength across the life course: Normative data from twelve British studies. *PLoS One* 2014;9(12):e113637.

21. Soriguer F, Gutiérrez-Repiso C, Rubio-Martín E, García-Fuentes E, Almaraz MC, Colomo N, et al. Metabolically healthy but obese, a matter of time? Findings from the prospective Pizarra study. *J Clin Endocrinol Metab* 2013;98(6):2318-25.

22. Olveira G. Manual de nutrición clínica y dietética. 3<sup>rd</sup> ed. Olveira G, ed. Madrid: Díaz de Santos; 2007. p. 188-9.

23. Durnin JV, Womersley J. Body fat assessed from total body density and its estimation from skinfold thickness: Measurements on 481 men and women aged from 16 to 72 years. *Br J Nutr* 1974;32(1):77-97.

24. Siri WE. Body composition from fluid spaces and density: Analysis of methods. 1961. *Nutrition* 1993;9(5):480-92.

25. Boadella JM, Kuijjer PP, Sluiter JK, Frings-Dresen MH. Effect of self-selected handgrip position on maximal handgrip strength. *Arch Phys Med Rehabil* 2005;86(2):328-31.

26. Johansson CA, Kent BE, Shepard KF. Relationship between verbal command volume and magnitude of muscle contraction. *Phys Ther* 1983;63(8):1260-5.

27. Mathiowetz V, Kashman N, Volland G, Weber K, Dowe M, Rogers S. Grip and pinch strength: Normative data for adults. *Arch Phys Med Rehabil* 1985;66(2):69-74.

28. Watanabe T, Owashi K, Kanauchi Y, Mura N, Takahara M, Ogino T. The short-term reliability of grip strength measurement and the effects of posture and grip span. *J Hand Surg Am* 2005;30(3):603-9.

29. Cederholm T, Bosaeus I, Barazzoni R, Bauer J, Van Gossum A, Klek S, et al. Diagnostic criteria for malnutrition - An ESPEN Consensus Statement. *Clin Nutr* 2015;34(3):335-40.

30. Mendes J, Alves P, Amaral TF. Comparison of nutritional status assessment parameters in predicting length of hospital stay in cancer patients. *Clin Nutr* 2014;33(3):466-70.

31. Lauretani F, Russo CR, Bandinelli S, Bartali B, Cavazzini C, Di Iorio A, et al. Age-associated changes in skeletal muscles and their effect on mobility: An operational

diagnosis of sarcopenia. J Appl Physiol 2003;95(5):1851-60.

32. Sallinen J, Stenholm S, Rantanen T, Heliövaara M, Sainio P, Koskinen S. Hand-grip strength cut points to screen older persons at risk for mobility limitation. J Am Geriatr Soc 2010;58(9):1721-6.

33. Günther CM, Bürger A, Rickert M, Crispin A, Schulz CU. Grip strength in healthy Caucasian adults: Reference values. J Hand Surg Am 2008;33(4):558-65.

34. Oliveira G, Oliveira C, Doña E, Palenque FJ, Porras N, Dorado A, et al. Oral supplement enriched in HMB combined with pulmonary rehabilitation improves body composition and health related quality of life in patients with bronchiectasis (Prospective, Randomised Study). Clin Nutr 2016;35(5):1015-22.

35. Bohannon RW. Grip strength: A summary of studies comparing dominant and nondominant limb measurements. Percept Mot Skills 2003;96(3 Pt 1):728-30.

**Nutrición  
Hospitalaria**

**Table I. General and anthropometric characteristics of the study population**

Variables	Total	Men	Women
	n = 817	n = 364	n = 453
	Mean ± SD	Mean ± SD	Mean ± SD
BMI (kg/m <sup>2</sup> )	28.8 (± 4.4)	29.3 (± 4.1)	28.4 (± 4.6)
Age (years)	49.7 (± 9.6)	50.9 (± 9.59)	48.8 (± 9.6)
Weight (kg)	76.3 (± 13.9)	84.2 (± 12.7)	70 (± 11.5)
Height (cm)	162.7 (± 4.4)	169.7 (± 7.1)	157.1 (± 5.9)
Arm circumference (cm)	30.2 (± 3.5)	31.1 (± 3.3)	29.6 (± 3.5)
Triceps skinfold (mm)	16.8 (± 7.5)	11.3 (± 5)	21.1(± 6.3)
Biceps skinfold (mm)	11.4 (± 5.8)	8.4 (± 3.9)	13.8 (± 5.9)
Subscapular skinfold (mm)	23.1 (± 7.3)	20.1 (± 6.9)	24.8 (± 7.2)
Abdominal skinfold (mm)	25.8 (± 8.2)	26.1 (± 8.5)	25.5 (± 7.9)
Arm muscle circumference (cm)	24.9 (± 3.7)	27.6 (± 3.2)	22.9 (± 2.7)
Fat mass percentage	34.1 (± 6.5)	29.1 (± 5.2)	38.1 (± 4.4)
Lean mass (kg)	50 (± 10.4)	59.1 (± 7.5)	42.6 (± 5.5)
Lean mass index (kg/m <sup>2</sup> )	18.7 (± 2.7)	20.5 (± 2.3)	17.2 (± 2)

n = number. BMI: Body mass index.

**Table II. Strength of the dominant hand by gender and age, measured with a Collin dynamometer**

Age group (years)	Grip strength (kg)									
	Mean $\pm$ SD	Maximum $\pm$ SD	P5	P10	P25	P50	P75	P90	P95	
<b>Men</b>										
Total n = 364	31.4 $\pm$ 13.6	34 $\pm$ 14	8	15	25	35	43.8	52	56	
Under 45 years n = 125	35.8 $\pm$ 12.6	38.3 $\pm$ 12.9	15	20.2	33	40	45	55	56.7	
From 45 to 60 years n = 164	32.2 $\pm$ 13	35 $\pm$ 13.4	11.3	15.5	26.3	36	43	50.5	55.8	
Over 60 years n = 71	22 $\pm$ 12.3	24.4 $\pm$ 12.9	6.6	8	16	24	30	45	50.8	
<b>Women</b>										
Total n = 384	10.3 $\pm$ 7.4	12 $\pm$ 7.6	2	3	6	11	17	22	25	
Under 45 years n = 161	12.6 $\pm$ 7	14.3 $\pm$ 7	3	5	8	15	20	24	26.9	
From 45 to 60 years n = 181	8.6 $\pm$ 7.3	10.5 $\pm$ 7.8	1.1	2	4.5	9	15	20	23.9	
Over 60 years n = 38	8.4 $\pm$ 7	10 $\pm$ 7.2	1	1.9	3	9.5	16.3	20.1	23.2	

SD: Standard deviation.

**Table III. Strength of the dominant hand by gender and age, measured with a Jamar dynamometer**


Age group (years)	Grip strength (kg)								
	Mean $\pm$ SD	Maximum $\pm$ SD	P5	P10	P25	P50	P75	P90	P95
<b>Men</b>									
Total n = 364	45.7 $\pm$ 9.9	47.8 $\pm$ 10.3	30	34	40	48	54	62	64.8
Under 45 years n = 125	47.2 $\pm$ 10	49.5 $\pm$ 10.4	32.6	37.6	42	48	57.5	64	64.7
From 45 to 60 years n = 164	47.2 $\pm$ 9.2	49.5 $\pm$ 9.5	34.5	37.5	44	50	55.8	62	66
Over 60 years n = 71	39.5 $\pm$ 9.3	40.9 $\pm$ 9.6	26.6	29.2	34	40	47	54	58.2
<b>Women</b>									
Total n = 453	24.2 $\pm$ 6.2	26 $\pm$ 6.3	16	18	22	26	30	34	36
Under 45 years n = 175	24.7 $\pm$ 5.4	26.4 $\pm$ 5.4	18	20	23	26	30	33.4	36.4
From 45 to 60 years n = 216	24.7 $\pm$ 6.6	26.4 $\pm$ 6.7	15	18	22	26	30	34	38
Over 60 years n = 58	21.3 $\pm$ 6.4	22.5 $\pm$ 6.7	12.8	14	18	22	28	31.1	34

SD: Standard deviation.

**Table IV. Hand grip strength for both dynamometers according to lean mass index as malnutrition criteria**

	Low LMI	Normal LMI	p value
Total			
Collin n = 649	14.3 ± 9 (n = 52)	21.1 ± 15.4 (n = 597)	0.002
Jamar n =709	26.6 ± 8.3 (n = 57)	34.6 ± 13.7 (n = 652)	< 0.001

LMI: Lean mass index. Low LMI interpreted as LMI < 17 kg/m<sup>2</sup> in men and LMI < 15 kg/m<sup>2</sup> in women.






Fig. 1. Relationship between hand grip strength measured by Collin dynamometer and Jamar dynamometer.




Fig. 2. Relationship between hand grip strength (Jamar) and lean mass.




Fig. 3. Relationship between hand grip strength (Jamar) and lean mass index.

**Nutricion  
Hospitalaria**